

International Conference and Exhibition on

Nanomedicine and Drug Delivery

May 29-31, 2017 Osaka, Japan

Exposure to excess phenobarbital negatively influences the osteogenesis of chick embryos

Yu Yan Jinan University, China

Phenobarbital is an antiepileptic drug that is widely used to treat epilepsy in a clinical setting. However, a long term of phenobarbital administration in pregnant women may produce side effects on embryonic skeletogenesis. In this study, we aim to investigate the mechanism by which phenobarbital treatment induces developmental defects in long bones. We first determined that phenobarbital treatment decreased chondrogenesis and inhibited the proliferation of chondrocytes in chick embryos. Phenobarbital treatment also suppressed mineralization in both *in vivo* and *in vitro* long bone models. Next, we established that phenobarbital treatment delayed blood vessel invasion in a cartilage template, and this finding was supported by the down-regulation of vascular endothelial growth factor in the hypertrophic zone following phenobarbital treatment. Phenobarbital treatment inhibited tube formation and the migration of human umbilical vein endothelial cells. In addition, it impaired angiogenesis in chick yolk sac membrane model and chorioallantoic membrane model. In summary, phenobarbital exposure led to shortened lengths of long bones during embryogenesis, which might result from inhibiting mesenchyme differentiation, chondrocyte proliferation, and delaying mineralization by impairing vascular invasion.

Biography

Yu Yan has completed her Bachelor's from South Medicine University of China. Currently, she is pursuing her Master's majoring in Histology and Anatomy at Medicine School of Jinan University. She has published 3 papers in reputed journals in 2016.

283329554@gg.com

Notes: