

## Stimulation of keratinocyte wound healing responses and re-epithelialization by novel epoxy-tiglanes via protein kinase C activation

Rachael L Moses<sup>1</sup>, Glen M Boyle<sup>2</sup>, Paul Reddell<sup>3</sup>, Robert Steadman<sup>1</sup> and Ryan Moseley<sup>1</sup>

<sup>1</sup>Cardiff University, UK

<sup>2</sup>QIMR Berghofer Medical Research Institute, Australia

<sup>3</sup>QBiotics Ltd., Australia

**N**ovel epoxy-tiglanes, EBC-46 and EBC-211, are sourced from seeds of the Fountain's Blushwood Tree, indigenous to Queensland. EBC-46 possess potent tumouricidal properties, through classical PKC activation, and is under development by our industrial partner, QBiotics Ltd., as a human and veterinary anti-cancer pharmaceutical. In clinical studies, EBC-46 also stimulated exceptional dermal healing, manifested as accelerated wound re-epithelialisation, closure and minimal scarring. This work describes epoxy-tigliane effects on keratinocyte wound healing responses and their underlying mechanisms of action.

Immortalized human epidermal keratinocytes (HaCaTs) were treated with EBC-46 or EBC-211 (0-10µg/ml). Cell cycle progression/proliferation were assessed by FACS analysis and MTT assay. HaCaT migration was assessed using in vitro scratch wound assays. Global gene expression changes induced by epoxy-tiglanes were quantified by Microarray analysis, with differentially expressed genes confirmed by protein level analysis. As epoxy-tiglanes mediate responses via classical protein kinase (PKC) activation, mechanistic studies were performed with BIM-1 (pan-PKC), Gö6976 (classical-PKC) and LY317615 (PKC-βI/PKC-βII) inhibitors. Western blotting confirmed phospho-PKC activation following epoxy-tigliane treatment.


Both epoxy-tiglanes induced significant HaCaT cell cycle progression and proliferation; and also promoted significant HaCaT scratch wound closure. Microarray analyses identified key genes differentially expressed in EBC-46/EBC-211-treated HaCaTs, which contribute to their stimulatory effects on keratinocyte proliferation and migration. Enhanced proliferative and migratory responses were significantly abrogated by BIM-1 and Gö6976, although LY317615 exhibited minimal inhibitory effects. PKC activation increased following epoxy-tigliane treatment.

Such findings explain the enhanced re-epithelialization responses in epoxy-tigliane-treated skin; and provide justification for their translational development as novel therapeutics for impaired wound re-epithelialisation.

### Speaker Biography

Rachael L Moses completed her PhD at Cardiff University in 2016 and is currently a Postdoctoral Research Associate at Cardiff University, UK. Dr. Moses's research focuses on elucidating the mechanisms underlying the novel epoxy-tigliane pharmaceuticals exceptional dermal wound healing responses. Dr. Moses was awarded a travel bursary to visit a medical research institute in Queensland, Australia, to undertake Microarray Analysis determining key genotypic changes following epoxy-tigliane treatment. Dr Moses has filed patents with an industrial partner in this sector (QBiotics Ltd.); and has been awarded conference prizes relating to this area.

e: MosesR@cardiff.ac.uk

 Notes: