

3rd International Conference on **Earth Science & Climate Change**

July 28-30, 2014 DoubleTree by Hilton Hotel San Francisco Airport, USA

Effect of seasonal changes on abundance and heavy metal accumulation in the marine polychaete *Nereis succinea*

Abd Allah A T¹, Moustafa M A², El Sheimy N³ and Said R²

¹Jazan University, KSA

²Al-Azhar University, Egypt

³Assiut University, Egypt

Effect of seasonal changes on uptake of dissolved ions of Pb, Cu, Cr, Zn, Mn and Fe as related to chemical and physical water criteria was investigated in the marine polychaete *Nereis succinea*. Studied species was collected from nonpolluted site; the marine biological station (MBS) and polluted site; the fish port station (FPS) at Hurghada, Red sea. The seasonal means of ecological factors were significantly differed between the two studied sites; pH ($P<0.05$), dissolved oxygen ($P<0.05$) turbidity ($P<0.01$) and conductivity ($P<0.01$). The abundance of *Nereis succinea* exhibited highly significant difference ($P<0.01$) between MBS and FBS sites. LSD multicomparison revealed high significant difference ($P<0.01$) between the means of *Nereis* abundance at all seasons except in winter and autumn between the two sites and winter and spring in MBS only. MANOVA test revealed that the seasonal means of concentration of all studied metals in *Nereis* tissue, exhibited high significance ($P<0.01$) at both sites, except in the case of lead. MANOVA showed that the BAFs of studied metals may be depending on the environmental factors as well as season and site variations. All BAFs at FBS gave the significant level ($P<0.01$), while at MBS the same level of significance was observed for Pb, Cd and Cu and ($P<0.05$) for Fe.

abd_allaht@hotmail.com