

# INFECTION CONTROL AND CLINICAL MICROBIOLOGY

September 25-26, 2017 Chicago, USA

## *Staphylococcus pseudintermedius* adenosine synthase inhibits the innate immune response and promotes bacterial survival in canine blood

Mohamed A Abouelkhair<sup>1,2</sup>, David A Bemis<sup>1</sup> and Stephen A Kania<sup>1</sup>

<sup>1</sup>University of Tennessee, USA

<sup>2</sup>University of Sadat City, Egypt

*Staphylococcus pseudintermedius* is an important opportunistic bacterial pathogen that is the most common cause of canine pyoderma. It is frequently associated with urinary tract, wound and surgical site infections and occasionally causes zoonotic infections in human beings. The development of a staphylococcal vaccine is challenging and prior infection with *S. pseudintermedius* is not associated with protective immunity. Identification of a novel virulence factor inhibiting phagocytosis and evasion of innate immunity could play an important role in the prevention or treatment of *S. pseudintermedius* infection. Here, through bioinformatics-based analysis of *S. pseudintermedius* genome sequences, we identified a putative adenosine synthase gene (*SpdsA*) encoding a 5'-nucleotidase. *S. pseudintermedius* *SpdsA* protein shares approximately 73.46 % similarity with that of *Staphylococcus aureus* and 46.44% similarity with that of *S. suis* type2. Like the orthologous protein in *S. aureus* it catalyzes the dephosphorylation of adenosine mono- and triphosphates and consequently produces the immune signaling molecule adenosine. Attenuation of this enzyme with selected amino acid substations result in diminished hydrolytic activity on adenosine mono- and triphosphates and low adenosine production. Adenosine perturbation enable escape of *S. pseudintermedius* from phagocytic clearance in dog blood. In contrast, the addition of *SpdsA* inhibitor or A2A receptor antagonist to phagocytic cells result in diminished ability of *S. pseudintermedius* to escape from phagocytic killing. Taken together, these results indicate that *SpdsA* may play an important role in promoting *S. pseudintermedius* survival and in inhibiting neutrophil activity by adenosine synthesis.

### Biography

Mohamed A Abouelkhair has completed his Master's degree from University of Sadat City, Egypt. He is an Assistant Lecturer in Bacteriology, Mycology and Immunology department, University of Sadat city, Egypt. Currently, he is a Graduate PhD student in Comparative and Experimental Medicine Graduate program, college of veterinary medicine, University of Tennessee, USA.

mabouelk@vols.utk.edu

### Notes: