

31ST ANNUAL WORLD DENTISTRY SUMMIT

August 14-16, 2017 | Toronto, Canada

Subcutaneous emphysema during Oral Treatment

Zamir Rahman Mayar
SNI Oral Care, Afghanistan

Extraction of third molars is the most widely recognized surgical technique performed in oral surgery once a day and, in spite of surgical abilities and aptitude, entanglements may happen. Difficulties saw amid or after third molar evacuation may incorporate torment, swelling, dying, contamination, sinus aperture and nerve harm. Luckily, with an appropriate administration and a decent surgical strategy, the rate of such occasions is low. Subcutaneous emphysema related with dental extraction happens when the air from the rapid dental handpiece is constrained into the delicate tissue through the reflected fold and attacks the neighboring tissues, prompting swelling, crepitus on palpation and every so often spreading through the tissue spaces of the

fascial planes. Albeit uncommon, iatrogenic subcutaneous emphysema can have genuine and conceivably hazardous results. Care ought to be taken when utilizing air-driven handpieces. The entrance of air into the facial tissues is not restricted to tooth extractions, but rather may likewise happen through different entries of passage, for example, endodontically treated teeth, periodontium and cuts of intraoral delicate tissues. At the point when subcutaneous emphysema happens, it must be immediately analyzed and legitimately figured out how to decrease the danger of further entanglements.

e: zamirrahman@hotmail.com