

4TH ANNUAL DIABETES CONGRESS



5th World Congress on

NURSING AND HEALTHCARE

April 15-16, 2019 | Milan, Italy

Therapeutic ketosis and the broad field of applications for the ketogenic diet: Ketone ester applications & clinical updates

Raffaele Pilla

St. John of God Hospital – Fatebenefratelli, Italy

It has been recently shown that nutritional ketosis is effective against seizure disorders and various acute/chronic neurological disorders. Physiologically, glucose is the primary metabolic fuel for cells. However, many neurodegenerative disorders have been associated with impaired glucose transport/metabolism and with mitochondrial dysfunction, such as Alzheimer's/Parkinson's disease, general seizure disorders and traumatic brain injury. Ketone bodies and tricarboxylic acid cycle intermediates represent alternative fuels for the brain and can bypass the rate-limiting steps associated with impaired neuronal glucose metabolism. Therefore, therapeutic ketosis can be considered as a metabolic therapy by providing alternative energy substrates. It has been estimated that the brain derives over 60% of its total energy from ketones when glucose availability is limited. In fact, after prolonged periods of fasting or ketogenic diet (KD), the body utilizes energy obtained from free fatty acids (FFAs) released from adipose tissue. Because the brain is unable to derive significant energy from FFAs, hepatic ketogenesis

converts FFAs into ketone bodies-hydroxybutyrate (BHB) and acetoacetate (AcAc)-while a percentage of AcAc spontaneously decarboxylates to acetone. Large quantities of ketone bodies accumulate in the blood through this mechanism. This represents a state of normal physiological ketosis and can be therapeutic. Ketone bodies are transported across the blood-brain barrier by monocarboxylic acid transporters to fuel brain function. Starvation or nutritional ketosis is an essential survival mechanism that ensures metabolic flexibility during prolonged fasting or lack of carbohydrate ingestion. Therapeutic ketosis leads to metabolic adaptations that may improve brain metabolism, restore mitochondrial ATP production, decrease reactive oxygen species production, reduce inflammation and increase neurotrophic factors' function. It has been shown that KD mimics the effects of fasting and the lack of glucose/insulin signaling, promoting a metabolic shift towards fatty acid utilization. In this work, the author reports a number of successful case reports treated through metabolic ketosis.

Biography

Raffaele Pilla is the Doctor Europaeus, received his Master's degree in Pharmacy at G. d'Annunzio University in Chieti-Pescara, Italy in 2005, where he also served internships at the Cell Physiology Laboratory and Molecular Biology Laboratory. He was hired as a postdoctoral scholar in the Department of Pharmacology and Physiology at the University of South Florida in Tampa, on two research grants funded by the Office of Naval Research (US Navy) and Divers' Alert Network. He has written and lectured widely worldwide. He has been involved in ongoing research at the University of South Florida with the use of ketone esters.

raf.pilla@gmail.com

Notes: