

Nashwa Mohammad Hafez Bareedy, Clin Dermatol Res J 2019, Volume: 4

4TH WORLD DERMATOLOGY CONGRESS



3rd International Conference on Diet & Nutrition

June 19-20, 2019 | Berlin, Germany

Hidradenitis suppurativa

Nashwa Mohammad Hafez Bareedy

Jaber Al Ahmad Hospital, Kuwait

idradenitis suppurativa is a chronic inflammatory skin condition which primarily affects intertriginous areas. It develops around puberty. Females are 3 times more than males. Life style has an important role in its pathogenesis. Aetiology and management will be discussed in the conference. No pathognomonic test exists and biopsy is rarely required. Diagnostic criteria, history, laboratory

studies and imaging studies may help in diagnosis of HS. Hurley staging system is the simplest and widely used system to assess severity and extent of HS. HiSCR has been recently used to evaluate therapeutic outcomes in patients with HS. Presence of TNFalfa, IL17and IL23 in HS plays an important role in its pathogenesis. Combined therapy has an important role in its management.

Biography

Nashwa Mohammad Hafez Bareedy has completed Master of Dermatology, Venereology and Cosmetology from Ain Shams University, Egypt and Laser Diploma from Cairo University, Egypt. Presently she is working at Jaber Al Ahmad Hospital, Kuwait. She has published papers in Indian Journal of Dermatology.

nashwabareedy@yahoo.com