



## Review Article

# Upper Semicontinuous Representability of Maximal Elements for Non total Preorders on Compact Spaces

Gianni Bosi<sup>1\*</sup>, Paolo Bevilacqua<sup>2</sup> and Magali Zuanon<sup>3</sup>

### Abstract

We discuss the possibility of determining all the maximal elements of a preorder on a compact topological space by maximizing all the functions in a suitable family of upper semicontinuous order-preserving functions.

### Keywords

Preorder; Order-preserving function; Weak utility; Maximal element; Upper semicontinuous function

### Introduction

White's theorem [1] is important since, for every maximal element  $x_0$  relative to a preorder  $\prec$  on set  $X$ , it guarantees the existence of an order-preserving function  $u$  on the preordered set  $(X, \prec)$  attaining its maximum at  $x_0$ , provided that an order-preserving function  $u'$  on  $(X, \prec)$  exists. So, at least theoretically, every maximal element is obtained by maximizing a real-valued order-preserving function. When this happens, it is clear that every maximal element is potentially optimal in the sense of Podinovski et al. [2] (i.e., for every maximal element there exists a total preorder extending the original preorder with respect to which such maximal element is best preferred).

In this paper, we generalize White's theorem to the upper semicontinuous case. This means that we present conditions on a preorder  $\prec$  on a topological space  $(X, \tau)$  under which, for every maximal element  $x_0$  relative to  $\prec$ , there exists an upper semicontinuous order-preserving function  $u$  on the preordered topological space  $(X, \tau, \prec)$  attaining its maximum at  $x_0$ , provided that an upper semicontinuous order-preserving function  $u'$  on  $(X, \tau, \prec)$  exists. It should be noted that Bevilacqua et al. [3] already characterized the property according to which every maximal element relative to a preorder on a compact topological space can be obtained by maximizing a transfer weakly upper continuous weak utility for its strict part (see the generalization of Weierstrass Theorem presented by Tian et al. [4]).

It is clear that these results are important due to the well-known fact that every upper semicontinuous (more generally transfer weakly upper continuous) function attains its maximum on a compact

topological space, and the nearly obvious consideration that a point  $x_0$  at which an order-preserving function  $u$  for a preorder (or, more generally, a weak utility for its strict part) attains its maximum is also a maximal element for  $\prec$ .

### Notation and preliminaries

Let  $X$  be a nonempty set (decision space). A binary relation  $\succsim$  on  $X$  is interpreted as a weak preference relation, and therefore, for any two elements  $x, y \in X$ , the scripture " $x \succsim y$ " has to be thought of as "the alternative  $y \in X$  is at least as preferable as  $x \in X$ ". As usual,  $<$  denotes the strict part of a binary relation  $\succsim$  (i.e., for all  $x, y \in X$ ,  $x < y$  if and only if  $(x \succsim y)$  and not  $(y \succsim x)$ ). A preorder is a reflexive and transitive binary relation. An anti-symmetric preorder  $\succsim$  is referred to as an order. Furthermore,  $\sim$  stands for the indifference relation (i.e., for all  $x, y \in X$ ,  $x \sim y$  if and only if  $(x \succsim y)$  and  $(y \succsim x)$ ). We have that  $\sim$  is an equivalence relation on  $X$  whenever  $\succsim$  is a preorder.

For every  $x \in X$ , we set

$$I(x) = \{z \in X : z < x\} \quad i(x) = \{z \in X : x < z\}$$

Given a preordered set  $(X, \succsim)$ , a point  $x_0 \in X$  is said to be a maximal element for  $\succsim$  if for no  $z \in X$  it occurs that  $x_0 < z$ . In the sequel we shall denote by  $X_M^{\succsim}$  the set of all the maximal elements of a preordered set  $(X, \succsim)$ . Please observe that  $X_M^{\succsim}$  can be empty.

We recall that a function  $u: (X, \succsim) \rightarrow (R, \leq)$  is said to be

- isotonic or increasing if, for all  $x, y \in X$ ,  $x \succsim y \Rightarrow u(x) \leq u(y)$ ;
- a weak utility for  $<$  if, for all  $x, y \in X$ ,  $x \succsim y \Rightarrow u(x) < u(y)$ ;
- Strictly isotonic or order-preserving if it is both isotonic and a weak utility for  $<$ .

Strictly isotonic functions on  $(X, \succsim)$  are also called Richter-Peleg representations of  $\succsim$  in the economic literature (see e.g. Richter et al. [5] and Peleg et al. [6]).

A preorder  $\succsim$  on a topological space  $(X, \tau)$  is said to be

- upper semicontinuous if, for all  $x, y \in X$ ,  $i(x) = \{z \in X : x \succ z\}$  is a closed subset of  $X$  for every  $x \in X$ ;
- Quasi upper semicontinuous if there exists an upper semicontinuous preorder  $\tilde{\succsim}$  on  $(X, \tau)$  such that  $< \subset \tilde{\prec}$ .

An upper semicontinuous preorder Ward et al. [7] or more generally a quasi-upper semicontinuous preorder Bosi et al. [8, Theorem 3.1]  $\succsim$  on a compact topological space  $(X, \tau)$  admits a maximal element. As usual, for a real-valued function  $u$  on  $X$ , we denote by  $\arg \max u$  the set of all the points  $x \in X$  such that  $u$  attains its maximum at  $x$  (i.e.,  $\arg \max u = \{z \in X : (u(z) \leq u(x)) \forall z \in X\}$ )

If  $\tau$  is a topology on a set  $X$ , and  $\prec$  is a preorder on  $X$ , then the triplet  $(X, \tau, \prec)$  will be referred to as a topological preordered space. The (natural) (interval) topology on the real line  $R$  will be denoted by  $\tau_{nat}$ .

Finally, we recall that a real-valued function  $u$  on a topological space  $(X, \tau)$  is said to be upper semicontinuous if  $u^{-1}((-\infty, \alpha]) = \{x \in X :$

\*Corresponding author: Gianni Bosi, Bruno de Finetti Department of Economics, Business, Mathematics and Statistics (DEAMS), University of Trieste, Italy, Tel: +39-040-558-7115; E-mail: gianni.bosi@deams.units.it

Received: October 20, 2017 Accepted: January 02, 2018 Published: January 08, 2018

$u(x) < \alpha$  is an open set for all  $\alpha \in \mathbb{R}$ . A very well known result guarantees that every upper semicontinuous real-valued function  $u$  on a compact topological space  $(X, \tau)$  attains its maximum.

### Maximal elements of preorders from maximization of upper semicontinuous functions

The following theorem was proved by White et al. [1]. Given any maximal element  $x_0$  relative to a preorder  $\sim$  on a set  $X$ , it guarantees the existence of some order-preserving function  $u$  attaining its maximum at  $x_0$ , provided that an order-preserving function  $u': (X, \sim) \rightarrow (R, \leq)$  exists. Therefore, in order to determine all the maximal elements of a preorder  $\sim$  on a set  $X$ , the agent maximizes all the functions  $u$  in a family  $\mathcal{U}$  of bounded order-preserving functions for  $\sim$ . Needless to say, this is a very important opportunity, at least theoretically.

**Theorem:** (White et al. [1]): Let  $(X, \sim)$  be a preordered set and assume that there exists an order-preserving function  $u': (X, \sim) \rightarrow (R, \leq)$ . If  $X_M^\sim$  is nonempty, then for every  $x_0 \in X_M^\sim$  there exists an order-preserving function  $u: (X, \sim) \rightarrow (R, \leq)$ . Such that  $\arg \max u = [x_0] = \{z \in X: z \sim x_0\}$ .

We now present a generalization of the above theorem to the "upper semicontinuous case".

**Theorem:** Let  $(X, \tau, \sim)$  be a topological preordered space, and assume that  $X_M^\sim$  is nonempty. Consider an element  $x_0 \in X_M^\sim$ . Then the following conditions are equivalent:

- i. There exists an upper semicontinuous order-preserving function  $u': (X, \tau, \sim) \rightarrow (R, \tau_{nat}, \leq)$  and  $[x_0] = \{z \in X: z \sim x_0\}$  is a closed subset of  $X$ ;
- ii. There exists an upper semicontinuous order-preserving function  $u: (X, \tau, \sim) \rightarrow (R, \tau_{nat}, \leq)$  such that  $\arg \max u = [x_0] = \{z \in X: z \sim x_0\}$ .

**Proof:** Consider a topological preordered space  $(X, \tau, \sim)$ .

(i)  $\Rightarrow$  (ii) Let  $u'$  be an upper semicontinuous order-preserving function on  $(X, \tau, \sim)$ . Without loss of generality, we can assume  $u'$  to be bounded. Consider a point  $x_0 \in X_M^\sim$  and define the real-valued function  $u$  on  $X$  as follows for any choice of a positive real  $\delta$ :

$$u'(x) \text{ if not } (x \sim x_0)$$

$$u(x) = \{$$

$$\sup u'(x) + \delta \text{ if } (x \sim x_0)$$

White et al. [1, Theorem 1] proved that the above function  $u$  is order-preserving for  $\sim$  as soon as  $u'$  is order-preserving for  $\sim$ . For the sake of completeness, let us recall here the arguments supporting this consideration. It is clear that  $u'(x) \leq u(x)$  for every  $x \in X$ . In order to show that  $u$  is increasing with respect to  $\sim$ , consider any two points  $x, y \in X$  such that  $x \sim y$ . If  $y \sim x_0$ , then it is clear that  $u(x) \leq u(y)$  from the definition of  $u$ . On the other hand, if  $\text{not}(y \sim x_0)$ , then it must be also  $\text{not}(x \sim x_0)$ , since  $x_0 \sim x \sim y$  would imply  $(x_0 \sim y)$ , and in turn  $x_0 \sim y$  due to the fact that  $x_0$  is a maximal element relative to  $\sim$ . Hence, since neither  $x \sim x_0$  nor  $y \sim x_0$ , we have that  $u(x) = u'(x) \leq u'(y) = u(y)$  from the definition of  $u$  and the fact that  $u'$  is increasing with respect to  $\sim$ . In order to show that  $u$  is a weak utility for  $\prec$ , consider any two points  $x, y \in X$  such that  $x \prec y$ . Then we have

that  $\text{not}(x \sim x_0)$ , since  $x_0 \sim x \prec y$  implies that  $x_0 \sim y$  (a contradiction, since  $x_0$  is assumed to be a maximal element for  $\sim$ ). Therefore, from the definition of  $u$  and the fact that  $u'$  is a weak utility for  $\prec$ , we have that  $u(x) = u'(x) < u'(y) \leq u(y)$ , which obviously implies that  $u(x) < u(y)$ . Further,  $u$  attains its maximum at  $x_0$  and actually, since  $\delta$  is a positive real, it is clear that (i)  $\arg \max u = [x_0] = \{z \in X: z \sim x_0\}$ . Therefore, it only remains to show that under our assumptions  $u$  is upper semicontinuous. It is clear that  $u$  is upper semicontinuous at every point  $x \in X$  such that  $x \sim x_0$ . Therefore, consider any point  $x \in X$  such that  $\text{not } x \sim x_0$  and  $\alpha \in \mathbb{R}$  such that  $u(x) < \alpha$ . We can limit our considerations to the case when  $\alpha \leq \sup u'(x) + \delta$ . Since in this case  $u(x) = u'(x)$ , from upper semi continuity of  $u'$  there exists an open set  $U_x'$  containing  $x$  such that  $u'(z) < \alpha$  for every  $z \in U_x'$ . Since  $[x_0]$  is closed, we have that  $U_x = U_x' \setminus [x_0]$  is an open set containing  $x$  such that  $u'(z) = u(z) < \alpha$  for every  $z \in U_x$ . Hence,  $u$  is an upper semicontinuous function.

(ii)  $\Rightarrow$  (i) Assume that there exists an upper semicontinuous order-preserving function  $u: (X, \tau, \sim) \rightarrow (R, \tau_{nat}, \leq)$  such that  $\arg \max u = [x_0] = \{z \in X: z \sim x_0\}$ . If  $[x_0]$  is not closed, then there exists an element  $z \in X \setminus [x_0]$  such that  $U_z \cap [x_0] \neq \emptyset$  for every neighborhood  $U_z$  of the element  $z$ . But this contradicts the fact that  $u$  is upper semicontinuous, since in this case  $u'(\cdot) - \infty, u(x_0)[\cdot]$ , an open set containing  $z$ , should contain an element  $z' \in [x_0]$ , for which  $u(z') = u(x_0)$ . This consideration completes the proof.

**Remark:** It is clear that Theorem 3.2 generalizes White's theorem, due to the fact that these two results precisely coincide when we consider the discrete topology  $\tau$  on  $X$ .

Since in order to determine a maximal element relative to preorder  $\sim$  on a set  $X$  it suffices to maximize a weak utility for the strict part  $\prec$  of  $\sim$ , the following corollary can be considered as useful. Indeed, the reader can easily verify that the implication "(i)  $\Rightarrow$  (ii)" in Theorem 3.2 is still valid if one considers weak utilities for  $\prec$  instead of order-preserving functions  $u'$  for  $\sim$ .

**Corollary:** Let  $(X, \tau, \sim)$  be a topological preordered space with  $\tau$  a compact topology. If there exists an upper semicontinuous weak utility  $u'$  for  $\prec$ , and  $[x] = \{z \in X: z \sim x\}$  is a closed set for all  $x \in X_M^\sim$ , then for every  $x_0 \in X_M^\sim$  there exists an upper semicontinuous weak utility  $u$  for  $\prec$  such that  $\arg \max [x] = \{z \in X: z \sim x_0\}$ .

Bosi et al. [8, Theorem 2.11] proved that there exists an upper semicontinuous weak utility for the strict part  $\prec$  of a quasi-upper semicontinuous preorder  $\sim$  on a second countable (i.e., with a countable base) topological space  $(X, \tau)$ . Hence, we get the following corollary.

**Corollary:** Let  $(X, \tau, \sim)$  be a topological preordered space. Assume that  $\tau$  is a compact and second countable topology, and that  $\sim$  is quasi upper semicontinuous. If  $[x] = \{z \in X: z \sim x\}$  is a closed set for all  $x \in X_M^\sim$ , then for every  $x_0 \in X_M^\sim$  there exists an upper semicontinuous weak utility  $u$  for  $\prec$  such that  $\arg \max [x] = \{z \in X: z \sim x_0\}$ .

### Conclusion

In this paper, following the spirit of a theorem of White et al. [1], we have presented some results concerning the representation of the set of all maximal elements of a preorder on a compact topological space by means of the maximization of all functions in a suitable family of bounded upper semicontinuous order-preserving functions. The more delicate problem of characterizing the possibility

of representing all the maximal elements for a preorder on a compact topological space by means of the maximization of finitely many upper semicontinuous order-preserving functions will be hopefully considered in a future paper.

#### Reference

1. White DJ (1980) Notes in decision theory: optimality and efficiency II. Eur J Oper Res 4: 426-427.
2. Podinovski VV (2013) Non-dominance and potential optimality for partial preference relations. Eur J Oper Res 229: 482-486.
3. Bevilacqua P, Bosi G, Zuanon M (2018) Maximal elements of preorders from maximization of transfer upper continuous weak utilities on a compact space. Far East J Math Sci 103: 213-221.
4. Tian G, Zhou J (1995) Transfer continuities, generalizations of the Weierstrass and maximum theorems: a full characterization. J Math Econ 24: 281-303.
5. Richter MK (1966) Revealed preference theory. Econometrica 34: 635-645.
6. Peleg B (1970) Utility functions for partially ordered topological spaces. Econometrica 38: 93-96.
7. Ward LE (1954) Partially ordered topological spaces. P Am Math Soc 5: 144-161.
8. Bosi G, Zuanon ME (2017) Maximal elements of quasi upper semi continuous preorders on compact spaces. Econ Theory Bulletin 5: 109-117.

#### Author Affiliation

[Top](#)

<sup>1</sup>Bruno de Finetti Department of Economics, Business, Mathematics and Statistics (DEAMS), University of Trieste, Italy

<sup>2</sup>Department of Engineering and Architecture, University of Trieste, Italy

<sup>3</sup>Department of Economy and management, University of Brescia, Italy

#### Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • [www.scitechnol.com/submission](http://www.scitechnol.com/submission)