



# Appraisal of Thrips Population Dynamics in Mango using Weather Based Indices

Gundappa\*, Adak T and Shukla PK

### Abstract

The aim of this investigation was to assess the dynamics of mango thrips population across 20 mango orchards based on thermal indices under subtropical Lucknow condition during 2013 and 2014. Wide variations in thrips populations were observed across different critical crop phenological stages and seasons. The thrips population was observed between 13<sup>th</sup> SMW to 24<sup>th</sup> SMW during 2013 cropping season, whereas it was found from 15<sup>th</sup> SMW to 28<sup>th</sup> SMW in the subsequent year. The mean thrips population across standard weeks was highest ( $6.18 \pm 0.14$ ) during 15<sup>th</sup> SMW followed by  $3.13 \pm 0.15$  in 16<sup>th</sup> and  $1.87 \pm 0.06$  in 17<sup>th</sup> SMW during 2013, whereas in the next year (2014), the highest values were  $4.67 \pm 0.09$  followed by  $4.19 \pm 0.10$  and  $4.07 \pm 0.15$  in 16<sup>th</sup>, 17<sup>th</sup> and 18<sup>th</sup> SMW respectively. Step wise regression analysis revealed that maximum temperature, minimum temperature, maximum relative humidity, minimum relative humidity, sunshine hours and evaporation could explained 55 per cent variations in thrips population. Different thermal indices viz., growing degree days (GDD), heliothermal units (HTU), photothermal units (PTU), were cumulated up to peak population density and a positive and significant correlation was revealed. The best fit polynomial regression analysis indicated that thrips population could be predicted up to 95 per cent using these thermal indices. Thus, it was concluded that use of all these robust indices may be useful in assessing the pest-weather dynamics of mango growing region and serve as a basis for real time pest management advisory.

### Keywords

Thermal indices; GDD; Mango; Thrips; Prediction model

## Introduction

Mango (*Mangifera indica* L.), one of the major fruit crops of India, is known as the king of fruits for its sweetness, excellent flavour, delicious taste and high nutritive value [1]. It is attacked by several pests during its vegetative and reproductive phases [2]. During the reproductive phase of the crop, pests like hoppers and thrips pose a threat to mango production. Under the era of changing climatic situation, thrips are becoming serious pest on mango. Several species of thrips are observed in mango orchards in Florida, the Caribbean, Central and South America, and Asia [3-5]. Among the thrips species, *Scirtothrips dorsalis* and *Thrips hawaiiensis* were recorded as severe pests of various vegetables, fruits and ornamental crops in

eastern Asia [6,7]. In India, *Thrips palmi* was recorded on mango inflorescence [8], from Andhra Pradesh three species of thrips viz., *Megalurothrips distalis* (Karny), *Thrips hawaiiensis* and *Haplothrips tenuipennis* are reported on mango [9-11]. Two species of thrips *Scirtothrips mangiferae* Hood and *S. dorsalis* Hood were reported by Kumar and Bhatt [12] respectively, from Gujarat on mango. Kumar et al. [13] reported from Cuttack that *Scirtothrips mangiferae*, *S. dorsalis*, *Rhipiphorotheirus cruentatus* Hood are occurring on mango. From Haryana *R. cruentatus* was reported by Dahiya and Lakra [14]. Patel et al. [15] reported three species of thrips on mango rootstock viz., *Pantachaetothrips* sp., *Selenothrips rubrocinctus* Giard, *Caliothrips impurus* Priesner, besides *Scirtothrips dorsalis*. From Kerala Ananthakrishnan and Muraleedharan [16] reported *Selenothrips rubrocinctus* on mango. Recently *Frankliniella schultzei* and *Thrips subnudula* reported on inflorescence of mango from Tamil Nadu [17]. Thrips pose an increasing threat to mango production, which can cause considerable economic loss in mango orchards. Indiscriminate use of synthetic insecticides for routine thrips control is detrimental to non-target organisms and causes pesticide residues on food and in the environment. Other destructive consequences include the development of resistant pest populations, pest resurgence and the outbreak of secondary pest infestations [18,19]. To minimize harmful effects of synthetic pesticides, development of non-polluting plant protection strategies is necessary. For rational use of synthetic pesticides forewarning/prediction models for the destructive pest like thrips are need of the hour.

Thermal characteristics may vary among species, populations, developmental stages, and with other ecological factors such as food source [20-23]. This effect of temperature can be described by specific rate functions of temperature for development (thermal models), which are used in predicting insect pests interaction with biotic and abiotic factors [24]. The pest forecasting models facilitate better preparedness to combat outbreaks of serious insect pests by developing effective pest management strategies well in advance. The temperature-response phenology models simulate the variability in insect development times within a population based on the detailed laboratory assessments of the insect's life history and thus, can provide better results on future pest activity [25]. In this context this study aimed to develop weather based prediction models for mango thrips.

## Materials and Methods

A field experiment was conducted for two consecutive seasons (2013 and 2014) on a *Typic ustochrepts* soil in Lucknow, Uttar Pradesh, India at 22 locations. The climate of the experimental site is semi-arid subtropical with hot dry summers and cold winters. Orchards of Mango cv. *Dashehari* of 20-35 years age were selected with planting 7.5×7.5 to 10×10 mt. Each orchard had at least 25 trees. Data on thrips population was recorded on weekly basis from 5 randomly selected trees in four direction of the tree. Thrips count was taken after gently tapping shoot or panicle by holding white paper in the palm. For analysis, mean count per shoot or panicle was taken and expressed as number of thrips per shoot or panicle. Daily weather data of temperature (maximum and minimum), relative humidity (morning and evening), rainfall, wind speed, bright sunshine hours

\*Corresponding author: Gundappa, ICAR- Central Institute for Subtropical Horticulture, Rehmankhera, Lucknow-226101, Uttar Pradesh, India, Tel: +91-522 284 1022; E-mail: gundappacish@gmail.com

Received: July 28, 2016 Accepted: August 10, 2016 Published: August 17, 2016

and evaporation rates were recorded in the agromet observatory located within the experimental site. Temperature, solar radiation and day length data were used to compute agro climatic models. These models were calculated as follows:

$$\text{Growing Degree days (GDD)} = \sum(T_{\text{Max}} + T_{\text{Min}}) / 2 - T_{\text{Base}}$$

Where  $T_{\text{Max}}$  and  $T_{\text{Min}}$  are the maximum and minimum temperatures ( $^{\circ}\text{C}$ ) of the day and  $T_{\text{Base}}$  is base temperature which was taken as  $15^{\circ}\text{C}$  [26].

$$\text{Heliothermal units (HTU)} = \text{GDD} \times \text{actual bright sunshine hour (n)}$$

$$\text{Photothermal units (PTU)} = \text{GDD} \times \text{day length (N)}$$

Where N indicates maximum possible bright sunshine hours or day length and calculated as

$$N = (24/\pi) \times Ws$$

Ws is the sunset hour angle (Radian) = Arc Cosine  $[-\tan(\Phi) \times \tan(\sigma)]$

$\Phi$  = Latitude in radian,

$\sigma$  = Solar declination in radiation, calculated as follows

$$\sigma = 0.409 \times \text{Sine} [(2 \times \pi \times J) / d - 1.39]$$

Where J = Julian days (1 to 365/366) and d = No. of days in the year

These models were computed on daily basis taking 1<sup>st</sup> September as base for each year since mango is harvested during June and July in Northern India and post harvest vegetative phase started. The thrips occurrence was also plotted against these models and cumulated up to peak/maximum thrips incidence and severity for generating regression based prediction models. Statistical analysis viz., regression equations, were carried out using MS Excel and SPSS packages (Version 12.0). The required statistically significant graphs were drawn using MS Excel/Power Point software packages.

## Results and Discussion

### Weather situation during study period

Weather parameters particularly maximum and minimum temperatures, rainfall, pan evaporation, relative humidity etc. has profound influence on mango crop phenology in turn pest-weather dynamics. Henceforth, analyses of weather parameters are essential in assessing the pest population dynamics. The analysis showed maximum temperature ranged between  $29.8$  to  $40.3^{\circ}\text{C}$  during 2013 crop season and  $25.7$  to  $43.4^{\circ}\text{C}$  during 2014 crop season at the time of first appearance of thrips population on mango and its peak values. However, the minimum temperature was recorded lower in 2014 season as compared to 2013. A range of  $9.5$  to  $26.8^{\circ}\text{C}$  was observed in both the seasons. The rainfall was widely distributed in 2013 than 2014 fruiting season. Higher pan evaporation was recorded in 2014 as compared to 2013 (Figure 1).

### Population dynamics of mango thrips

A wide variation in thrips occurrence on the mango crop at its different phenological stages was revealed in 20 mango fixed orchards during two consecutive mango cropping season (2013 and 2014). The first appearance of thrips population was observed on 13<sup>th</sup> standard meteorological week (SMW) and 15<sup>th</sup> SMW in 2013 and 2014 fruiting season respectively (Table 1). The infestation of thrips on mango

crop observed till 24<sup>th</sup> SMW during 2013 and up to 28<sup>th</sup> SMW during 2014. Among the 20 mango orchards, highest thrips population density ( $13.6$  /panicle) was recorded in 15<sup>th</sup> SMW at Kakori orchard (II), followed by  $10.5$  /panicle and  $10.3$  /panicle in Kakori (I) and NB Dhanewa (I) orchard in 2013 season. The higher values of thrips population density were recorded in 15<sup>th</sup> and 16<sup>th</sup> SMW in 2013 while in the next season, it was observed from 17<sup>th</sup> to 20<sup>th</sup> SMW. Some of the mango orchards viz., Malihabad ( $9.45$  and  $9.15$  /panicle), Kanar-II ( $9.4$  /panicle) [21] and Allapur-I ( $8.8$  /panicle) showed higher thrips infestation at flowering and initial fruit setting stage. In 2014 mango season, highest thrips infestation was recorded in CISH Block II as  $13.0$  /panicle followed by  $10.1$  /panicle (Allapur (II)). Mango orchards




Figure 1: Weather dynamics during study period (2013-14).

of CISH Block III had thrips density of 8.95 to 9.3/panicle followed by Ulrapur-II (8.85/panicle) and Mahmood Nagar-I (8.75 /panicle). Such wide spatio-temporal variations in thrips populations across 20 mango orchards and seasons may be due to differences in cultural practices adopted by the mango growers and weather dynamics. The mean thrips population across standard weeks was highest ( $6.18 \pm 0.14$ ) during 15<sup>th</sup> SMW followed by  $3.13 \pm 0.15$  in 16<sup>th</sup> and  $1.87 \pm 0.06$  in 17<sup>th</sup> SMW during 2013, whereas in the next year (2014), the highest values were  $4.67 \pm 0.09$  followed by  $4.19 \pm 0.10$  and  $4.07 \pm 0.15$  in 16<sup>th</sup>, 17<sup>th</sup> and 18<sup>th</sup> SMW respectively (Table 2). Although there were significant variations in thrips population in SMW and orchards, a range of 0.15 to 13.6 in 2013 and 0.15 to 13.0 in 2014 was found. Peak occurrence of thrips was found during the flowering and fruiting developmental stages of mango. Appearance of thrips population on mango was delayed one week during the year 2014 compared


to 2013 (Figure 2), this is attributed to the maximum temperature which was higher and influenced the faster growth and development of thrips. Higher temperature also play key role in the development of vegetative as well as reproductive growth of the mango. Prolonged lower night temperature (minimum temperature) may be responsible factor for the longer stay of thrips on mango for one more month in 2014 cropping season. The peak infestation of thrips on mango was synchronized with the flowering, fruit development and vegetative flushes of mango in the Lucknow region. Changes in the phenology of crop may also influence the pest occurrence. The impact of temperature and rainfall on the mango crop phenology was observed in Lucknow region [27] and pest-weather dynamics during critical crop phenological stages was also reported by Ravishankar et al. [28]. It is obvious that abiotic factors had profound influence on the dynamics of thrips population across regions. In a study in

**Table 1:** Mean density of mango thrips population in 20 orchards during 2013 and 2014 at Lucknow, Uttara Pradesh.

Year	2013												2014													
	Standard meteorological weeks																									
Location	13	14	15	16	17	18	19	20	21	22	23	24	15	16	17	18	19	20	21	22	23	24	25	26	27	28
Malihabad (Fixed I)	0	4.55	9.15	0.85	0.65	0.4	1	0.5	1	1.1	0	0.4	0	3.95	4.25	4.9	1.75	1.25	0.6	1.2	1	0	0.65	0.75	0.4	0
Malihabad (Fixed II)	0	5	9.45	2.15	0.55	0.25	0	0.8	0.65	1	0	0	1.05	4.45	4.6	6.95	1.45	0.5	0.4	1	1	0.4	0.5	0	0.5	0
Navipana (Fixed I)	0	3.35	3.95	1.15	0.6	0	0	0	0.6	0.25	0	0.6	6.9	5.05	2.35	0.7	0.75	0	0.8	0.65	1.5	0.4	0	0.65	0	0
Navipana (Fixed II)	0	5.85	6.2	3.1	0.75	0	0	0.35	1.2	0.5	0	0.85	6.75	6.6	2.5	1.85	1.75	1	0.75	1	1	0.65	0	0	0	0
Methe Nagar (Fixed I)	0	1.35	5.65	3.45	2.35	0.6	0.15	0.95	1.05	0	0.7	1	0	6.4	3.3	0.95	2.8	1	1	0	0.75	0.6	0	0	0	0
Methe Nagar (Fixed II)	0	2.75	5.6	0.6	2.05	0.8	0	0.55	1.3	2.5	1.15	1.15	2.7	5.5	4.7	3.65	1.2	0.5	0.25	0.8	1.15	0	0	0	0	0
Hafizkhera (Fixed I)	0	0	5.3	1.85	4.25	0.85	0.35	1.15	0.95	0.55	0.25	1.05	1.4	5.7	3.15	0.15	0.5	1	0.5	0.85	1.15	0	0	0	0	0
Hafizkhera (Fixed II)	0	0	4.25	1.55	4.8	1.2	0.25	1.3	1.05	0.65	0.45	0.85	3.1	5.3	3.15	2.75	3	1.15	3.7	0.9	0.75	0.85	0.6	0	0	0
Mahmood Nagar (Fixed I)	0	6	6.9	2.75	1.8	1.15	0.4	0.6	1	1	0	0.4	1.05	3.65	4.55	8.75	2.85	0.65	0.75	1.25	0.75	0	0	0.4	1.1	0
Mahmood Nagar (Fixed II)	0	2.15	7.55	1.2	0.25	0.9	0.65	0.75	1	0.75	0	0	0	4.95	2.75	5.2	1.8	0	0.6	1.15	0	1	0	0.4	0	0
Kakori (Fixed I)	0	6.4	13.6	10.5	3.25	0.85	0.7	0.65	2.5	1	0.9	0	0	0	4.3	7.3	3.25	0	0.4	1	0.6	0	0	0	0	0
Kakori (Fixed II)	0	0.7	8.15	4.55	1.35	1.55	0.95	0.75	1.15	1	0.95	0.25	2.7	5.05	5.95	4.4	3.3	1	1.1	1.55	1.25	1.15	0.9	0.9	1	0
Ulrapur (Fixed I)	0	3.95	1.1	1.4	2.5	0.6	0	0.7	1.25	1.05	0.8	0	5	2.9	7.85	4	1.45	0.8	3	1.25	0.6	0.35	0.4	0	0	0
Ulrapur (Fixed II)	0	3.75	6.45	1.2	3.1	0.25	0.35	0.35	0.25	0.65	0.6	0.4	2.65	3.05	8.85	0	1.3	0.65	0.35	1.65	1	0	0	0	4	0
Kanar (Fixed I)	0	1.1	4.5	5.8	2.3	0.7	1.1	0.65	0.65	0.6	1	0.4	0	5.05	4.4	4.6	1.1	1	0.5	1	0.9	0.9	0	0.25	0	0
Kanar (Fixed II)	0	1.55	7.3	9.4	0.8	0.55	0.55	0.6	1.15	1.5	1.35	0	1.1	4.1	4.85	3.4	1.55	0.25	1.3	0.5	0	0.75	0.75	0	0	0
NB Dhanewa (Fixed I)	1.6	2.75	10.3	1.65	1.95	0.6	1.45	1.05	0.5	1.2	0	1.1	5.1	8.2	1.75	1.5	1	1.25	1.65	0.75	0.75	0.8	0.85	0.4	0	0
NB Dhanewa (Fixed II)	1.75	2.65	6.6	2.25	1.05	0.25	2.25	0.6	1.05	0.55	0	0.75	5.95	5.55	2.4	0.75	0	0	1.15	0	0.85	0	0	0.65	0	0
CISH Block-III (Fixed I)	3.35	7.15	1	1.2	2.25	0.25	2.95	6.2	1.4	1.35	1.65	0	7.4	6.65	8.95	9.3	7.45	6.5	5.35	2	0	0	1.35	0	1.4	0.4
CISH Block-II (Fixed II)	0	1.4	3.8	2.7	3.05	1.85	0.4	0.6	6.2	2.25	0.75	0	1.5	3.9	2.85	3.15	4.1	13	2.75	4.65	1	1	0.9	4.25	0	0
Allupur (Fixed I)	0	1.05	4.9	0.65	0.85	0.8	0.4	1.05	0.85	1.4	0.75	0	0	1.7	2.55	5.15	0.65	0.35	0.95	1.35	2	0	0.65	0.9	0.4	0
Allupur (Fixed II)	0	1.55	4.2	8.8	0.55	1.25	0	0.5	0.6	1	0.8	0	1.95	4.95	2.1	10.1	5.25	2.55	2.5	1.6	1.25	0	0.75	1.15	0	0.9

**Table 2:** Univariate statistical analysis for the population of mango thrips across the standard meteorological weeks (SMW) during 2013 and 2014 at Lucknow, Uttara Pradesh.

SMW	Mean	Range	2013		SMW	Mean	Range	2014	
			Skeweness	Kurtosis				Skeweness	kurtosis
13	0.30 ± 0.04	3.35-1.60	2.91	8.43	15	2.56 ± 0.13	7.40 - 1.05	0.74	-0.79
14	2.95 ± 0.11	7.15-0.70	0.50	-0.83	16	4.67 ± 0.09	8.20 - 1.70	-0.69	1.51
15	6.18 ± 0.14	13.60-1.00	0.51	1.04	17	4.19 ± 0.10	8.95 - 1.75	1.23	0.82
16	3.13 ± 0.15	10.50-0.60	1.61	1.55	18	4.07 ± 0.15	10.10 - 0.15	0.52	-0.54
17	1.87 ± 0.06	4.80-0.55	0.74	-0.11	19	2.19 ± 0.09	7.45 - 0.65	1.57	2.96
18	0.71 ± 0.02	1.85-0.25	0.61	0.18	20	1.56 ± 0.14	13.00 - 0.35	3.45	12.60
19	0.63 ± 0.04	2.95-0.15	1.85	3.50	21	1.38 ± 0.07	5.35 - 0.40	1.80	3.06
20	0.94 ± 0.06	6.20-0.35	4.25	19.19	22	1.19 ± 0.05	4.65 - 0.40	2.64	10.17
21	1.24 ± 0.06	6.20-0.25	3.76	15.67	23	0.88 ± 0.02	2.00 - 0.60	-0.12	1.09
22	0.99 ± 0.03	2.50-0.25	1.02	1.67	24	0.40 ± 0.02	1.15 - 0.35	0.38	-1.51
23	0.55 ± 0.03	1.65-0.25	0.38	-0.81	25	0.38 ± 0.02	1.35 - 0.40	0.60	-0.79
24	0.42 ± 0.02	1.15-0.25	0.46	-1.35	26	0.49 ± 0.05	4.25 - 0.25	3.54	14.40
-	-	-	-	-	27	0.40 ± 0.05	4.00 - 0.40	3.33	12.52
-	-	-	-	-	28	0.06 ± 0.01	0.90 - 0.40	3.76	14.53



Malaysia, three peak populations of mango thrips were recorded and the population was significantly influenced by abiotic factors like temperature, relative humidity on thrips population dynamics [29].

**Thrips population relationship with weather parameters**

For development of efficient pest management approach, understanding of association between pest population and weather parameters is pre-requisite for any particular location [30-33]. Many workers have reported that weather factors as one amongst the major contributing factor to the outbreak of pests on mango across the country, workers have described the relationship of ecological factor with the pest population under specific location [34-37]. Weather variables including rainfall, temperature, relative humidity and wind

have been reported as important factors that significantly affect thrips population [38,39]. In our study thrips population was found negatively correlated with maximum relative humidity ( $r = -0.792^{**}$ ) and minimum relative humidity ( $r = -0.670^{**}$ ) during the year 2013. Whereas during the year 2014 thrips population was found negatively correlated with minimum temperature ( $r = -0.865^{**}$ ), minimum relative humidity ( $r = -0.634^{**}$ ), maximum relative humidity ( $r = -0.577^{**}$ ) and evaporation ( $r = -0.581^{**}$ ) (Table 3). Similar results were reported by Duraimurugam and Jagadish [40], Shukla [41], Panickar and Patel [42], Nandini et al. [43] and Vanisree et al [44], Meena et al. [45] on various crops. Step wise regression analysis revealed that maximum temperature, minimum temperature and maximum relative humidity, minimum relative humidity, sunshine

hours and evaporation could explained 55 per cent variations in thrips population ( $p < 0.009$ ) (Table 4). Similar finding was reported in case of coffee borer [46] where correlation studies indicated that different weather parameters influence pest population differently across seasons.

**Relationship between thermal indices of mango vis-à-vis thrips population**

Different thermal indices (growing degree days, Photothermal units and heliothermal unit) were calculated for mango based on the weather data of experimental station. It was estimated that the thermal heat accumulation (GDD) and two other indices (HTU and PTU) were 1007.28°Cday, and 8951.67 and 12608.22 at the time of peak occurrence of thrips population during 2013 while the corresponding values were 1123.04, 8273.42 and 14256.98 of GDD, HTU and PTU respectively during 2014 crop season (Table 5). The significantly positive relationship was observed between thermal indices of mango viz., growing degree days ( $r = 0.946^{**}$ ), Photothermal units ( $r = 0.915^{**}$ ) and heliothermal unit ( $r = 0.940^{**}$ ) with the thrips population. The different thermal indices of mango along with thrips population were subjected for the regression analysis to develop prediction models. This revealed that different regression

equation (Linear, Logarithmic and exponential) had explained good amount of variation. During the year 2013 exponential model had explained highest variation for GDD ( $y = 0.0421e^{0.0024x}$ ;  $R^2 = 0.97^{**}$ ), HTU ( $y = 0.0394e^{0.0003x}$ ;  $R^2 = 0.90^{**}$ ) and PTU ( $y = 0.0616e^{0.0002x}$ ;  $R^2 = 0.90^{**}$ ) compare to linear and logarithmic models. During the year 2014, compared to exponential and logarithmic models linear models for GDD ( $y = 0.0033x - 3.8391$ ;  $R^2 = 0.91^{**}$ ), HTU ( $y = 0.0004x - 3.6391$ ;  $R^2 = 0.90^{**}$ ) and PTU ( $y = 0.0002x - 3.3693$ ;  $R^2 = 0.90^{**}$ ) had explained highest variation (Table 6). Stepwise regression analysis between different thermal indices and thrips population had explained considerable amount of variation. Growing degree days of mango along with photothermal units ( $y = 0.046X_1 - 0.002X_3 - 9.56$ ;  $R^2 = 0.97^{**}$ ) had explained about 97 per cent of variation. All thermal indices parameters together had also explained up to 97 per cent of variation (Thrips population =  $0.042 \times GDD - 0.0003 \times HTU - 0.002 \times PTU - 8.97$ ;  $R^2 = 0.97^{**}$ ) (Table 7). It indicated that GDD and PTU are important thermal indices play a key role for prediction of mango thrips population. Based on three years study (2011-14) at Lucknow, Uttar Pradesh Region with 20-35 years old mango orchards of cv. *Dashehari*, it was found that mango hopper populations could be predicted 80 to 90% using thermal indices [47,48].

**Table 3:** Correlation coefficients (r) between mango thrips population and weather parameters.

Weather parameter	2013	2014
Temperature (Min)	NS	-0.865**
Relative humidity (Max)	-0.792**	-0.634*
Relative humidity (Min)	-0.670*	-0.577*
Evaporation	NS	-0.581*

\*\* . Correlation is significant at the 0.01 level (2-tailed).  
\* . Correlation is significant at the 0.05 level (2-tailed).

**Table 4:** Step-wise regression analysis for mango thrips population with weather parameters.

Regression equation	R <sup>2</sup>	F	Sig.
$y = 8.76 + 0.17X_1 - 0.22X_2 - 0.09X_3 + 0.02X_4 - 0.29X_5 - 0.10X_6$	0.55	3.99	$p < 0.009$
$y = 8.91 + 0.12X_1 - 0.25X_2 - 0.08X_3 + 0.03X_4 - 0.24X_5$	0.54	4.85	$p < 0.004$
$y = 9.48 + 0.03X_1 - 0.23X_2 - 0.09X_3 + 0.05X_4$	0.52	5.76	$P < 0.002$
$y = 10.77 - 0.05X_1 - 0.12X_2 - 0.06X_3$	0.50	7.60	$p < 0.001$
$y = 4.23 + 0.08X_1 - 0.27X_2$	0.44	9.24	$p < 0.001$

Y = Population ; X<sub>1</sub> = Temperature (max) ; X<sub>2</sub> = Temperature (min); X<sub>3</sub> = Relative humidity (max); X<sub>4</sub> = Relative humidity (min); X<sub>5</sub> = Sunshine hours; X<sub>6</sub> = Evaporation

**Table 5:** Thermal indices during the incidence of mango thrips across two consecutive seasons.


Std. week	2013				2014				
	Thrips Population	GDD	Helio thermal unit	Photothermal Unit	Std. week	Thrips Population	GDD	Helio thermal unit	Photothermal unit
13	0.30	870.0214	6694.368	10567.99	15	2.56	1052.11	9675.61	13168.96
14	2.95	930.3071	9265.216	11473.83	16	4.67	1123.04	8273.42	14256.98
15	6.18	1007.286	8951.676	12608.22	17	4.19	1197.54	12301.28	15408.25
16	3.13	1096.25	8888.737	13917.22	18	4.07	1295.87	12484.18	16883.10
17	1.87	1178.986	10749.92	15169.45	19	2.19	1395.37	12743.66	18388.41
18	0.71	1283.707	14010.01	16724.74	20	1.56	1499.96	15589.56	19968.88
19	0.63	1391.707	13303.82	18340.2	21	1.38	1606.84	15983.27	21580.19
20	0.94	1505.086	15319.32	20037.27	22	1.19	1714.41	17596.56	23190.68
21	1.24	1623.95	15449.41	21810.18	23	0.88	1835.74	18933.47	24967.19
22	0.99	1753.864	15990.74	23724.45	24	0.40	1966.50	15888.69	26840.67
23	0.55	1859.586	13759.69	25291.35	25	0.38	2079.82	8703.79	28432.51
24	0.42	1969.121	11779.05	26876.4	26	0.49	2200.33	18683.86	30067.42
					27	0.40	2318.12	9914.10	31600.91
					28	0.06	2420.79	10717.02	32857.57



**Table 6:** Linear and non-linear regression analysis for prediction of thrips population as a function of thermal indices.

	2013			2014		
	Equation	R <sup>2</sup>	r	Equation	R <sup>2</sup>	r
<b>GDD</b>						
Linear	y = 0.004x - 3.7961	0.75	0.867	y = 0.0033x - 3.8391	0.91	0.958
Logarithmic	y = 5.067ln(x) - 34.772	0.67	0.802	y = 5.2482ln(x) - 37.092	0.85	0.925
Exponential	y = 0.0421e <sup>0.0024x</sup>	0.97	0.986	y = 0.0146e <sup>0.0025x</sup>	0.88	0.939
<b>HTU</b>						
Linear	y = 0.0004x - 3.5193	0.60	0.776	y = 0.0004x - 3.6391	0.90	0.948
Logarithmic	y = 4.4634ln(x) - 40.121	0.52	0.724	y = 5.0615ln(x) - 46.168	0.84	0.917
Exponential	y = 0.0394e <sup>0.0003x</sup>	0.90	0.950	y = 0.0169e <sup>0.0003x</sup>	0.86	0.932
<b>PTU</b>						
Linear	y = 0.0003x - 3.1569	0.75	0.866	y = 0.0002x - 3.3693	0.90	0.951
Logarithmic	y = 4.3919ln(x) - 41.188	0.65	0.812	y = 4.6993ln(x) - 45.193	0.83	0.912
Exponential	y = 0.0616e <sup>0.0002x</sup>	0.97	0.986	y = 0.02e <sup>0.0002x</sup>	0.89	0.943

y= Population; GDD – Growing degree days; HTU- Heliothermal units; PTU- Photothermal units;



**Figure 3:** Prediction models for mango thrips using different thermal indices.

**Table 7:** Step-wise regression analysis between the thrips population and weather based indices.

Linear Regression equation	R <sup>2</sup>	F	Sig.
y = 0.003X <sub>1</sub> -3.58	0.89	205.38	p<0.000
y = 0.00042X <sub>2</sub> -3.73	0.83	123.35	p<0.000
y = 0.00023X <sub>3</sub> -3.07	0.88	181.00	p<0.000
y = 0.009X <sub>1</sub> -0.0007X <sub>2</sub> -2.93	0.93	166.36	p<0.000
y = 0.046X <sub>1</sub> -0.002X <sub>3</sub> -9.56	0.97	375.27	p<0.000
y = 0.00068X <sub>2</sub> -0.0005X <sub>3</sub> -1.74	0.91	117.03	p<0.000
y = 0.042 X <sub>1</sub> -0.0003X <sub>2</sub> -0.002X <sub>3</sub> -8.31	0.97	311.80	p<0.000

y= Population X<sub>1</sub> = Growing degree days (GDD) ; X<sub>2</sub>= Heliothermal units (HTU); X<sub>3</sub> = Photothermal units (PTU)

**Prediction models for mango thrips**

Positive relation between thrips populations up to its peak value was observed against these thermal indices, the progressive changes in peak thrips incidence and population. Henceforth, cutting across the years, the peak thrips population were pooled (Figure 3), and was predicted by exponential regression equations of following types;

Mango thrips population = 0.0304e<sup>0.002xGDD</sup> (n = 26, R<sup>2</sup> = 0.91\*\*)

Mango thrips population = 0.0232e<sup>0.0003xHTU</sup> (n = 26, R<sup>2</sup> = 0.93\*\*)

Mango thrips population = 0.0417e<sup>0.0002xPTU</sup> (n = 26, R<sup>2</sup> = 0.92\*\*)

The thermal indices during the reproductive stages particularly, during flowering and fruit set, varied across the seasons and thereby influences the thrips populations. Normally with the progress of reproductive phase in mango viz., flowering, panicle emergence, fruit set and development, hopper started shoots up based on exiting hydrothermal conditions. Application of thermal indices in this study confirmed that variations in thrips populations may be predicted >93 %. Thus, it was concluded that use of all these robust indices may be useful in assessing the pest-weather dynamics of region [48]. Based on all these information, region specific crop simulation dynamics models may be developed to predict and forecast the thrips population so that farmers can adopt control measures well in advance to save the fruit crop being lost.

**Acknowledgements**

Authors are thankful to the Director, CISH, Lucknow for providing facilities for carrying out the work. Thanks are also due to Indian Council of Agricultural

Research, New Delhi for providing financial assistance under National Initiative of Climate Resilient Agriculture (NICRA) Project entitled, "Understanding the changes in host pest interactions and dynamics in mango under climate change scenario", which facilitated carry out this study.

## References

1. Singh LB (1968) The Mango: botany, cultivation and utilization. World Crop Books, Leonard Hill, London, United Kingdom.
2. Tandon PL, Verghese A (1985) World List of insect, mite and other pests of mango. Technical Document, IHR, Bangalore, Karnataka, India.
3. Aliakbarpour H, Che-salmah M R, Dieng H (2010) Species composition and population dynamics of thrips (thysanoptera) in mango orchards of northern peninsular malaysia. *Environ Entomol* 39: 1409-1419.
4. Galán-saúco V (2009) El cultivo del mango. Segunda Edición. Mundi Prensa, Madrid, Spain.
5. Peña JE, Mohyuddin AI, Wysoky M (1998) A review of the pest management situation in mango agroecosystems. *Phytoparasitica* 26: 1-20.
6. Seal DR, Ciomperlik MA, Richards ML, Klassen W (2006) Distribution of chilli thrips, *scirtothrips dorsalis* (thysanoptera: thripidae), in pepper fields and pepper plants on st. vincent fla *Entomol* 89: 311- 320.
7. Reynaud P, Balmes V, Pizzol J (2008) Thrips hawaiiensis (morgan, 1913) (Thysanoptera: Thripidae), an asian pest thrips now established in europe. *Bulletin* 38: 155-160.
8. Verghese A, Tandon PL, Rao GSP (1988) Ecological studies relevant to the management of thrips palmi karny on mango in india. *Tropical Pest Mgt* 34: 55-58.
9. Kannan M, Rao NV (2006a) Seasonal incidence and population fluctuation of dipteran, isopteran, hymenopteran and thysanopteran pests of mango. *J PI Prot Envir* 3: 50-55.
10. Kannan M, Rao NV (2006b) Influence of age and stage of the host plant on insect pests of mango (*mangifera indica* L.). *Int J Agril Sci* 2: 351-353.
11. Ramasubbarao V, Thamiraju NB (1994) New record of blossom thrips *megalurothrips distalis* on mango (*mangifera indica*). *Ind J Agril Sci* 64: 417-418.
12. Kumar S, Bhatt RI (1999) Field evaluation of plant leaf extracts, oil and neem products against mango hopper (*amritodus atkinsoni* Lethierry) and thrips (*scirtothrips mangiferae* hood). *Allelopathy* J6: 271-276.
13. Kumar S, Nair AG, Bhatt RI (2002) Evaluation of promising and released mango hybrids for multiple pest resistance. *J Appl Zool Res* 13: 66-68.
14. Dahiya KK, Lakra RK (2001) Seasonal occurrence and succession of thrips, *rhipiphorothrips cruentatus* hood in important horticultural crops of haryana. *Crop Res (Hisar)* 21: 112-114.
15. Patel JR, Patel MB, Radadia GC, Shah AA, Pandya HV (1997) Insect pest management in mango nursery. *J Appl Hort (Navsari)* 3: 125-128.
16. Ananthkrishnan TN, Muraleedharan N (1974) On the incidence and effects of infestation of *selenothrips rubrocinctus* (giard) (thysanoptera: heliothripinae) on the free amino-acids of some susceptible host plants. *Curr Sci* 43: 216-218.
17. Krishnamoorthy A, Ganga, Visalakshi PN (2012) Record of thrips on mango. *J Hortl Sci* 7: 110-111.
18. Desneux N, Decourtye A, Delpuech JM (2007) The sub-lethal effects of pesticides on beneficial arthropods. *Annu Rev Entomol* 52: 81-106.
19. Horrigan L, Lawrence RS, Walker P (2002) How sustainable agriculture can address the environmental and human health harms of industrial agriculture. *Environ Health Perspect* 110: 445-456.
20. Adak T, Chakravarty NVK (2010) Quantifying the thermal heat requirement of *brassica* in assessing biophysical parameters under semi-arid microenvironments. *Int J Biometeorol* 54: 365-377.
21. Bazzocchi GG, Lanzoni A, Burgio G, Fiacconi MR (2003) Effects of temperature and host on the pre-imaginal development of the parasitoid *diglyphus isaea* (hymenoptera: eulophidae). *Biol Control* 26: 74-82.
22. Gilbert N, Raworth DA (1996) Insect and temperature, a general theory. *Can. Entomol* 128: 1-13.
23. Roy M, Brodeur J, Cloutier C (2002) Relationship between temperature and development rate of *stethorus punctillum* (coleoptera: coccinellidae) and its prey *tetranychus mcdanieli* (Acari: Tetranychidae). *Environ Entomol* 31: 177-187.
24. Jervis MA, Copland MJW (1996) The life cycle. Insect natural enemies-practical approaches to their study and evaluation. Chapman and Hall, London 63-161.
25. Wagner TL, Wu HI, Sharpe PJH, Schofield RM, Coulson RN (1984) Modelling insect development rates: A literature review and application of a biophysical model. *Ann Entomol Soc Am* 77: 208-225.
26. Whiley AW, Rasmussen TS, Wolstenholme BN, Saranah JB, Cull BW (1991) Interpretation of growth responses of some mango cultivars grown under controlled temperature. *Acta Horti* 291: 22-31.
27. Rajan S (2012) Phenological responses to temperature and rainfall: A case study of mango. Bioversity International, New Delhi, India.
28. Ravishankar H, Adak T, Singh VK, Pandey BK., Singh AK, et al. (2015) Empirical appraisal of some weather parameters' dynamics for their possible implications on mango production in some important mango growing regions with special reference to lucknow region of uttar pradesh., CRC Press (Apple Academic Press), Taylor & Francis Group, USA.
29. Aliakbarpour H, Rawi CS (2012) Seasonal abundance of *thrips hawaiiensis* (morgan) and *scirtothrips dorsalis* (Hood) (Thysanoptera: Thripidae) in mango orchards in malaysia. *Pertanika J Trop Agric Sci* 35: 637- 645.
30. Gupta MP, Nair MK, Srivastava AK (2009) Studies on seasonal activity of whitefly (*bemisia tabaci* genn) population and its association with weather parameters in bundelkhand zone of madhya Pradesh. *J Agrometeorol* 11: 175-179.
31. Sahu KR, Katlam BP, Chaudhary JL (2010) Impact of climatic factors on infestation of leaf eating caterpillar of cashew in chhattisgarh. *J Agrometeorol* 12: 105-107.
32. Dhawan AK, Aneja A, Singh J, Saini S (2009) Population dynamics of different pest on bt cotton *vis-a-vis* meteorological parameters in punjab. *J Agrometeorol* 11: 180-182.
33. Youngman RR, Toscano NC, Jones VP, Kido K, Natwick ET (1980) Correlations of seasonal trap counts of *bemisia tabaci* (homoptera: aleyrodidae) in southeastern california. *J Econ Entom* 79: 67-70.
34. Joshi, Kumar S (2012) Effect of some meteorological factors on seasonal abundance of *idioscopus nitidulus* (walker) (hemiptera: cicadellidae) in mango orchards of haridwar (India). *New York Sci. J* 5: 101- 103.
35. Kumari AD, Lakshmi BK., Reddy MGS, Reddy L (2009) Influence of abiotic factors on the incidence of hopper and chemical control strategies in mango. Karnataka. *J Agric Sci* 22: 601-602.
36. Shukla RP, Shukla PK, Varma S, Rajkumar B, Gundappa, et al. (2013) Population dynamics of major insect pests of mango and their natural enemies in relation to weather. Proceedings of International Conference on "Impact of Technological Tools on Food Security under Global Warming Scenario", Meerut, Uttar Pradesh, India.
37. Varshneya A, Rana KS (2008) Effect of some abiotic factors on population buildup of *idioscopus clypealis* (lethierry) in western uttar Pradesh. *J Environ Biol* 29: 811-812.
38. Ananthkrishnan TN (1993) Bionomics of thrips. *Ann Rev Entomol* 38: 1-92.
39. Kirk WDJ (1997) Distribution, abundance and population dynamics. Thrips as Crop Pests. CABI, Oxon, United Kingdom.
40. Duraimurugan P, Jagadish A (2002) Seasonal incidence and effect of weather parameters on the population dynamics of illi thrips *Scirtothrips dorsalis* Hood (Thysanoptera: Thripidae) on rose, Resources management in plant protection during twenty first century, Hyderabad, India: 180-183.
41. Shukla A. (2006) Seasonal activity of thrips, *scirtothrips dorsalis* on chilli crops. *Indian J of Tropical Biodiversity* 14: 177-174.
42. Panickar BK, Patel JR (2001) Population dynamics of different species of thrips on chilli, cotton and pigeonpea. *Indian J of Entomol* 63: 170-175.
43. Nandini, Giraddi RS, Mantur SM., Patil K., Mallapur CP, et al. (2010) Population dynamics and extent damage of pests of *capsicum* under protected cultivation. *Karnataka J Agril Sci* 25: 150-151.

44. Vanisree K., Rajashekar P., Rao R G. and Rao S V ( 2011) Seasonal incidence of thrips and its natural enemies on chilli (*capsicum annuum* L.) in andhra pradesh. *The Andhra Agril J* 58: 185-191.
45. Meena RS, Ameta OP, Meena BL (2013) Population dynamics of sucking pests and their correlation with weather parameters in chilli *capsicum annuum*. *Crop An In Quart. Js. of Life Sci: The Bioscan* 8: 177-180.
46. Jayakumar M, Rajavel M (2014) Weather based forewarning models for coffee borer incidence. *J Agrometeorol* 16: 140-143.
47. Adak T, Gundappa, Balaji R (2014) Application of thermal indices in predicting mango hopper dynamics in lucknow region of uttar pradesh. *Proceedings of International symposium on "New-dimensions in Agrometeorology for sustainable Agriculture"*. GBPUAT, Pantnagar, India.
48. Adak T, Singh VK, Ravishankar H (2014) Radiation dynamics and agroclimatic models as a tool to predict impact of climate change on dynamics of flowering in mango. *CISH, Lucknow, India*.

### Author Affiliations

[Top](#)

ICAR- Central Institute for Subtropical Horticulture, Rehmankhera, Lucknow, Uttar Pradesh, India

### Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 50 Journals
- ❖ 21 Day rapid review process
- ❖ 1000 Editorial team
- ❖ 2 Million readers
- ❖ Publication immediately after acceptance
- ❖ Quality and quick editorial, review processing

Submit your next manuscript at • [www.scitechnol.com/submission](http://www.scitechnol.com/submission)