



Biofilm Environmentome: A Survival Experiment

Juan Bueno*

Abstract

Microbial biofilms are communities of constant symbiosis and interaction in adaptation with their environment. This signaling network determines the chemistry as well as the physiology of the biofilm, it also confers protection to the microorganisms embedded in this formation, which improves the chances of survival in the face of external aggression. This mechanism known as Quorum Sensing (QS) responds to both intracellular and extracellular signals and modulates the response of the biofilm to the immune system. This chemical response to the environmental factors involved, together with the genetic factors of the genome in the development of an individual constitute the environmentome, it which represents a new source of promising therapeutic targets for the development of new treatments that prevent the emergence of antimicrobial resistance (AMR). The aim of this comment paper is briefly to explore the initial elements of an approach that can be applied in the development of new tools for clinical diagnosis and treatment, as well as in other areas such as ecology.

Keywords

Biofilm; Environmentome; Microbial pathogenesis; Biosensors

Introduction

Biofilms are an organized spatial structure, made up of several groups of microorganisms in exchange, these symbiotic relationships constitute a sociomicrobiological network that allow the evolution and survival of the organisms that persist inside [1]. In this order of ideas, biofilms contain a large number of microbial cells in different physiological states with a variety of information and expression, both genotypic and phenotypic, that modulates various metabolic pathways involved in adaptive mutation and selection responses to environmental agents [2]. In fact, it has been demonstrated that this formation helps cells to survive during exposure to toxic compounds as metals [3], it has also been described that the production of substances such as taurine, ectoine, and hydroxyectoine give protection against osmotic stress in the biofilms [4]. As a consequence, this resistance and tolerance that characterizes this microbial association is defined by the development of metabolic shifts as the primary response [5]. Likewise, as the biofilm interacts with the external environment, it is able to modulate the environment inside the structure in order to increase the persistence, in this way present two different internal domains, namely: an internal region with a low amount of oxygen that has dormant cells and an outer layer with cells in active metabolic

state and greater amount of oxygen, the presence of these two populations should always be taken into account for the design of any study and approach, because it represents the biofilm in a mature state [6]. Equally the production of microbial metabolites inside the biofilm microenvironments can alter the surrounding acidity by altering the microbiome and favoring tissue invasion [7], so this process can be considered as an adaptive phenomenon of microorganisms aimed at aggregation as a factor of survival and persistence [8].

Quorum sensing and environmentome

In this way, bacteria produce various compounds that modify their environment to increase their resilience, such is the case of *Pseudomonas aeruginosa* quorum sensing (PQS) system that use 2-alkyl-4(1H)-quinolone molecular family that modulates the production of toxins and the biofilm conformation [9]. These environmental changes modulated by QS are essential for survival and adaptation [10] and control various microbial processes as are: motility, biofilm structure, immune evasion, iron scavenging, cytotoxicity and antibiotic resistance [11].

Microbial glycomics

On the other hand microbial surface adhesion is an important pathogenic factor that favors cell invasion and lysis [12]. Adhesion is considered an important step in the establishment of biofilm and a need for microbial cells, which makes it an important factor of interaction with the environment for obtaining nutrients [13]. In addition, adhesion is determined by specific carbohydrate ligands lectin type with the ability to bind to glycans on human glycoproteins [14]. In this way the final attachment induced by microbial adhesins determines the production of the exopolysaccharide matrix that consolidates irreversible adhesion towards a mature formation that protects against external antimicrobial agents [15].

Ironfilm

There is evidence that the biofilm is not only a process of adaptation to the environment but a way of using environmental resources in favor of persistence, an example of this case is the iron, it have been observed that iron supplementation increase AMR to antimicrobial drugs tigecycline and tobramycin, also it has been shown that the absence of iron blocks the formation of the biofilm, these data suggest that the inhibition of iron metabolism may be an important therapeutic target for the development of new antibiotic drugs that prevent the development of AMR [16]. In this order of ideas two siderophores have been detected in *P. aeruginosa* (pyoverdine and pyochelin) which are determined by PQS [9].

Microbial fuels cell-based biosensor

This remarkable ability of biofilms to modulate the environment through a network of sociomicrobiological interactions can be applied as a powerful environmental monitoring tool that allows the detection of multiple pollutants, as the case of devices that include microbial fuel cell (MFC)-based biosensors with great advantages in selectivity and stability [17]. This technology have promising applications as are: bioremediation, biosensors, wastewater treatment and energy production [18]. Equally, MFC-sensors have several advantages between them: simplicity, results in a short period of time, wide range

*Corresponding author: Juan Bueno, Research Center of Bioprospecting and Biotechnology for Biodiversity Foundation (BIOLABB), Colombia, E-mail: juanbueno@biolabb.org, juangbueno@gmail.com

Received: January 01, 2018 Accepted: January 29, 2018 Published: February 05, 2018

of measurements and the possibility of conducting studies in the field; but the materials for its manufacture are expensive [19].

In this way the study of biofilm environmentome is an exciting research field where is possible to integrate multiple disciplines to obtain applicable knowledge that leads to innovation and sustainable development.

References

1. Hansen SK, Rainey PB, Haagenen JA, Molin S (2007) Evolution of species interactions in a biofilm community. *Nat* 445: 533-536.
2. Stewart PS, Franklin MJ (2008) Physiological heterogeneity in biofilms. *Nat Rev Microbiol* 6: 199-210.
3. Harrison JJ, Ceri H, Turner RJ (2007) Multimetal resistance and tolerance in microbial biofilms. *Nat Rev Microbiol* 5: 928-938.
4. Mosier AC, Justice NB, Bowen BP, Baran R, Thomas BC, et al. (2013) Metabolites associated with adaptation of microorganisms to an acidophilic, metal-rich environment identified by stable-isotope-enabled metabolomics. *mBio* 4:1-8.
5. Booth SC, Workentine ML, Wen J, Shaykhtudinov R, HJ Vogel, et al. (2011) Differences in metabolism between the biofilm and planktonic response to metal stress. *J Proteome Res* 10: 3190-3199.
6. Rahman T, Yarnall B, Doyle DA (2017) Efflux drug transporters at the forefront of antimicrobial resistance. *Eur Biophys J* 46: 647-653.
7. Takahashi N, Washio J, Mayanagi G (2012) Metabolomic approach to oral biofilm characterization—a future direction of biofilm research. *J Oral Biosci* 54: 138-143.
8. Hope EA, Amorosi CJ, Miller AW, Dang K, CS Heil, et al. (2017) Experimental evolution reveals favored adaptive routes to cell aggregation in yeast. *Genetics* 206: 1153-1167.
9. Popat R, Harrison F, Da Silva AC, Easton SA, McNally L, et al. (2017) Environmental modification via a quorum sensing molecule influences the social landscape of siderophore production. *Proc Biol Sci* 284:1-18.
10. Hawver LA, Jung SA, Ng WL (2016) Specificity and complexity in bacterial quorum-sensing systems. *FEMS Microbiol Rev* 40: 738-752.
11. Lee J, Zhang L (2015) The hierarchy quorum sensing network in *Pseudomonas aeruginosa*. *Protein & cell* 6: 26-41.
12. Reid CW, Fulton KM, Twine SM (2010) Never take candy from a stranger: the role of the bacterial glycome in host-pathogen interactions. *Future Microbiol* 5: 267-288.
13. Zhang B, Powers R (2012) Analysis of bacterial biofilms using NMR-based metabolomics. *Future Med Chem* 4: 1273-1306.
14. Lipke PN (2016) Glycomics for Microbes and Microbiologists. *mBio* 7: 1-3.
15. Khelissa SO, Abdallah M, Jama C, Faille C, Chihib NE, et al. (2017) Bacterial contamination and biofilm formation on abiotic surfaces and strategies to overcome their persistence. *J Mater Environ Sci* 8: 3326-3346.
16. Oglesby-Sherrouse AG, Djapgne L, Nguyen AT, Vasil AI, Vasil, ML et al. (2014) The complex interplay of iron, biofilm formation, and mucoidy affecting antimicrobial resistance of *Pseudomonas aeruginosa*. *Pathog Dis* 70: 307-320.
17. Zhou T, Han H, Liu P, Xiong J, Tian F et al. (2017) Microbial Fuels Cell-Based Biosensor for Toxicity Detection: A Review. *Sensors* 17: 2230.
18. Logroño W, Guambo A, Pérez M, Kadier A, Recalde C, et al. (2016) A terrestrial single chamber microbial fuel cell-based biosensor for biochemical oxygen demand of synthetic rice washed wastewater. *Sensors* 16: 101.
19. Chouler J, Bentley I, Vaz F, O'Fee A, Cameron, PJ, et al. (2017) Exploring the use of cost-effective membrane materials for Microbial Fuel Cell based sensors. *Electrochim Acta* 231: 319-326.

Author Affiliations

Top

Research Center of Bioprospecting and Biotechnology for Biodiversity Foundation (BIOLABB), Colombia

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission