

Journal of Addictive Behaviors, Therapy & Rehabilitation

A SCITECHNOL JOURNAL

Research Article

Cannabis Associated Elation of Manic Symptoms in a University Cohort (Non-Clinical)

Muhammad T Khalily^{1,2*}, Matthias Schwannauer¹ and Brian Hallahan³

Abstract

Cannabis is the most frequently used drug worldwide and is a wide spread problem among the young adults. The use of cannabis in some countries such as Pakistan is a traditional habit for the elation of mood and the users are not aware of the link between cannabis use and manic symptoms. The present study aims to investigate the presence of manic symptoms in non-clinical population particularly adolescents and young adults, who use cannabis in a social context and did not report psychiatric illness/ or have no previous psychiatric history. For this purpose a cross sectional design study sample which comprised of non-clinical 100 participants from a general population ,which include 50 individuals using Cannabis for social purpose and with no psychiatric diagnosis, and an equal number of non-cannabis users without cannabis use history and no psychiatric diagnosis were considered for this study. Our results indicated a positive correlation (p<.01) between cannabis use and the presence of manic symptoms and subsequently demonstrate association between cannabis use and manic symptoms. Furthermore our results indicated a significant difference (p<.01) between cannabis users and non-cannabis users on the basis of manic symptoms and impulsiveness. To conclude our findings highlighted the prevalence of manic symptoms in individuals who ingest cannabis in the social context but did not report other psychiatric disorders.

Keywords

Cannabis; Pakistan; Psychosis; Elation; Depression

Introduction

Cannabis products are the most commonly abused drugs on the illicit drug market worldwide [1-3], with increasing use demonstrated in most western countries over the last two decades, particularly in young adults [4-6]. A recent national survey conducted by United Nations Office on Drug and Crime (UNODC) in 2013 [7] indicated that cannabis is the most popular illegal psycho-active substance in Pakistan and is ingested by approximately four million people (3.6% of the adult population), with use particularly prevalent in younger individuals [7]. Data relating to trends in cannabis use in Pakistan are unavailable, given the lack of data prior to 2013, although given its low price and easy availability, it was previously thought to be

Received: June 14, 2015 Accepted: October 19, 2015 Published: October 24, 2015



All articles published in Journal of Addictive Behaviors, Therapy & Rehabilitation are the property of SciTechnol, and is protected by copyright laws. Copyright © 2015, SciTechnol, All Rights Reserved.

the most commonly abused illegal psycho-active substance [7,8]. Cannabis use in Pakistan is markedly more prevalent amongst men and in those who have attained lower levels of education although use and abuse occurs across both genders and all socio-economic classes [7]. The possession, exportation, importation, trading and cultivation of cannabis products are illegal in Pakistan, with no official statistics on cannabis cultivation, production and eradication available to date. However, it is known that large amounts of cannabis are transported into Pakistan from Afghanistan [7]. The pattern of cannabis in Pakistan is predominantly by inhalation of hashish (Charas) but also occurs from consumption of "bhang", a drink made from cannabis leaves.

There are multiple motives for young individuals to use cannabis in Pakistan. These include peer influence, using cannabis as a coping strategy secondary to psycho-social stressors, and using cannabis to experience its euphoric inducing effects [7]. Euphoric feelings secondary to cannabis ingestion in Pakistan include anecdotal reports such as "elation of the king" [9]. In addition, the consumption of "bhang" has previously been associated with symptoms of excitement, grandiosity and psychosis [10]. International studies have also suggested that cannabis is ingested predominantly for its euphoric inducing effects', its facilitation of social interactions and as a coping mechanism secondary to psycho-social stressors [11-13]. The use of cannabis for its induction of euphoria has also previously been noted in Iran where a "solid pie" called "Majoon Birjandi" is ingested to induce euphoria and make individuals "feel special" [14].

There is a burgeoning evidence base suggesting that cannabis use contributes to the development of psychosis and is associated with a poorer prognosis for those with a pre-existing vulnerability to psychosis [15-17]. Emerging evidence also suggests that cannabis use or dependence may be associated with inducing manic or depressive episodes in individuals both with and without an established bipolar disorder [18-22]. In addition, cannabis use has been associated with longer durations of acute episodes (including manic episodes) and increased relapse rates for individuals with bipolar disorder [23-25]. However, cannabis (and other psycho-active substances) has also been utilized by individuals to self-medicate in relation to manic or depressive symptoms [18] and consequently disentangling the mood altering effects of cannabis from non-psychoactive substance induced mood disorder can be problematic.

Consequently, there is an increasing evidence of a putative relationship between cannabis misuse and its effect on mood and mood disorders. In this study, we investigate in a young university student population in Pakistan, any putative association between recreational cannabis use (non-dependent on cannabis) and the production of hypomania or manic symptoms compared to a matched non-cannabis using university student population.

Materials and Methods

Participants

Individuals were recruited by advertisement (Facebook, e-mail platforms; posters in computer laboratories) from Universities in Islamabad. All individuals provided informed consent and

^{*}Corresponding author: Prof. Muhammad T Khalily, Post Doc Fellow, Clinical and Health Psychology School of Health in Social Sciences, The University of Edinburgh and Professor Department of Psychology International Islamic University Islamabad, Tel: 00923475710446; E-mail: khalily64@gmail.com

ethical approval was attained from the Department of Psychology International Islamic University Islamabad and the School of Health in Social Sciences University of Edinburgh UK. Exclusion criteria comprised individuals actively attending the mental health services, a past Axis I major mental illness, a first degree relative with a major affective disorder (e.g. bipolar disorder, recurrent depressive disorder, schizoaffective disorder), or a history of harmful use or dependence on alcohol, opiates or other (non-cannabis) illegal psycho-active substances. Participants included 50 individuals with low to moderate cannabis use (scoring 1 or 2 on Cannabis Abuse screening Test [26] and 50 non-cannabis users (score of 0 on CAST). As this study investigated students without a history of harmful use or dependence on cannabis, we excluded individuals who scored \geq 3 on the CAST. We similarly excluded individuals who harmfully used or were dependent on alcohol (a score \geq 3 on the Fast Alcohol Screening Test [27].

Demographic and clinical data

Demographic and clinical data collected included age, gender, employment, residential and marital status. Clinical data included past psychiatric and family psychiatric history and a history of psycho-active substance use. Psychometric scales were utilized to measure cannabis (CAST) and alcohol use (FAST) and evaluated the social context of cannabis [28] and alcohol usage [29]. In addition, we also measured symptoms of elation (Internal State Scale; ISS [30], impulsivity (Barratt Impulsivity Scale II; BIS-II [31]; Patton, Stanford, & Barratt, 1995) and the presence of distress or psychological symptoms across nine symptom dimensions [32,33]. A brief description of these instruments is provided below.

Psychometric instruments

Cannabis Abuse screening Test (CAST)

The CAST is a 6 item self-administered scale utilized to screen problematic patterns of cannabis use (range 0-4 on each item) and is has demonstrated high internal consistency including in young populations (cronbach's α =0.72-0.92) [26,34,35]. A score ≥3 indicates harmful use or dependence.

Fast Alcohol Screening Test (FAST)

The FAST is a 4-item self-report screening test (range 0-4 on each item), derived from the Alcohol Use Disorders Identification Test (AUDIT) to detect alcohol misuse and dependence and is associated with good reliability (cronbach's alpha=0.77, [36]. and sensitivity indices of >90% [27].

Social Context of Cannabis Use Scale (SCCUS)

The SCCUS is a 24 item self-report scale that assesses the reasons why and the variety of situations where cannabis is used. It includes four sub-scales: social facilitation, peer acceptance, emotional pain and sex seeking with good internal consistency demonstrated for all sub-scales, including in a student university population (cronbach's α 's=0.71 to 0.95, [12].

Social Context of Drinking Scale (SCDS)

The SCDS is a 30 item self-report scale that quantifies the frequency of alcohol use in 6 different contexts and is associated with good internal consistency (cronbach's α =0.76-0.93) [29,37].

Internal State Scale (ISS)

The ISS is a 15 item self-report 100 point visual analogue scale quantifying participants experiences of their mood state over the previous 24 hours. It consists of four subscales: Activation (ACT), Perceived Conflict (PC), Well-Being (WB) and the Depression Index (DI). The ACT and WB sub-scales are validated for discriminating mood states and the ACT and DI correlate with clinical measures of mania and depression. The ISS has good internal consistency (Cronbach's a's of 0.81 to 0.92) for the various sub scales, [30] and good inter-rater reliability (κ =0.62-0.94; [30,38] in a wide variety of populations [39], although lower sensitivity has been noted in inpatient settings in patients with acute mania [40].

Brief Symptom Inventory (BSI)

The BSI is a 53 item self-report inventory derived from the SCL-90-R [41]. and was developed for use in a wide variety of settings. It encompasses 9 primary symptom dimensions: somatisation (SOM), obsessive compulsive (O-C), interpersonal sensitivity (I-S), depression (DEP), anxiety(ANX), hostility (HOS), phobic anxiety (PHOB), paranoid ideation (PAR), and psychoticism (PSY) [32,42]. It also includes three global indices of distress, the General Severity Index (GSI), the Positive Symptom Distress Index (PSDI) and the Positive Symptom Total (PST). All subscales of the BSI are associated with good internal consistency ranging from 0.75 to 0.89 [33].

Barratt Impulsiveness Scale-II (BIS)

The BIS is a 34-item self-report questionnaire with three subscales measuring attentional motor and non-planning impulsivity. It is associated with good internal consistency (cronbach's $\alpha = 0.79$ to 0.83) and has been used across several populations including university students [31]. Anonymised data was collected electronically [43] and all individuals were instructed to complete psychometric scales and demographic data when they had not ingested alcohol or cannabis for at least 48 hours.

Statistical analysis

Statistical analysis was performed using the Statistical Package for Social Sciences 20.0 for Windows (SPSS Inc., IBM, New York, USA). We utilized the student t-test for parametric data and the Chi Square (χ^2) test for non-parametric categorical data where appropriate.

Results

Demographic and clinical data

Out of 153 respondents, 50 individuals fulfilled inclusion and exclusion criteria for the cannabis user group (CAST score of 1 or 2) and 48 individuals fulfilled criteria for the non-cannabis using group (CAST=0). Five individuals were excluded due to a CAST score \geq 3 and 48 individuals were excluded due to a FAST score of \geq 3. Demographic data are presented in Table 1; with no differences noted between the groups in age, gender, or other socio-demographic indices measured.

Cannabis use and clinical symptoms

In addition to higher scores on the CAST and SCCUS (Table 2), the cannabis using group had marginally higher scores on the FAST and SCDS (relating to use of alcohol in social contexts). Higher total scores on the ISS (p<0.001) and higher sub-scale scores (particularly the ACT and WC), (Table 2) were present in the cannabis-using

doi:http://dx.doi.org/10.4172/2324-9005.1000144

group. Higher motor (p=0.016) and attentional impulsivity (p=0.012), and higher scores on the BSI (p=0.043) and a number of subscales

Table 1: Demographic Data.						
Variables	Cannabis Users	Non-Cannabis Users	Statistics			
variables	F (%)	F (%)	χ², df	р		
Age (years)						
16-20	8 (34.8)	15 (62.2)	4 66 3	0 104		
21-24	26 (59.1)	18 (40.9)	4.00, 5	0.134		
25-30	10 (43.5)	13 (56.5)				
31.34	6 (60.0)	4 (40.0)				
Gender			0.55 1	0.112		
Male	30 (60)	14 (28)	2.55, 1	0.112		
Female	2 (4)	4 (8)				
Education Completed						
Secondary School	5 (10)	1 (2)	11.20, 2	0.004		
Undergraduate Degree	28 (56)	16 (32)				
Higher Degree	17 (34)	34 (66)				
Employment Status						
Employed	10 (20)	15 (30)	0.48, 3	0.92		
Unemployed	2 (4)	2 (4)	,			
Student	38 (76)	33 (66)				
Residential Status						
Urban	31 (62)	26 (52)	3.30, 2	0.129		
Sub-urban	6 (12)	11 (22)				
Rural	13 (26)	13 (26)				
Marital Status						
Single	45 (90)	38 (76)	4.05.0	0.226		
Married	4 (8)	11 (22)	4.35, 3			
Separated	1 (2)	1 (2)				

p < .05, **p < .01; CAST = Cannabis Abuse screening Test, FAST = Screening for alcohol problem, SCCUS = Social Context of Cannabis Use Scale, SCDS = Social Context of Drinking Scale, ISS = Internal State Scale, ACT = Activation, OC = Personal Conflict, WB = Well being, DI = Depression Index, BIS = Barratt Impulsiveness Scale-II, BSI = Brief Symptom Inventory, SOM = Somatization, O-C = Obsessive Compulsive, I-S = Interpersonal Sensitivity, DEP = Depression, ANX = Anxiety, HOS = Hostility, PHOB = Phobic Anxiety, PAR =Paranoid Ideation, PSY = Psychoticism

Table 2:	Psychometric	Measures.
----------	--------------	-----------

	Cannabis Users (n=50) M (SD)	Non-Cannabis Users		Statistics			
Variables		(n=48) M (SD)	Т	р	Cohen's d		
CAST	8.42 (5.60)	.58 (0.06)	10.13	<0.001	1.72		
FAST	2.90 (2.52)	1.00 (0.34)	5.36	<0.001	1.06		
SCCUS	34.17 (11.56)	2.44 (0.76)	27.51	<0.001	3.90		
SCDS	19.65 (14.69)	6.11 (1.6)	5.69	<0.001	1.30		
ISS				<0.001			
Total	52.65 (29.39)	21.94 (20.44)	6.59	<0.001	1.25		
ACT	18.60 (11.07)	8.64 (6.75)	6.36	<0.001	1.20		
PC	13.67 (7.13)	7.66 (6.77)	5.78	<0.001	1.11		
WT	15.51 (10.92)	7.00 (5.38)	5.15	<0.001	0.94		
DI	4.85 (4.65)	2.48 (1.53)	4.60	<0.001	0.90		
BIS							
Total	48.97 (16.53)	45.90 (19.80)	0.88	0.378	0.16		
Motor	13.23 (5.04)	10.73 (5.55)	2.45	0.016	0.47		
Non-planning	17.69 (7.89)	20.73 (8.69)	1.89	0.060	0.36		
Attentional	18.04 (6.95)	14.43 (7.72)	2.54	0.012	0.49		
BSI							
Total	35.57 (31.46)	32.62 (22.55)	2.04	0.043	0.10		
SOM	4.13 (3.98)	3.75 (2.37)	2.02	0.046	0.11		
O-C	4.39 (3.95)	4.66 (3.15)	1.84	0.142	0.07		
I-S	4.30 (4.90)	3.99 (2.67)	2.02	0.046	0.09		
DEP	5.02 (4.58)	4.38 (3.07)	2.17	0.032	0.16		
ANX	4.29 (3.81)	3.91 (2.80)	1.94	0.054	0.11		
HOS	3.14 (3.05)	3.60 (2.42)	1.10	0.270	0.16		
PHOB	3.48 (3.42)	3.42 (2.20)	1.88	0.063	0.01		
PAR	4.50 (4.32)	3.87 (2.60)	2.28	0.024	0.17		
PSY	5.05 (4.92)	4.77 (2.92)	2.02	0.045	0.06		

p < .05, ***p* < .01;

CAST = Cannabis Abuse screening Test, FAST = Screening for alcohol problem, SCCUS = Social Context of Cannabis Use Scale, SCDS = Social Context of Drinking Scale, ISS = Internal State Scale, BIS = Barratt Impulsiveness Scale-II, BSI = Brief Symptom Inventorys

Table 3: Correlation between psychometric instrumen	s in cannabis and n	on-cannabis using groups.
---	---------------------	---------------------------

	Cannabis users (n = 50)					Non-Cannabis users (n = 48)				
	CAST	FAST	SCCUS	SCDS	BSI	CAST	FAST	SCCUS	SCDS	BSI
CAST	-	-0.10	-0.12	-0.10	-0.07	-	-0.08	-0.05	-0.04	-0.06
FAST		-	-0.12	0.29 [*]	-0.17		-	0.21	-0.09	0.28
ISS										
Total	0.07	-0.34 [*]	0.35⁺	-0.47**	-0.36*	-0.01	0.25	0.23	0.14	0.34 [*]
ACT	0.11	-0.34 [*]	0.33 ⁻	-0.52**	-0.43**	0.08	0.15	0.24	0.17	0.28
PC	0.02	-0.32 [*]	0.34 [*]	-0.37*	-0.33*	0.02	0.36*	0.20	0.01	0.40 [*]
WB	0.07	-0.25	0.27	-0.45**	-0.40 [*]	-0.12	0.09	0.12	0.12	0.17
DI	0.00	-0.26	0.29	-0.32*	-0.03	-0.05	0.43**	0.33*	0.28	0.34 [*]
SCCUS	-0.12	-0.12	-	0.08	-0.16	-0.05	0.21	-	-0.05	-0.11
SCDS	-0.10	0.29 [*]	0.08	-	0.33 [*]	-0.04	-0.09	-0.05	-	-0.06
BIS										
Total	0.03	0.00	-0.16	-0.05	0.22	0.11	-0.07	0.11	-0.19	0.26
Motor	0.03	-0.14	0.04	-0.27	-0.14	0.02	0.00	0.13	-0.07	0.32
Non-planning	-0.01	0.22	-0.30	0.21	0.44**	0.15	-0.19	0.07	-0.22	0.03
Attentional	0.07	-0.17	-0.08	-0.12	0.22	0.12	0.01	0.09	-0.22	0.42**
BSI	-0.07	-0.17	-0.16	0.33 [*]	-	0.69	0.08	0.48	0.70	-

particularly PAR (p=0.024) was found in the cannabis using group.

In Table 3, we present correlation analysis between the psychometric measures utilized in this study. In the cannabis using group, the highest correlations (r>0.4, p<0.01) were noted between the ACT and WB sub-scales of the ISS and the total BSI score, between the ACT and WB subscales of the ISS and total SCDS scale and between non-planning impulsivity and the BSI scale. For non-cannabis users, the highest correlation was between the DI subcomponent of the BSA and the FAST, with other correlations (r>0.4, p<0.01) noted between the PC sub-scale of the ISS and the BSI and attentional impulsivity and the BSI.

Discussion

In this study, university students without a history of major mental illness or substance dependence (alcohol or illegal psychoactive substances) who were recreational cannabis users demonstrated increased scores on measures of psychological well-being and in particular in relation to measures of elated mood. These symptoms were present in a cohort of students who previously had no reported psychopathology and who had never sought or attained support for any mental health related difficulty.

Although, not extensively investigated, our findings of cannabis induced mood elation are consistent with some previous research [22]. Whilst all sub-scales of the ISS were increased in the cannabis-using group, those pertaining to mood elation (ACT, WB) demonstrated the largest differences between the groups, with these sub-scales also correlated with total BSI score (the presence of psychological symptoms). Overall scores on the BSI were relatively low in both groups with a very minimal total score increase in the cannabisusing group compared to the non-cannabis using group. This potentially suggests that the most prominent symptoms experienced by the cannabis using group related to symptoms of elation, a finding consistent with some previous research [19]. The cannabis using group demonstrated higher levels of motor and attentional impulsivity, a finding consistent with previous research [25]. Such impulsivity relates to impaired impulse control and is potentially associated with this group engaging in cannabis mis-use [44,45].

Limitations

There are a number of limitations with this study. Firstly, measures of functionality and in-depth measures relating to educational achievement in university were not collected, and consequently a longitudinal follow-up study to ascertain the link between cannabis use and university performance and functionality will be undertaken. Secondly, we only included a homogenous group of individuals who recreationally ingested cannabis or consumed alcohol, rather than individuals who harmfully used or were dependent on these substances. This reduces the generalizability of our findings to nonrecreational users. Thirdly, there was a minimal, albeit significant difference in the alcohol consumption between the two groups as measured on the FAST and SCDS which potentially could explain differences between the groups on the ISS. A re-analysis (ANCOVA) controlling for these factors did not however significantly alter our findings. Indeed, in the non-cannabis using group a correlation between the depression index (rather than the ACT or WB sub-scales) of the ISS and alcohol use (FAST) was noted. Finally, we cannot outrule that a proportion of the students investigated were intoxicated on cannabis when completing this study and that this may have affected the accuracy of their results. However, it is probable that this could have affected several psychometric instruments and confounds more than just the findings related to elation and impulsivity.

Conclusion

Cannabis use was demonstrated to be associated with symptoms of elation and impulsivity in a university student population who had no prior history of treatment by the mental health services. A greater awareness of the link between cannabis use and symptoms of elation and potentially bipolar disorder by clinicians in suggested.

References

 Boydell J, van Os J, Caspi A, Kennedy N, Giouroukou E, et al. (2006) Trends in cannabis use prior to first presentation with schizophrenia, in South-East London between 1965 and 1999. Psychol Med 36: 1441-1446.

- Sequoia Ecosystem and Recreation Preserve Act of 1999 (1999) H.R. 2077, US House of Representatives.
- 3. Macleod J (2007) Cannabis use and psychosis: the origins and implications of an association. Adv Psychiatr Treat 13: 400-411.
- 4. EMCDDA: Annual report (2005) The State of the Drugs problems in the European Union and Norway. Eu Publications, Luxembourg.
- Hall W, Degenhardt L (2007) Prevalence and correlates of cannabis use in developed and developing countries. Curr Opin Psychiatry 20: 393-397.
- SAMHSA (Substance Abuse and Mental Helath Services Administration) (2013) Results from the 2012 National Survey on Drug Use and Health: Summary of National Findings. HHS, Rockville, MD, USA.
- 7. Drug Use in Pakistan (2013) United Nations Office on Drugs and Crime (UNODC). Islamabad, Pakistan.
- Illicit Drug Trends in Pakistan (2008) United Nations Office on Drugs and Crime (UNODC) Islamabad, Pakistan.
- 9. Khalily MT (2010) Developing a coordinated response to drug abuse in Pakistan. J Interprof Care 24: 168-172.
- Chaudry HR, Moss HB, Bashir A, Suliman T (1991) Cannabis psychosis following bhang ingestion. Br J Addict 86: 1075-1081.
- Simons JF (1998) Cell wall 1,6-beta-glucan synthesis in Saccharomyces cerevisiae depends on ER glucosidases I and II, and the molecular chaperone BiP/Kar2p. EMBO J 17: 396-405.
- Beck KH, Caldeira KM, Vincent KB, O'Grady KE, Wish ED, et al. (2009) The social context of cannabis use: relationship to cannabis use disorders and depressive symptoms among college students. Addict Behav 34: 764-768.
- Buckner JD, Shah SM, Dean KE, Zvolensky MJ (2015) Cannabis use frequency and use-related impairment among African-American and White users: the impact of cannabis use motives. Ethn Health.
- Mehrpour O, Karrari P, Afshari R (2012) Recreational use and overdose of ingested processed cannabis (Majoon Birjandi) in the eastern Iran. Hum Exp Toxicol 31: 1188-1189.
- Arseneault L, Cannon M, Witton J, Murray RM (2004) Causal association between cannabis and psychosis: examination of the evidence. Br J Psychiatry 184: 110-117.
- Smit F, Bolier L, Cuijpers P (2004) Cannabis use and the risk of later schizophrenia: a review. Addiction 99: 425-430.
- Rehman IU, Farooq S (2007) Schizophrenia and comorbid self reported cannabis abuse: impact on course, functioning and services use. J Pak Med Assoc 57: 60-64.
- Strakowski SM, DelBello MP, Fleck DE, Arndt S (2000) The impact of substance abuse on the course of bipolar disorder. Biol Psychiatry 48: 477-485.
- Henquet C, Krabbendam L, De Graaf R, Ten HM, Van OSJ (2006) Cannabis use and expression of mania in the general population. J Affect Disord 95: 103-110.
- Caldeira KM, Arria AM, O'Grady KE, Vincent KB, Wish ED (2008) The occurrence of cannabis use disorders and other cannabis-related problems among first-year college students. Addict Behav 33: 397-411.
- 21. van Rossum I, Boomsma M, Tenback D, Reed C, van Os J; EMBLEM Advisory Board (2009) Does cannabis use affect treatment outcome in bipolar disorder? A longitudinal analysis. J Nerv Ment Dis 197: 35-40.
- 22. Gibbs M, Winsper C, Marwaha S, Gilbert E, Broome M, et al. (2015) Cannabis use and mania symptoms: a systematic review and meta-analysis. J Affect Disord 171: 39-47.
- Arendt M, Munk-Jorgensen P (2004) Heavy cannabis users seeking treatment- prevalence of psychiatric disorders. Soc Psychiatry Psychiatr Epidemiol 39: 97-105.
- Isaac M, Isaac M, Holloway F (2005) Is cannabis an anti-antipsychotic? The experience in psychiatric intensive care. Hum Psychopharmacol 20: 207-210.
- 25. Sundram S (2006) Cannabis and neurodevelopment: implications for psychiatric disorders. Hum Psychopharmacol 21: 245-254.
- 26. Legleye S, Piontek D, Kraus L, Morand E, Falissard B (2007) Validation of

doi:http://dx.doi.org/10.4172/2324-9005.1000144

the CAST, a general population Cannabis Abuse Screening Test. J Subst Use 12: 233-242.

- Hodgson R, Alwyn T, John B, Thom B, Smith A (2002) The FAST Alcohol Screening Test. Alcohol Alcohol 37: 61-66.
- Beck KH, Thombs DL, Summons TG (1993) The social context of drinking scales: construct validation and relationship to indicants of abuse in an adolescent population. Addict Behav 18: 159-169.
- Perkins HW, Haines MP, Rice R (2005) Misperceiving the college drinking norm and related problems: a nationwide study of exposure to prevention information, perceived norms and student alcohol misuse. J Stud Alcohol 66: 470-478.
- Bauer MS, Crits-Christoph P, Ball WA, Dewees E, McAllister T, et al. (1991) Independent assessment of manic and depressive symptoms by self-rating. Scale characteristics and implications for the study of mania. Arch Gen Psychiatry 48: 807-812.
- Patton JH, Stanford MS, Barratt ES (1995) Factor structure of the Barratt impulsiveness scale. J Clin Psychol 51: 768-774.
- Derogatis LR, Melisaratos N (1983) The Brief Symptom Inventory: an introductory report. Psychol Med 13: 595-605.
- Meachen SJ, Hanks RA, Millis SR, Rapport LJ (2008) The reliability and validity of the brief symptom inventory-18 in persons with traumatic brain injury. Arch Phys Med Rehabil 89: 958-965.
- Legleye S, Piontek D, Kraus L (2011) Psychometric properties of the Cannabis Abuse Screening Test (CAST) in a French sample of adolescents. Drug Alcohol Depend 113: 229-235.
- Piontek D, Kraus L, Klempova D (2008) Short scales to assess cannabisrelated problems: a review of psychometric properties. Subst Abuse Treat Prev Policy 3: 3-25.
- Meneses-Gaya C, Crippa JA, Zuardi AW, Loureiro SR, Hallak JE, et al. (2010) The fast alcohol screening test (FAST) is as good as the AUDIT to screen alcohol use disorders. Subst Use Misuse 45: 1542-1557.
- Thombs DL, Beck KH, Mahoney CA (1993) Effects of social context and gender on drinking patterns of young adults. J Couns Psychol 40: 115-111.
- Bauer MS, Vojta C, Kinosian B, Altshuler L, Glick H (2000) The Internal State Scale: replication of its discriminating abilities in a multisite, public sector sample. Bipolar Disord 2: 340-346.
- 39. Bauer MS (2008) The Internal State Scale (ISS): Version 2 and the ChronoBook, Scoring key and Summary, USA.
- Altman E, Hedeker D, Peterson JL, Davis JM (2001) A comparative evaluation of three self-rating scales for acute mania. Biol Psychiatry 50: 468-471.
- Derogatis LR, Rickels K, Rock AF (1976) The SCL-90 and the MMPI: a step in the validation of a new self-report scale. Br J Psychiatry 128: 280-289.
- Johnson ME, Brems C, Mills ME, Fisher DG (2007). Psychiatric symptomatology among individuals in alcohol detoxification treatment. Addict Behav 32: 1745-1752.
- 43. Schwannauer M, Noble A, Fraser G (2011) Behavioural risk of bipolar disorder in an analogue population: the role of cognitive, developmental and interpersonal factors. Clin Psychol Psychother 18: 411-417.
- 44. Patock-Peckham JA, King KM, Morgan-Lopez AA, Ulloa EC, Filson Moses JM (2011) The gender specific mediational links between parenting styles, parental monitoring, impulsiveness, drinking control, and alcohol-related problems. J Stud Alcohol Drugs 72: 247–258.
- Potenza MN, de Wit H (2010) Control yourself: alcohol and impulsivity. Alcohol Clin Exp Res 34: 1303-1305.

Author Affiliations

¹Department Clinical & Health Psychology School of Health in Social Sciences, University of Edinburgh, UK

²Department of Psychology, International Islamic University Islamabad ³School of Medicine, Clinical Science Institute, National University of Ireland Galway, Galway, Ireland, Department of Psychiatry, University Hospital Galway, Galway Roscommon Mental Health Services, Ireland

Тор