



Research Article

# Effect of Aqueous Extract of *Protium heptaphyllum* (Burseraceae) on *Lutzomyia longipalpis* (Diptera: Psychodidae), a Proven Vector of Visceral *Leishmaniasis* in Brazil

Yrllan Ribeiro Sincurá<sup>1</sup>, Amanda Oliveira Baracho<sup>1</sup>, Cristiane Fernanda Fuzer Graef<sup>2</sup>, Emerson Cotta Bodevan<sup>3</sup> and Ricardo Andrade Barata<sup>1\*</sup>

## Abstract

**Objective:** To evaluate the toxicity of aqueous extract of *Protium heptaphyllum* on *Lutzomyia longipalpis* a proven vector of visceral leishmaniasis (VL).

**Methods:** Aqueous extract of the leaves of *P. heptaphyllum* were prepared at 0.25, 0.10, 0.05 and 0.025 mg/mL concentrations, and 100 µL was applied to adults of *Lu. longipalpis* (n=20). Controls included 20 adults that were treated with 100 µL distilled water (negative control) and 20 adults that were treated with 100 µL alpha-cypermethrin at 196 µg/mL concentration (positive control). The percentage of dead insects in each group was observed at 1, 2, 4, 8, 12, 24, 48 and 72h after application.

**Results:** The mortality rate of *Lu. longipalpis* after 2h of exposure to the extract at a concentration of 0.1 mg/mL was 52%, behaving similarly to the positive controls (cypermethrin), and the extract at a concentration of 0.25 mg/ml was able to kill 56% of insects within the first hour of exposure. After 72h, the aqueous extract of *P. heptaphyllum* showed a mortality rate of about 81% and 83%, at concentrations of 0.1 and 0.25 mg/mL, respectively.

**Conclusion:** The extract of *P. heptaphyllum* contains terpenes, a molecule with recognized toxicity in insects, and which may be responsible for killing the phlebotomines. These results demonstrate the potential use of this extract in the control of this vector.

## Keywords

Phlebotominae; Sand flies; Insecticidal activity; Bioinsecticide; Vector control

## Introduction

Visceral leishmaniasis (VL) is an infectious parasitic disease that mainly affects hematopoietic organs such as liver, spleen and bone

\*Corresponding author: Ricardo Andrade Barata, Universidade Federal dos Vales do Jequitinhonha e Mucuri, Laboratório de Parasitologia, Departamento de Ciências Biológicas, Campus JK, Rodovia MGT-367, Km 583, 5000, Alto da Jacuba, 39100-000, Diamantina, MG, Tel: 0-000 Diamantina, MG, Brasil, Tel: 55 38 3532-1236; E-mail: ricbarata@hotmail.com

Received: September 23, 2017 Accepted: October 03, 2017 Published: October 10, 2017

marrow causing hepatosplenomegaly, malnutrition, jaundice and can lead the individual to death, if not treated in time [1]. In the Americas, about 90 % of human cases have been recorded in Brazil [2], where the disease is distributed in all regions and is present in 26 of the 27 federative units [3-5].

In the Americas, the protozoan *Leishmania infantum* is transmitted to humans primarily through blood feeding of female *Lutzomyia longipalpis*, a sand fly that has shown high ecological plasticity and adaptability to various environments [6-8], and therefore, it has been incriminated as one of those vectors responsible for the urbanization phenomenon of the disease in Brazil [9,10].

The use of indoor residual insecticides has been an action used by the Ministry of Health to control VL. Currently, synthetic pyrethroids such as cypermethrin and deltamethrin are the most commonly used insecticides in Brazil. However, low efficiency in vector control was observed and some studies have pointed to resistance of *Lu. longipalpis* for these compounds, which increases the interest in new chemical products [11-14].

Popularly known as pitch-black, mastic, wild musk, among others, *Protium heptaphyllum* is an evergreen tree species found in areas of forest and savannah ("cerrado"), highly very widespread in Brazil and South America, present from the Guianas to Argentina [15]. Studies show their sedative, anxiolytic, antidepressant, anti-inflammatory, antimicrobial and acaricidal activities [16-19]. Considering so many positive effects, this study aimed to investigate the insecticidal activity of the aqueous extract of *P. heptaphyllum* leaves on *Lutzomyia longipalpis*, a vector of visceral leishmaniasis (VL) in Brazil.

## Methods

### Collection, identification, and preparation of plant material


In June 2015, *Protium heptaphyllum* leaves were collected in rock field area in the city of Diamantina (Minas Gerais, Brazil) (18° 11' S - 43°34' O). The stuff was deposited in the DIAM / UFVJM herbarium in Diamantina under no. DIAM n° 5878 and its identification made based on morphological characters [15].

### Obtaining the plant extract

The collected material was subjected to drying in a forced circulation oven and air renewal to 60°C and milled in a knife mill to obtain a fine powder of uniform granulation. To prepare the aqueous extract was added to the glass beaker, the powder obtained after grinding and distilled water at room temperature, keeping at rest for 48h.

### Collection and identification of phlebotomine sand flies

The phlebotomine sand flies were collected in Diamantina / MG in the town of Aroeira (18° 88' S - 43° 38' W), using light traps HP [20] type exposed for two nights. The specimens were transported to the Laboratório de Parasitologia of Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM) and put in cages. From this material, a sample was taken for identification according to the classification proposed by Young & Duncan [21]. The sand flies remained at rest for 24h until perform the test, being offered to them sugar solution of distilled water and honey (1:1).



**Figure 1:** Proportion of *Lutzomyia longipalpis* (%) killed at four concentrations of the aqueous extract of *Protium heptaphyllum*: 0.25 mg/ml; 0.1 mg/ml; 0.05 mg/ml; 0.025 mg/ml; PC (positive control) = 196 µg/ml; NC (negative control) = 300 µl of distilled water; (n = 20). ▲ = values with non-significant difference ( $p > 0.05$ ) in relation to PC (equality test of proportions test - chi-square, with continuity correction or Fisher's exact test).

### Biological tests on *Lutzomyia longipalpis*

The aqueous extract of *P. heptaphyllum* was applied on the filter paper in the following concentrations: 0.025 mg/mL; 0.05 mg/mL; 0.1 mg/mL; 0.25 mg/mL and introduced inside translucent plastic pots. Controls included 20 adults that were treated with 100 µL distilled water (negative control) and 20 adults that were treated with 100 µL alpha-cypermethrin at 196 µg/mL concentration (positive control). Tests, including controls, were performed in duplicate, with 15 males and 15 females in each pot. After 1h, 2h, 4h, 8h, 12h, 24h, 48h and 72h the dead sand flies were counted. The overall temperature of the laboratory was 28°C.

### Statistical analysis

The statistical test used to compare proportions between groups at each time point was the proportion of equality test (chi-square), with continuity correction or Fisher's exact test. The software used for statistical analysis was the R (version 2015) [22], considering the 0.05 significance level.

### Results

The aqueous extract of *Protium heptaphyllum* demonstrated toxic properties in all concentrations tested. The mortality rate of *Lu. longipalpis* after 2h of exposure to the extract at a concentration of 0.1 mg/mL was 52%, behaving similarly to the positive controls (cypermethrin), and the extract at a concentration of 0.25mg / ml was able to kill 56% of insects within the first hour of exposure. The aqueous extract of *P. heptaphyllum* showed a mortality rate of about 81% and 83% after 72 h, at concentrations of 0.1 and 0.25 mg/mL, respectively (Figure 1).

### Discussion

The use of pyrethroids insecticides, mainly deltamethrin and alpha-cypermethrin, to eliminate phlebotomines constitutes a strategy of visceral leishmaniasis control. Deltamethrin has shown low efficacy on *Lu. longipalpis* [23-25]. Similarly, the alpha-cypermethrin had

low mortality rates in the wall, which led the authors to assume the possibility of resistance of *Lu. longipalpis* strain used or even the low efficacy of this insecticide [26].

Studies of plants with insecticidal action effects on *Lu. longipalpis* have been performed [14]. It showed the capacity of the neem seed oil (*Azadirachta indica*) on the development of *Lu. longipalpis*. In the same way, *Antonia ovata* and *Derris amazonica* displayed significant insecticide effect against *Lu. longipalpis* [27]. At the same time, studies have demonstrated the effects of plant extracts on other organisms. Tests with *P. heptaphyllum* on *Candida krusei*, *Cryptococcus neoformans*, *Staphylococcus aureus* and *Tetranychus urticae* demonstrate a fungicidal, bactericidal, acaricidal action, anti-inflammatory and healing activity [12,16-19,28,29-32]. In these works, the action effect found was associated with the presence of terpenes, components that are found in plants, used in defense against natural enemies [5]. Terpenes have been identified in large numbers in leaves of *P. heptaphyllum*, with β-caryophyllene being the most abundant component [33,34]. In view of the above, the presence of terpenes in extracts of *P. heptaphyllum* leaves, a molecule with recognized toxicity in insects, has been shown to be responsible for killing the phlebotomines and that the significant mortality rates observed in the present study show that *P. heptaphyllum* merits further study as to its potential use as an bioinsecticide.

### References

1. Brasil (2014) Manual de Vigilância e Controle da Leishmaniose Visceral (Ministério da Saúde). Ministério da Saúde, Brasília.
2. Grimaldi G, Tesh RB (1993) Leishmaniasis of the New World: current concepts and implications for future research. *Clin Microbiol Rev* 6: 230-250.
3. Gontijo CMF, Melo MN (2004) Leishmaniose visceral no Brasil: quadro atual, desafios e perspectivas. *Rev Bras Epidemiol* 7: 338-349.
4. Dantas-Torres F, Brandão-Filho SP (2006) Expansão geográfica da leishmaniose visceral no Estado de Pernambuco. *Rev Soc Bras Med Trop* 39: 352-356.
5. Cavalcante GM, Moreira AFC, Vasconcelos SD (2006) Potencialidade inseticida de extratos aquosos de essências florestais sobre mosca-branca. *Pesq Agropec Bras* 41: 9-14.

6. Jeronimo SMB, Oliveira RM, Mackay S, Costa RM, Sweet J, et al. (1994) An urban outbreak of visceral leishmaniasis in Natal, Brazil. *Trans R Soc Trop Med Hyg* 88: 386-388.
7. Ximenes MFFM, Silva VPME, Queiroz PVS, Rego MM, Cortez AM, et al. (2007) Flebotomíneos (Diptera: Psychodidae) e leishmanioses no Rio Grande do Norte, Nordeste do Brasil: reflexos do ambiente antrópico. *Neotrop Entomol* 36: 128-137.
8. Rangel EF, Vilela ML (2008) *Lutzomyia longipalpis* (Diptera, Psychodidae, Phlebotominae) and urbanization of visceral leishmaniasis in Brazil. *Cad Saúde Pública* 24: 2948-2952.
9. Costa CHN, Pereira HF, Araújo MV (1990) Epidemia de leishmaniose visceral no nordeste do Piauí, Brasil, 1980-1986. *Rev Saúde Pública* 24: 361-372.
10. Lainson R, Rangel EF (2005) *Lutzomyia longipalpis* and the eco-epidemiology of American visceral leishmaniasis, with particular reference to Brazil - a review. *Mem Inst Oswaldo Cruz* 100: 881-827.
11. Mazzarri MB, Feliciangeli MD, Maroli M, Hernandez A, Bravo A (1997) Susceptibility of *Lutzomyia longipalpis* (Diptera: Psychodidae) to selected insecticides in an endemic focus of visceral leishmaniasis in Venezuela. *J Am Mosq Control Assoc* 13: 335-341.
12. Viegas-Júnior C (2003) Terpenos com atividade inseticida: uma alternativa para o controle químico de insetos. *Quim. Nova* 26: 390-400.
13. Amóra SS, Bevilaqua CML, Feijó FMC, Alves ND, Maciel M (2009) Control of phlebotomine (Diptera: Psychodidae) leishmaniasis vectors. *Neotrop Entomol* 38: 303-310.
14. Maciel MV, Morais SM, Bevilaqua CML, Silva RA, Barros RS, et al. (2010) Atividade inseticida in vitro do óleo de sementes de nim sobre *Lutzomyia longipalpis* (Diptera: Psychodidae). *Rev Bras Parasitol Vet* 19: 7-11.
15. Domene VD, Mattos PP, Salis SM (2010) Fenologia e crescimento de almécega no Pantanal da Nhecolândia, Mato Grosso do Sul. *Comun. Técnico Embrapa Florestas* 263: 5.
16. Júnior GMV, Souza CML, Chaves MH (2005) Resina de *Protium heptaphyllum*: isolamento, caracterização estrutural e avaliação das propriedades térmicas. *Quim Nova* 28: 183-187.
17. Aragão GF, Carneiro LM V, Junior PF, Vieira LC, Bandeira PN, et al. (2006) A possible mechanism for anxiolytic and antidepressant effects of alpha- and beta- amyryn from *Protium heptaphyllum* (Aubl.) March. *Pharmacol Biochem Behav* 85: 827-834.
18. Holanda-Pinto SA, Pinto LMS, Cunha GMA, Chaves MH, Santos FA, et al. (2007) Anti-inflammatory effect of alpha, beta-amyryn, a pentacyclic triterpene from *Protium heptaphyllum* in rat model of acute periodontitis. *Inflammo Pharmacol* 15: 1-5.
19. Oliveira FA, Vieira-Junior GM, Chaves MH, Almeida FRC, Florencio MG, et al. (2004) Gastroprotective and anti-inflammatory effects of resin from *Protium heptaphyllum* in mice and rats. *Pharmacol Res* 49: 105-111.
20. Puggedo H, Barata RA, França-Silva JC, Silva JC, Dias ES (2005) HP : um modelo aprimorado de armadilha luminosa de sucção para a captura de pequenos insetos. *Rev Soc Bras Med Trop* 38: 70-72.
21. Young DG, Duncan MA (1994) Guide to the identification and geographic distribution of *Lutzomyia* sand flies in Mexico, the West Indies, central and south america (Diptera: Psychodidae). *Mem Am Entomol Inst* 54: 1-881.
22. Team RC (2015) R: A Language and Environment for Statistical Computing.
23. Seyedi-Rashi MA, Yezdan PH, Shah H, Jeradi M (1975) Susceptibility of *Phlebotomus papatasi* (Diptera: Psychodidae) to DDT in some foci of cutaneous leishmaniasis in Iran. *J Am Mosq Control Assoc* 8: 99-100.
24. Rahman SJ, Wattal BL, Mathur KK, Joshi GC, Kumar K (1982) Susceptibility of laboratory reared strain of *Phlebotomus papatasi* (Scopoli) to organochlorine insecticides. *J Commun Dis* 14: 122-124.
25. Alexander B, Usma MC, Cadena H, Quesada BL, Solarte Y, et al. (1995) Phlebotomine sandflies associated with a focus of cutaneous leishmaniasis in Valle del Cauca. *Colombia Med. Vet Entomol* 9: 273-278.
26. Nery-Guimarães F, Bustamante FM (1953) A aplicação domiciliar de DDT como base da profilaxia das leishmanioses. Estudo de um foco de leishmaniose muco-cutânea cinco anos depois da aspersão periódica com aquele inseticida. *Rev Bras Malariol Doenças Trop* 6: 127-130.
27. Luitgards-Moura JF, Bermudez EGC, Rocha AFI, Tsouris P, Rosa-Freitas MG (2002) Preliminary assays indicate that *Antonia ovata* (Loganiaceae) and *Derris amazonica* (Papilionaceae), ichthyotoxic plants used for fishing in Roraima, Brazil, have an insecticide effect on *Lutzomyia longipalpis* (Diptera: Psychodidae: Phlebotominae). *Mem Inst Oswaldo Cruz* 97: 737-742.
28. Vinaud MC, Souza R, Junior L (2008) Activity of *Stryphnodendron polyphyllum*, a plant from the Brazilian savannah, against hemocytes of *Biomphalaria glabrata*, an intermediate host of *Schistosoma mansoni*. *Rev Patol Trop* 37: 237-246.
29. Siani AC, Ramos MF, Menezes-de-Lima O, Ribeiro-dos-Santos R, Fernandez-Ferreira E, et al. (1999) Evaluation of anti-inflammatory-related activity of essential oils from the leaves and resin of species of *Protium*. *J Ethnopharmacol* 66: 57-69.
30. Susunaga GS, Siani AC, Pizzolatti MG, Yunes RA, DelleMonache F (2001) Triterpenes from the resin of *Protium heptaphyllum*. *Fitoterapia* 72: 709-711.
31. Pohlit AM, Quinard ELJ, Nunomura SM, Tadei WP, Hidalgo ADF, et al. (2004) Screening of plants found in the State of Amazonas, Brazil for activity against *Aedes aegypti* larvae. *Acta Amaz* 34: 97-105.
32. Lima-Júnior RC, Oliveira FA, Gurgel LA, Cavalcante IJ, Santos KA, et al. (2006) Attenuation of visceral nociception by  $\alpha$ - and  $\beta$ -amyryn, a triterpenoid mixture isolated from the resin of *Protium heptaphyllum*, in mice. *Planta Med* 72: 34-39.
33. Citó A, Costa F, Lopes J (2006) Identificação dos constituintes voláteis de frutos e folhas de *Protium heptaphyllum* Aubl (March). *Rev Bras PI Med* 8: 4-7.
34. Bandeira PN, Pessoa ODL, Trevisan MTS, Lemos TLG (2002) Metabólitos secundários de *Protium heptaphyllum* March. *Quim Nova* 25: 1078-1080.

## Author Affiliations

Top

<sup>1</sup>Laboratório de Parasitologia, Departamento de Ciências Biológicas, Universidade Federal dos Vales dos Jequitinhonha e Mucuri, Diamantina, MG, Brazil

<sup>2</sup>Laboratório de Farmacognosia, departamento de Farmácia, Universidade Federal dos Vales dos Jequitinhonha e Mucuri, Diamantina, MG, Brazil

<sup>3</sup>Laboratório de Matemática, Departamento de Matemática e Estatística, Universidade Federal dos Vales dos Jequitinhonha e Mucuri, Diamantina, Minas Gerais, Brazil

## Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • [www.scitechnol.com/submission](http://www.scitechnol.com/submission)