



Effect of Resistance Band Exercises on Neck Pain, Disability and Forward Head Posture in Dentists with Chronic Neck Pain

Pooja Pancholi*, Joginder Yadav and Sheetal Kalra

Abstract

Background: Dentists experience daily neck pain largely due to their awkward postures and high work demand. This puts extra strain on the muscles and predispose to development of faulty posture and muscle imbalance. So to avoid all these problems an exercise plan is needed, which can be easily performed even at the workplace. The purpose of this study was to determine the effects of resistance band exercises on neck pain, disability, and forward head posture in dentists with chronic neck pain.

Methodology: Fifty dentists between the ages of 25-50 participated in this study. They were randomly assigned to either an experimental (25) or a control group (25). The experimental group participated in resistance band exercises program while control performed conventional exercises for 10 weeks. Evaluation of neck pain and disability was done by Numeric pain rating scale (NPRS) and Neck Disability Index (NDI) respectively. Forward head posture was measured by digital photography technique.

Results: The paired t-test was used to compare within group differences and unpaired t-test was used to compare the between group differences. Significant differences found between experimental and control group in all three variables ($p < 0.05$).

Conclusion: The results of this study suggest that the use of resistance band exercises may be helpful for reducing neck pain, disability, and correcting forward head posture in dentists.

Keywords

Resistance band; Chronic neck pain; Forward head posture

Introduction

Musculoskeletal disorders have become increasingly common worldwide during the past decades. These problems are caused by repetitive, awkward, or stressful motions. Among the healthcare professionals, dentists are at high risk for developing profession-related disorders such as musculoskeletal injuries. Dentists experience daily neck pain largely due to their awkward postures and high work demand. common musculoskeletal disorders include: muscle strain, abhorrent neck and back postures, repetitive

stress injuries of the neck and shoulder girdles, and psychosocial stressors for back, neck, and shoulder complaints [1]. According to Fejer et al. lifetime prevalence of neck pain is 71% with an annual adult experience of between 12% and 34% [2]. In a study by Fejer et al. it was shown that 55% of the dentists experience neck pain. A career in dental profession is considered an occupational risk factor for neck pain, which can begin as early in their educational training [3]. Repeated unnatural, deviated, or inadequate working postures, forceful hand movements, inadequate equipment or workplace designs, and inappropriate work patterns are likely to be the particular risk factors for musculoskeletal disorders among dental professionals [4]. Occupational health problems in India estimates have shown that musculoskeletal disorders contributes to about 40% of all costs towards the treatment of work related injuries. Musculoskeletal problems have become significant issue for the profession of dentistry [5]. Neck disability index (NDI) has been found to be both a reliable and valid outcome tool [6]. Forward head and rounded shoulder posture in dentistry is very common that predisposes them to a muscle imbalance which develop between the neck and shoulder muscles. These muscles tend to fatigue quickly and weaken with prolonged forward head posture and rounded shoulder posture. Thus forward head posture is the most common cause of craniofacial pain, headache, neck ache, and shoulder pain and leads to decline in cervical joint motion. Digital photography is a form of photography that uses cameras containing arrays of electronic photodetectors to capture images focused by a lens. Many researchers have used digital photography technique as a method of assessment for forward head posture [7].

There are many treatments options available for alleviating neck pain such as: electrotherapy including TENS, Hotpacks, laser, manipulation, ultrasound and acupuncture therapy. According to Ferreira et al. multimodal physiotherapy programme (including exercises, education, and ergonomics) was found as a better treatment option for reduction of chronic neck pain [8]. Strengthening exercises may decrease pain and increase neck range of motion and muscle performance. Isometric exercise is commonly used to increase muscle performance. Posture correction is recommended to dentists with poor neck postures. This will lead to physical and psychological comfort to dentist during execution of clinical act. In dentists protracted posture causes overstress in the lower part of cervical spine so retraction exercises and nodding exercises are recommended as posture correction exercises [9,10]. Body weight, resistive bands, pulleys and weight machines are a few modes of dynamic resistive exercises. Strength training is beneficial in decreasing neck pain and also specific strength training with theraband has been considered preventive regarding neck pain among military pilots and office workers [11,12]. A Theraband is an elastic band used for strength training. Resistance band training is now used widely as part of general fitness and strength training. Typically the bands are color coded to show different levels of resistance and users need to select an appropriate level and are simple to use and their light weight allows people to easily carry them if traveling and continue with routine sessions for strength training.

*Corresponding author: Pooja Pancholi, Department of Physiotherapy, Shree Guru Gobind Singh Tricentenary University, Gurugram, India, Tel: 8059331625; E-mail: poojasharma6101994@gmail.com

Received: April 02, 2018 Accepted: April 27, 2018 Published: May 04, 2018

Material and Methods

Purpose of the study

The purpose of this study was to determine the effects of resistance band exercises on neck pain, disability, and forward head posture in dentists with chronic neck pain.

Neck pain is the major problem in dentists so to avoid neck related problems like pain, disability and faulty posture they need to stick to the exercises. There is dearth of the studies which have evaluated the efficacy of theraband based exercises on neck pain, disability and forward head posture (FHP) in dentists with chronic neck pain.

Study design

Comparative study design

Study centre

The study was conducted at Physiotherapy OPD of SGT Hospital, Budhera, Gurgram

Sample

A total of 50 subjects were randomly selected and assigned into two groups. Each group having 25 subjects named as Group A and Group B.

Group A-Conventional exercises (Control group- Stretching exercises, strengthening and posture correction exercises were given)

Group B-Theraband exercises+Conventional exercises (Experimental group)

Sampling

Convenience random sampling

Inclusion criteria

Age group

25-50 years

Chronic mechanical neck pain, Symptoms more than three months in duration.

NDI Score 14\50, NPRS Score 5/10

Males and females both

Forward head posture (CV angle <50 degrees)

Exclusion criteria

Previous history of whiplash injury, Previous history of injury to cervical spine, Diagnosed with osteoporosis, Malignancy or other spinal infection, Patients on medication for neck pain, Diagnosed with cervical radiculopathy

Scales

Numeric pain rating scale (NPRS)

Neck pain disability index (NPDI)

Digital photography technique

Procedure

The subjects who fulfilled the inclusion criteria were included in the study. The whole procedure was explained to them and the

informed consent was taken from them. 50 subjects were randomly selected using convenience random sampling and assigned into two groups named as group A and group B.

Intervention

Group A: Control Group

Hot fomentation was given by applying cervical hot pack over the area in prone lying position for 15 min 45 along with conventional exercises. These include Isometrics exercises for neck including flexors, extensors, lateral flexors and Postural correction exercises included- Nodding, shoulder shrugs and shoulder bracing exercises. Stretching of trapezius muscle was also given to control group. Hot fomentation and postural correction exercises was given to both the groups.

Group B: Experimental group

Resistance band exercises- In addition to conventional exercises, resisted exercises using elastic band were given. The training program included four training exercises for the prime movers of the neck during cervical flexion, extension and lateral flexion. Exercises was performed with a head harness using different color-coded elastic resistance bands). Cervical flexion was performed seated (Figure 1).

Both groups performed exercises 5 times a week for 10 weeks. Resistance band exercise groups gradually increased resistance to next level when they were able to performed 5 to 6 sets of 8 to 12 repetitions. Exercise was performed with the color band that was prescribed for the patient, or with a color that allows him or her to complete 2 to 3 sets of 10 to 15 repetitions with mild fatigue on the last set. Then Progression was done to the next color band when patient was able to complete the 3 sets of 10 to 15 repetitions [13].

Data was be collected on baseline, 5th week and last day of 10th week. Data was analyzed using SPSS version 21.

Results

The Mean and standard deviations for age ,height, weight, and body mass index (BMI) are shown for both groups in Table 1.




Figure 1: Demonstration of resistance band exercises for neck side flexion.

Comparison of change in NPRS

The between group analysis of the NPRS using unpaired t-test showed that there was no significant difference seen between both groups at baseline and 5th week ($p > 0.005$), but a highly significant difference was seen on 10th week ($p < 0.01$) (Table 2).

Comparison of change in NDI between both groups

The between group analysis of the NDI using unpaired t-test showed that there was no significant difference seen at baseline and 5th week ($p > 0.05$), but a highly significant difference was seen on 10th week ($p < 0.01$) (Table 3).

Between group comparison of CV angle

The between group analysis of the CV angle using unpaired t-test

showed that a significant difference was seen at 5th week ($p < 0.05$), but a highly significant difference was seen on 10th week ($p < 0.01$) (Table 4).

Discussion

A work-related musculoskeletal disorder (WRMD) is defined as a musculoskeletal injury that results from a work-related event. This may result from prolonged work time. These types of injuries are common among dentists [14]. Musculoskeletal problems have become a significant issue for the profession of dentistry and dental hygiene. Reported prevalence of work-related musculoskeletal disorders (WRMD) is about 63 to 93 percent among dental professionals and the prevalence of general musculoskeletal pain ranges between 64% and 93%. The most prevalent regions for pain in dentists have been shown to be the back (36.3-60.1%) and neck

Table 1: Mean comparison of Age, Height, Weight and BMI.

	Group	Mean ± Standard Deviation	t-Value	p-Value
Age	Group A	37.8 ± 7.12	1.862	0.15 ^{NS}
	Group B	40.72 ± 7.17356		
Height	Group A	1.66 ± 0.03187	0.552	0.58 ^{NS}
	Group B	1.6608 ± 0.03187		
Weight	Group A	58.56 ± 4.112	0.472	0.68 ^{NS}
	Group B	58.12 ± 4.12634		
BMI	Group A	21.2 ± 1.681	0.095	0.98 ^{NS}
	Group B	21.088 ± 1.62923		

* $p < .05$ (Significant) NS (Not significant)

Table 2: Between group comparison of NPRS.

	Group	Mean ± Standard deviation	t-value	p-value
Baseline NPRS	Control	7.92 ± 1.038	0.140	0.889 ^{NS}
	Experimental	7.96 ± .978		
5th week NPRS	Control	5.80 ± 1.041	0.141	0.889 ^{NS}
	Experimental	5.76 ± .970		
10th week NPRS	Control	4.44 ± .651	12.97	0.000 ^{**}
	Experimental	1.52 ± .918		

** $p < 0.01$ HS (Highly significant)

NS-not significant

Table 3: Between group comparison of NDI.

	Group	Mean ± Standard deviation	t-value	p-value
Baseline NDI	Control	38.76 ± 2.047	0.944	0.350 ^{NS}
	Experimental	39.36 ± 2.430		
5th week NDI	Control	32.00 ± 3.253	0.778	0.440 ^{NS}
	Experimental	31.36 ± 2.515		
10th week NDI	Control	23.84 ± 3.325	7.674	0.000 ^{**}
	Experimental	18.16 ± 1.625		

** $p < 0.01$ HS (Highly significant)

NS-not significant

Table 4: Between group comparison of CV angle.

	Group	Mean ± Standard deviation	t-value	p-value
Baseline Cv angle	Control	25.48 ± 3.864	1.002	0.321 ^{NS}
	Experimental	26.56 ± 3.754		
5th week Cv angle	Control	30.08 ± 2.753	4.549	0.02 [*]
	Experimental	34.08 ± 3.761		
10th week CV angle	Control	34.44 ± 2.256	10.809	0.000 ^{**}
	Experimental	44.92 ± 4.291		

** $p < 0.01$ HS (Highly significant)

NS-not significant

(19.8-85%). This study compared two types of exercises protocols (conventional versus resistance band) in reducing pain, disability and forward head posture in dentists with chronic neck pain. The subjects in this study had similar baseline values of all dependent variables suggesting that all groups had homogenous distribution of patients. The results of this study revealed that although both the groups improved significantly at the end of the 10th week but addition of "resistance" band exercises (group B) to the conventional exercises (Group A) yielded improved NPRS and NDI outcomes, and an increase in forward head angle measured by craniocervical angle compared to control group (Group A).

In summary, the results of our study lead us to reject null hypothesis thereby confirming that both treatment exercise groups (conventional exercises and resistance band exercises) were effective in reducing neck pain (1.92%), disability (29%), and forward head posture in dentists with chronic neck pain. Improvement in both groups was found to be Statistically significant. But it was a strong finding of the intervention that there was more reduction in pain, disability and forward head posture, which was found to be (1.92%), (29%) and (23%) improved in experimental group after the intervention as compared to group A. The results of our study are in accordance with the result of the study done by Jull et al. In this study they compared the effect of neck muscles strength training on neck pain and disability. They found that 4 weeks of strength training resulted in reducing pain, disability in subjects suffering from chronic neck pain [15]. Resistance training with resistance band for shoulder muscles have shown increase in muscle power in novice lifters. Other studies have also demonstrated an increase in power generation in the lower body following EB (elastic band) training. This study showed that there are positive effects of resistance training on muscular performance of novice lifters [16]. Our results are further supported by the study done by Seguin R et al. they investigated the effects of strength training on pain sensitivity. In this experiment, 62 women (40 with shoulder pain, 20 without) participated in either a general exercise program or specific strength training for their shoulders and results of the study revealed that Pain tolerance increased in response to strength training in the women who started out with pain [17]. Strength training with elastic bands have been proven effective in reducing pain according to the results of a study done by Mats Hagberg et al. [18]. They studied the effect of strength training on function of chronically painful muscles. Results of the study revealed that the strength training as therapy for shoulder pain had positive results in women with shoulder pain, researchers found that "specific strength training with the aid of resistance band relieved pain and increases maximal activity and their pain was reduced by 42-49% [18]. Ylinen et al. studied the effect of specific training with elastic bands to Reduce Chronic Neck Pain and disability. They divided 180 female office workers with chronic neck pain into three groups: one group did strength training with elastic bands (lateral shoulder raise) another did endurance training, and a third did nothing. They found that "both strength and endurance training decreased perceived neck pain and disability in office workers [19].

Magni et al. studied the effects of resistance training on muscle strength, joint pain, and hand function in individuals with hand osteoarthritis. They found that resistance training has a slight significant effect on grip strength or hand function [20]. In the study performed by Magistro D et al. In their study they found the effects of resistance band training on women having sedentary life style. According to this study, healthy adaptations can occur using

resistance band exercise. The results indicate that a resistance exercise can provide healthy muscle adaptations [21]. Gupta et al. in his study concluded that in people with neck pain, addition of conventional strength and high-load endurance retraining of muscles leads to reduction in neck pain and associated symptoms [22]. In a study performed by Lidegaard et al. involving office workers, it was found that only a little exercises say for 2 min daily for 10 weeks with resistance band can effectively reduce neck and shoulder pain. In this study subjects were asked to perform resistance band based exercises for 2 minutes daily for a period of 10 weeks, and it has led to a significant decrease in neck and shoulder pain [23]. Our results are also supported by Kim et al. they investigated the effects of elastic band exercises program on the posture of subjects with rounded shoulder and forward head posture. The results of the study suggested that the strength training with the aid of elastic band was helpful in correcting rounded shoulder and forward head posture and it showed a significant result [24]. To conclude both groups were found to be effective in reducing pain, disability and forward head posture. Results of the study revealed that addition of resistance band exercises as a mode of resistance training program when added to conventionally given exercises which includes (stretching, isometrics and posture correction exercises) brings better improvements in terms of reducing pain, disability and improving forward head posture.

Limitations of the Study

1. Sample size was small.
2. Follow up of the subjects was not done to see if the effects are retained.
3. Specification of the work is not mentioned.

Relevance to Clinical Practice

This study showed a significant overall improvement in neck pain, disability and reduction in forward head posture in both control and experimental groups but the results of resistance band training is more significant than in conventional exercises group. Hence, the result of this study provide the evidence that the Resistance band training may be the valuable and useful tool in clinical practice and is consistence with the current use by clinical physiotherapist in reducing pain ,disability and extent of forward head posture.

Future Research

1. Study can be done on the wider sample.
2. Study can be done on different subjects and different age groups and other population like bankers, IT workers etc.
3. Study can be done on the basis of specifications of the work like in oral surgeons, endodontic etc.
4. Further research can be done on the long term effects of the resistance band training on the subjects with chronic neck pain.

Conclusion

This study compared the effectiveness of conventionally provided exercises to resistance band exercises in reducing pain, disability and forward head posture in dentists with chronic neck pain. The results show that resistance band exercises are more effective. They are beneficial for muscle strengthening and possess the added advantages of their ease of use, variability of tension, and portability. Our research hypothesis has been verified and our null hypothesis rejected.

1. Neck pain decreased in both the groups ,Group B improved much better than group A.

2. Neck disability was reduced in both the groups but group B showed better results than group A.

3. There was a reduction in extent of forward head posture in both the groups, but group B shows more better results.

Overall we conclude that Resistance band exercises are more effective than conventionally given exercises in reducing pain, disability and forward head posture. Thus research hypothesis has been verified that resistance band exercises are more effective in reducing neck pain, disability and forward head posture as compared to conventional exercises and null hypothesis is rejected that there is no significant difference between resistance band exercises and conventional exercises. Resistance bands has many advantages as exercises using resistance band are very helpful in building strength and it comes with variety of resistance in different color codes according to the need of the patient. resistance bands are light weight and easily portable so it can be carried while travelling and it is cost effective too. Resistance bands can replace the weights and it is easy to use alone as it can be used at home and work place. Resistance bands are ideal for exercising any time.

References

- Hayes MJ, Smith DR, Cockrell DJ (2010) An international review of musculoskeletal disorders in the dental hygiene profession. *Int Dent J* 60: 343-352.
- Fejer R, Kyvik KO, Hartvigsen J (2006) The prevalence of neck pain in the world population: a systematic critical review of the literature. *Eur Spine* 15: 834-848.
- Fejer R (2006) Neck pain- Prevalence, genetic and environmental factors. *Ugeskr. Laeg* 168.
- Finsen L, Christensen H, Bakke M (1998) Musculoskeletal disorders among dentists and variation in dental work. *Appl Ergon* 29: 119-125.
- Leggat PA, Kedjarune U, Smith DR (2007) Occupational health problems in modern dentistry. *Ind Health* 45: 611-621.
- Pietrobon R, Coeytaux RR, Carey TS, Richardson WJ, DeVellis RF (2002) Standard scales for measurement of functional outcome for cervical pain or dysfunction. *Spine* 27: 515-522.
- Croft PR, Lewis M, Papageorgiou AC, Thomas E, Jayson MIV, et al. (2001) Risk factors for neck pain. a longitudinal study in the general population. *Pain* 93: 317-32.
- Ferreira GE, Barreto RG, Robinson CC, Plentz R,DM, Silva MF (2016) Global Postural Reeducation for patients with musculoskeletal conditions: a systematic review of randomized controlled trials. *Braz J Phys Ther* ;20: 194-220.
- Warren N (2010) Causes of musculoskeletal disorders in dental hygienists and dental hygiene students A study of combined biomechanical and psychosocial risk factors *Work* 35: 441-454.
- León-Hernández JV, Martín-Pintado-Zugasti A, Frutos LG, Alguacil-Diego IM, Llave-Rincón AL, et al. (2016) Immediate and short-term effects of the combination of dry needling and percutaneous TENS on post-needling soreness in patients with chronic myofascial neck pain. *Braz J Phys Ther* 20.
- Rush PJ, Shore A (1994) Physicians perceptions of the Value of Physical Modalities in the Treatment of Musculoskeletal Disease. *Br.J.Rheumatol* 33: 566-568.
- van den Oord MHAH , De Loose V, Meeuwse n T , Sluiter JK, Frings-Dresen MHW (2010) Neck pain in military pilots: prevalence and associated factors. *Mil Med* 175: 55-60.
- Murray M, Lange B, Nørnberg BR, Sogaard K, sjogaard G (2015) Specific exercise training for reducing neck and shoulder pain among military helicopter pilots and crew members. a randomized controlled trial protocol. *BMC Musculoskelet Disord* 16: 198.
- Rambab T, Suneetha K (2014) Prevalence of Work Related Musculoskeletal Disorders Among Physicians, Surgeons and Dentists: A Comparative Study. *Ann Med Health Sci Res* 4: 578-582.
- Jull GA, Falla D, Vicenzino B, Hodgesa PW (2009) The effect of therapeutic exercise on activation of the deep cervical flexor muscles in people with chronic neck pain. *Man Ther* 14: 696-701.
- Markus DJ, Emil S, Christoffer HA, Mette KZ, Peter M, et al. (2012) Evaluation of Muscle Activity During a Standardized Shoulder Resistance Training Bout in Novice Individuals. *J Strength Cond Res* 26: 2515-2522.
- Seguin R, Nelson ME (2003) American journal of preventive medicine. The benefits of strength training for older adults. *Am J Prev Med* 25: 141-149.
- Hagberg M, Harms-Ringdahl K, Nisell R, Hjelm EW (2002) Rehabilitation of neck-shoulder pain in women industrial workers: A randomized trial comparing isometric shoulder endurance training with isometric shoulder strength training. *Arch Phys Med Rehabil* 81: 1051-1058.
- Yiinen JJ, Hakkinen AH, Takala EP, Nykanen MJ, Kautianen HJ, et al. (2006) Effects of neck muscle training in women with chronic neck pain: one year follow up study. *J Strength Cond. Res* 20: 6-13.
- Magni NE, McNair PJ, Rice DA (2017) The effects of resistance training on muscle strength, joint pain, and hand function in individuals with hand osteoarthritis: a systematic review and meta-analysis. *Arthritis Res Ther* 19: 131.
- Magistro D, Liubicich ME, Candela F, Ciairano S (2014) Effect of Ecological Walking Training in Sedentary Elderly People: Act on Aging Study. *The Gerontologist* 54: 611-623.
- Gupta BD, Aggarwal S, Gupta B, Gupta M, Gupta N (2013) Effect of Deep Cervical Flexor Training vs. Conventional Isometric Training on Forward Head Posture, Pain, Neck Disability Index In Dentists Suffering from Chronic Neck Pain. *J Clin Diagn Res* 7: 2261-2264.
- Lidegaard M, Jensen RB, Andersen CH, Zebis MK, Colado JC, et al. (2013) Effect of Brief Daily Resistance Training on Occupational Neck/Shoulder Muscle Activity in Office Workers with Chronic Pain: Randomized Controlled Trial. *BioMed Res Int* 2013: 1-11.
- Kim TW, An DA, Lee HY, Jeong HY, Sung YH (2016) Effects of elastic band exercise on subjects with rounded shoulder posture and forward head posture. *J Phys Ther Sci* 28: 1733-1737.

Author Affiliations

Top

Department of Physiotherapy, Shree Guru Gobind Singh Tricentenary University, Gurugram

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission