



Short Communication

Evaluation of Diagnostic Value of Amniotic Fluid Ferritin and Its Relation to the Pregnancy Complications

Khadije Rezaie Keikhaie¹, Kolsoum rezaie Keikhaie², Leili rezaie Keikhaie^{2*}, Ashrafali rezaie Keikhaie³ and Morteza salarzaei⁴

Abstract

Introduction: The health of the child is in close relation with the health of the mother and his access to health care. Among these services, it is possible to perform prenatal care with the aim of identifying risk factors and predicting and preventing pregnancy complications.

Methodology: This descriptive-analytic study was performed on all pregnant women referring to the clinic of Amiral-Momenin Hospital in Zabol in 2017; pregnant mothers with gestational age more than 20 weeks.

Results: The present study was conducted on 100 pregnant women referring to the clinic of Amiral-Momenin Hospital in Zabol in 2017; the mean age of studied subjects turned out to be 29.86 ± 6.22 years.

Discussion and conclusion: The results of this study showed that the mean ferritin level in women with preeclampsia and premature rupture of membrane was significantly higher than healthy women without complications of pregnancy; however, ferritin in the other complications of pregnancy, including Gestational diabetes, fetal abnormalities, placental abruption, and oligohydramnios were not statistically different with healthy pregnant women.

Keywords

Ferritin; Pregnancy; Zabol; Placental abruption; Abnormalities

Introduction

The health of the child is in close relation with the health of the mother and his access to health care. Among these services, it is possible to perform prenatal care with the aim of identifying risk factors and predicting and preventing pregnancy complications [1]. Here's a brief overview of some of the common complications of pregnancy.

Methodology

This descriptive-analytic study was performed on all pregnant women referring to the clinic of Amiral-Momenin Hospital in

Table 1: Comparison of mean ferritin level in pregnant women with preeclampsia and eclampsia and healthy subjects.

Group	ferritin	P value
Positive preeclampsia	92.25 ± 14.81	0>0001
Negative preeclampsia	37.48 ± 13.29	
Positive diabetes	8.38 ± 33.28	0.148
Negative diabetes	37.48 ± 13.29	
Positive oligohydramnios	10.29 ± 46	0.219
Negative oligohydramnios	37.480. ± 13.29	
Positive placental abruption	9.19 ± 35.5	0.605
Negative placental abruption	37.48 ± 13.29	
Positive fetal abnormalities	6.42 ± 31	0.148
Negative fetal abnormalities	37.48 ± 13.29	
Positive premature rupture of membrane	15.32 ± 88.90	0.148
Negative premature rupture of membrane	37.48 ± 13.29	

Zabol in 2017; pregnant mothers with gestational age more than 20 weeks, single-strike pregnancy and satisfaction to enter the study were the main inclusion and proven systemic infections, diabetes, hypertension, pre-pregnancy, internal disease, mental illness, smoking, non-Iranian nationality, Corticosteroids consumption and immune stimulants in the last 4 weeks and the use of Non-steroidal anti-inflammatory medicines were the main exclusion criteria. The collected data was analyzed using SPSS18 and descriptive statistics in form (frequency tables and charts) (Table 1).

Findings

The present study was conducted on 100 pregnant women referring to the clinic of Amiral-Momenin Hospital in Zabol in 2017; the mean age of studied subjects turned out to be 29.86 ± 6.22 years.

Discussion and Conclusion

The health of the child is in close relation with the health of the mother and her access to health care. Among these services, it is possible to perform prenatal care with the aim of identifying risk factors and predicting and preventing complications of pregnancy [2]. The aim of this study was to evaluate the diagnostic value of amniotic fluid lactate dehydrogenase and its relationship with pregnancy complications in pregnant women with gestational age greater than 20 weeks referred to the clinic of Amiral-Mu-menin Hospital in Zabol in 1395. The mean age of studied subjects turned out to be 29.86 ± 6.22 years. The mean ferritin level in women with preeclampsia was 92.25 ± 14.81, which turned out to be significantly higher than healthy pregnant women. The mean ferritin level in women with gestational diabetes mellitus was 33.28 ± 8.38, but the mean of pregnant women with diabetes mellitus was not significantly different from healthy subjects. The mean ferritin level in women with oligohydramnios was 46 ± 10.29, but. The mean ferritin level in women with placental abruption was 35.5 ± 9.19, but it was not significantly different from healthy subjects. The mean ferritin level in women with fetal abnormality was 31 ± 6.42; it was not significantly different from healthy subjects. The mean ferritin level in women with premature rupture of the membrane was 88.90 ± 15.32, which turned out to be significantly different from the control group. In general, the results of this study showed that the mean ferritin level in

*Corresponding author: Leili rezaie Keikhaie, Zabol University of Medical Sciences, Zabol, Iran, E-mail: tahghighatt2@gmail.com

Received: January 27, 2018 Accepted: March 08, 2018 Published: March 20, 2018

women with preeclampsia and premature rupture of membrane was significantly higher than healthy women without complications of pregnancy; however, ferritin in the other complications of pregnancy, including Gestational diabetes, fetal abnormalities, placental abruption, and oligohydramnios were not statistically different with healthy pregnant women [3-6].

References

1. Behzadmehr R, Keikhaie KR, Pour NS (2017) The Study of Pregnant Women's Attitude toward Using Ultrasound in Pregnancy and its Diagnostic Value based on the Demographic Features in Amir-al-Momenin Hospital of Zabol. *Int J Adv Res Biol Sci* 4: 58-63.
2. Kahkhaie KR, Keikhaie KR, Vahed AS, Shirazi M, Amjadi N (2014) Randomized comparison of nylon versus absorbing polyglactin 910 for fascial closure in caesarean section. *Iranian Red Crescent Med J* 16.
3. Rezaie Keikhaie K, Ghorbani S, Hosseinzadeh Z, Hassanshahian M (2017) Antimicrobial activity of methanol extract of *Citrullus colocynthis* against antibiotic-resistant *Staphylococcus aureus*. *Advan Herb Med* 3: 1-6.
4. Shirazi M, Hantoush-Zadeh S, Rezaie-Keikhaie K, Pirjani R (2016) Spontaneous Uterine Rupture and Live Fetus in 21th Week of Pregnancy with Hemorrhagic Shock Due to Placenta Percreta: A Case Report. *Case Reports in Clinical Pract* 1: 19-21.
5. Behzadmehr R, Keikhaie KR, Pour NS. *INTERNATIONAL JOURNAL OF PHARMACY & LIFE SCIENCES*.
6. Keikhaie KR, Behzadmehr R, Salarzaeis M (2017) Comparison of Two Doses of Oxytocin Regimes to Prevent Post-Caesarean Bleeding Due to Uterine Atony in Pregnant Women Referring to Amir-al Momenin Hospital of Zabol in 2016. *Ann Med Health Sci Res*.

Author Affiliations

Top

¹Department Obstetrics and Gynecology, Maternal and Fetal Health Research Center, Zabol University of Medical Sciences, Zabol, Iran

²Zabol University of Medical Sciences, Zabol, Iran

³Zabol University, Zabol, Iran

⁴Student Research Committee, Zabol University of Medical Sciences, Zabol, Iran

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission