



# Insect and Mammalian Pests of Large Cardamom (*Amomum subulatum* Roxburgh) in Sikkim Himalaya

Deka TN, Gudade BA\*, Saju KA and Bora SS

### Abstract

Large cardamom is an important cash crop of Sikkim and Darjeeling district of West Bengal. About 3863 MT of large cardamom are produced annually from an area of 26,459 ha in Sikkim region. Insect pests are one of the constraints of large cardamom production. Surveillance was carried out in the major large cardamom growing tracts during 2010 - 12 and incidences of major pests were recorded. Minor insect pests, mammalian pests and new pests in large cardamom ecosystem were documented. Entomopathogens of leaf caterpillar (*Artona chorista*) were identified.

### Keywords

Pests; Large cardamom; *Artona*; Shoot fly; Sikkim Himalaya

### Introduction

Large cardamom (*Amomum subulatum* Roxburgh) belonging to family *Zingiberaceae* is an important spices crop of North East India and often referred as currency crop [1]. Now-a-days neighboring countries of India like Nepal, Bhutan and Myanmar have also started cultivation of large cardamom [2]. In Sikkim Himalayas it is cultivated since time immemorial and believed to be the native of the state [3]. Sikkim has the largest area as well as production of large cardamom in the globe and act as a big player in the world market [4]. It grows well at altitude ranges from 1000 to 2200 m MSL with well distributed rainfall spread around 200 days with a total of about 3000-3500 mm/year [5-7]. Large cardamom is a sciophyte i.e. the plant is grown under shade [8,9]. Crop prefers humid subtropical, semi evergreen forests hills of eastern sub-Himalayan region [10]. It is essentially a cross-pollinated crop due to the heterostyly nature of its flowers. Bumble bee (*Bombus breviceps*) is the major pollinator in large cardamom due to its high pollination efficiency attributed to its big body size and foraging habit [11,12]. In Sikkim the average productivity of large cardamom ranges from 240 to 260 kg/ha [13]. The crop grows somewhat wild in nature and damages due to insect pests are common. It is infested by various pests and diseases causing considerable amount of monetary loss [14-16]. About 3,863 MT of large cardamom are being produced annually from an area around 26,459 ha in Sikkim region [17]. Here we have presented

our observations recorded on insect pests of the crop and their IPM management schedule in Sikkim Himalaya.

### Materials and Methods

Surveillance was carried out in major large cardamom growing tracts of Sikkim and Darjeeling district of West Bengal during 2010-12. Data were recorded based on two different types of surveillance viz., fixed plot and roving surveillance. Fixed plot surveillance was conducted at Kabi (1567 m amsl), North Sikkim and Pangthang (1952 m amsl), East Sikkim at monthly interval.

Roving surveillance was carried out in four districts of Sikkim, Darjeeling and Kalimpong sub division in Darjeeling district of West Bengal. Data recorded in two gardens from each district / sub divisions and per cent incidence were recorded. Roving surveillance was carried out twice in a year in the months of April - May and October - December.

In both the surveillance, twenty five clumps of large cardamom per garden were selected through random sampling and pests of major and minor importance recorded. Numbers of infested tillers per clumps were noted and per cent incidence for important pests was derived. Natural enemies (entomopathogens) associated with *Artona chorista* in field condition were recorded and pathogenicity test was carried out.

### Results and Discussion

In the fixed plot surveillance, occurrence (per cent) of three important insect pests viz., stem borer (*Glyphipterix* sp.), shoot fly (*Merchlorops dimorphus*) and leaf caterpillar (*Artona chorista*) was recorded. *M. dimorphus* was recorded highest among the different pests recorded in the surveyed areas (Table 1). It was 57.00 per cent in Pangthang and 46.30 per cent in Kabi. Roving surveillance was carried out in four districts of Sikkim, Darjeeling and Kalimpong sub division in Darjeeling district of West Bengal. Data on pest incidence and traditional practice of pest management was recorded. Occurrence of shoot fly (*M. dimorphus*) was recorded the highest (44.00 per cent) among the other major pests recorded during roving surveillance followed by leaf caterpillar (*A. chorista*) and stem borer (*Glyphipterix* sp.). From the data it is revealed that occurrence of *M. dimorphus* and *Glyphipterix* sp. was highest in East Sikkim than other districts / sub division of West Bengal. Infestation of *A. chorista* was observed more in Darjeeling district of West Bengal than other surveyed areas (Table 2). In both the surveillance incidence of

Table 1: Incidence of insect pests at fixed plots in Sikkim region during 2010 -12.

Pests	Incidence percent ( Range)	
	Kabi	Pangthang
Stem borer ( <i>Glyphipterix</i> sp.)	0-16	0-20
Shoot fly ( <i>M. dimorphus</i> )	16-64	16-72
Leaf caterpillar ( <i>A. chorista</i> )	0-24	0-16

\*Corresponding author: BA Gudade, Indian Cardamom Research Institute, Regional Research Station, Spices Board, Tadong, Gangtok-737 102, Sikkim, India, Tel +91-89007 22294; E-mail: bgudade@gmail.com

Received: July 27, 2016 Accepted: August 10, 2016 Published: August 17, 2016

**Table 2:** Incidence of insect pest in large cardamom field in different districts of Sikkim and Darjeeling district of West Bengal during 2010-12.

Name of the districts/ Sub Division	Incidence (percent) – Range		
	Insect pests		
	Stem borer ( <i>Glyphipterix</i> sp.)	Shoot fly ( <i>M. dimorphus</i> )	Leaf caterpillar ( <i>A. chorista</i> )
North	12-20	32-40	8-12
East	16-24	40-48	12-16
South	4-12	36-40	0-12
West	8-20	32-40	8-16
Kalimpong	4-8	12-28	0 - 4
Darjeeling	4-16	28-44	16-28

**Table 3:** Wild mammalian pests of large cardamom in Sikkim region during 2010-12.

Name of the pests	Vernacular name	Period of occurrence	Name of the infested parts	Places
Rodents	Musha	Aug-Dec	Spikes / capsules	Kabi, Pangthang and Dzongu
Himalayan palm civet	Kala	Aug –Dec	capsules	Kabi and Pangthang
Indian wild boar	Bongdel	April- Sept	Clumps	Kabi
Musk deer	Kasturi Mirga	Aug-Dec	Spikes and capsules	Dzongu and Kabi
Monkey	Bandar	April-Sept	Clumps	Kewzing and Namchi

**Table 4:** New pest recorded in the cardamom ecosystem in Sikkim during 2010-12.

SI. No.	Pest	Host	Remarks
1	Aphid ( <i>Mollitrichosiphum</i> sp.)	Utis ( <i>Alnus nepalensis</i> )	Utis is an important shade tree of large cardamom ecosystem

**Table 5:** Traditional practices of pest management in Sikkim.

SI./No.	Traditional practice	Against pest
1.	Collection and burning of infested plant parts	Leaf caterpillar, stem borer and shoot fly.
2.	Leave extract – Titepati, Chillowney, Indreni and Tobacco	Chewing and sucking Pests
3.	Field exposures to Sun shine & Bird	Soil insects mainly white grub
4.	Application of ash and ash solution	Ants and aphids
5.	Covering of Cardamom clumps with dried leaves / small branches	Mammalian Pests viz., civet, wild boar and deer
6.	Crop rotation	White grub and other soil borne insects
7.	Drainage / Flooding	Soil insects – white grubs
8.	Kerosene	House hold pests / Ants
9.	Mint leaves	House hold pests / House fly


mammalian pests and also a new pest in cardamom ecosystem was recorded. Mammalian pests became a cause of concern in some areas. Presence of five mammalian pests was recorded based on damage caused by them and identified with the help of local farmers (Table 3). Utis (*Alnus nepalensis*) is an important shade tree for large cardamom [18]. An aphid species is commonly observed in the twigs of the tree. The identification of the pest was confirmed from Network Project on Insect Biosystematics-NPIB), Indian Agricultural Research Institute, New Delhi as *Mollitrichosiphum* sp. (Registered in IARI, New Delhi as RRS NO. 2133-2162/11) (Table 4 and Plate 1). Occurrence of some other minor insect pests viz., lace wing bug, fruit borer, aphid, mealy bug, hairy caterpillar, scale insects, grass hopper, rhizome weevil and thrips were also recorded during the surveillance. Attempt was also made to find out the association of natural enemies with these pests. Dead caterpillars of *A. chorista* was observed in the field conditions and collected for isolating the associated pathogens. Four entomopathogens namely *Cladosporium* sp., *Metarrhizium* sp., *Penicillium* sp. and *Verticillium* sp. were isolated and identified. In the pathogenicity test *Cladosporium* sp. was found promising under laboratory conditions. Large cardamom is basically grown organically in the Sikkim Himalayan region. Nine different practices are adopted by the farmers to manage the agricultural and house hold pests in this region (Table 5). These traditional practices may serve as important input for valid scientific investigation, standardization and large scale use in integrated pest management.

#### Acknowledgements

The authors are thankful to the large cardamom farmers of Sikkim and Darjeeling district of West Bengal, who provided valuable information and cooperation during the course of study. They are also thankful to IARI, New Delhi for the help to identify the insect pest.

#### References

- Gupta U, Chhetri P (2013) Performance of selected accessions of large cardamom germplasm under preliminary evaluation trial. *Green Farming Int J* 4: 670-671.
- Deka TN, Gudade BA, Saju KA, Vijayan, AK, Chhetri P (2014) Foraging behavior of Bumble bee (*Bombus breviceps*) in large cardamom (*Amomum subulatum* Roxb.) at Dzongu, North Sikkim. *Ecol Environ Conser* 20 : 197-199.
- Gudade BA, Harsha KN, Vijayan AK, Chhetri P, TN Deka, et al. (2015) Effect of soil application of Zn, Mn and Mg on growth and nutrient content in large cardamom (*Amomum subulatum* Roxb.) at Sikkim. *Int J Farm Sciences*: 51-55.
- Gupta U, Gudade BA, Chhetri P and Harsha KN (2012) Large cardamom- the lifeline in Sikkim. *Indian Horticulture* 57: 7-10.
- Rao YS, Gupta U, Kumar A, and Naidu RA (1993) A note on large cardamom (*Amomum subulatum* Roxb.) germplasm collection. *J Spic Arom Crop* 2: 77-80.
- Gudade BA, Chhetri P, Gupta U and Deka TN (2013) Establishment of large



**Plate 1:** Photographs of insect and pests of large cardamom at Sikkim Himalaya. (a) Large cardamom flowering clump; b. Congregation of *A. chorista*; c. Large cardamom leaf damaged by *A. chorista*; d. *M. dimorphus* infested shoot tip; e. Symptoms of *Glyphipterix* sp. infestation; f. *Mollitrichosiphum* sp. on *Alnus* twig.

- cardamom *Amomum subulatum* Roxb) sucker nursery at Sikkim. Popular Kheti 1: 1-3.
7. Gudade BA, Chhetri P, Gupta U, Deka TN, Vijayan AK, et al. (2014) The study of eco-friendly practices of large cardamom cultivation in Sikkim and Darjeeling. *Ecol Environ Conserv* 20: 119-123.
  8. Gudade BA, Chhetri P, Gupta U, Harsha KN (2013) More large cardamom under *Alnus*. *Indian Horticulture* 58: 19-20
  9. Gudade BA, Chhetri P, Deka TN, Gupta U, Vijayan AK (2013) Large cardamom (*Amomum subulatum* Roxb): a spice crop with multiple uses. *Indian J Arecanut Spices Med Plant* 15: 15-17.
  10. Vijayan AK, Gudade BA, Chhetri P, Deka TN (2014) Status of viral diseases of large cardamom (*Amomum subulatum* Roxb.) and its management in Sikkim and Darjeeling district of West Bengal. *J. Mycol. Pl. Pathol.*, 44 (4): 438-441
  11. Deka TN, Sudharshan MR, Saju KA (2011) New record of bumble bee, *Bombus breviceps* Smith as a pollinator of large cardamom. *Current Science* 100: 926- 928.
  12. Kishore K, Kalita H, Rinchen D, Pandey B (2012) Floral biology of large cardamom (*Amomum subulatum*). *Indian J Agricultural Sci* 82: 578-582
  13. Gupta U, Chhetri P, Gudade BA (2012) Collection, characterization, evaluation and conservation of germplasm of large cardamom. *Green Farming Int J* 3: 499-500.
  14. Subba JR (1984) *Agriculture in the Hills of Sikkim*. Sikkim Science Society, India
  15. Thakur Azad NS, Sachan JN (1987) Insect pests of large cardamom (*Amomum subulatum* Roxb) in Sikkim. *Bulletin of Entomology* 28: 46-58.
  16. Thakur Azad NS (1980) Pest complex on large cardamom in Sikkim. *Krishi Samachar* 3: 19-20.
  17. Gudade BA, Chhetri P, Gupta U, Deka TN, Vijayan AK (2013) Traditional practices of large cardamom cultivation in Sikkim and Darjeeling. *Life Sciences Leaflets* 9: 62-68.
  18. Gudade BA, Chhetri P, Gupta U, Harsha KN (2012) More large cardamom under *Alnus*. *Ind Horticulture* 58: 19-20.

### Author Affiliations

[Top](#)

*Indian Cardamom Research Institute, Regional Research Station, Spices Board, Tadong, Gangtok-737 102, Sikkim, India*

### Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 50 Journals
- ❖ 21 Day rapid review process
- ❖ 1000 Editorial team
- ❖ 2 Million readers
- ❖ Publication immediately after acceptance
- ❖ Quality and quick editorial, review processing

Submit your next manuscript at • [www.scitechnol.com/submission](http://www.scitechnol.com/submission)