



Pattern of Lipid Profile in Pregnancy and its Impact on the Gestational Course

Ashraf Reda¹, Awani Gamal¹, Mohamed Rezk² and Gehad Gamal^{3*}

¹Department of Cardiology, Faculty of Medicine, Menoufia University, Menoufia Governorate, Egypt

²Department of Obstetrics and Gynecology, Faculty of Medicine, Menoufia University, Menoufia Governorate, Egypt

³Department of Cardiology, Dar AlFouad Hospital, Giza, Egypt

*Corresponding author: Gehad Gamal, Department of Cardiology, Dar AlFouad Hospital, Giza, Egypt. E-mail: gehadgamal83@gmail.com

Received Date: November 02, 2018; Accepted Date: February 13, 2019; Published Date: February 20, 2019

Abstract

Background: Major changes in lipid profile occur during normal pregnancy with significant correlation could be detected with gestational complications.

Objective: We aimed to screen lipid profile in a sample of Egyptian pregnant and assess the correlation between second trimester maternal lipid profile and gestational outcome.

Methods: This cohort study included 94 women with maternal lipid profile assessed during the second trimester (16-18 weeks). Serial antenatal visits were conducted to record the development of maternal gestational hypertension, preeclampsia, gestational diabetes mellitus and low birth weight.

Results and discussion: Lipid profile components in pregnant is higher than normal references. Meanwhile, gestational hypertension (GH) was the most common complication in the study affecting 13 patients (13.8%) followed by preeclampsia (PE) in 8 patients (8.5%), gestational DM (GDM) in 7 patients (7.4%) and lastly low birth weight in 4 neonates (4.3%). Positive association between second trimester lipid profile components and GH, PE and GDM was found.

Conclusion: Measurement of lipid profile during the second trimester is highly recommended and could be a useful predictor of hypertensive disorders and GDM among Egyptian women.

Keywords: Lipid profile; Preeclampsia; Hypertensive disorders during pregnancy; Gestational diabetes mellitus

Introduction

Major changes in lipid profile occur during normal pregnancy with progressive increase in maternal plasma cholesterol and triglyceride levels during the late second and the third trimester, returning to normal after delivery [1-4].

Pregnancy-associated hyperlipidemia was linked with maternal morbidity with development of gestational diabetes mellitus and

preeclampsia [5-7]. To the authors' knowledge, there were no studies in the literature addressing lipid profile in pregnant women in Egypt. We aimed to screen the lipid profile pattern in the pregnant then assess the correlation between second trimester maternal lipid profile and adverse pregnancy outcome among cohort of Egyptian pregnant women.

Patients and Methods

Patient populations

This prospective study was conducted at the department of Cardiology in collaboration with the Obstetrics and Gynecology department, Menoufia University hospital, Menoufia governorate, Egypt during the period between June 2016 and June 2018. The study protocol was formally reviewed and approved by the ethics committee for human research at Menoufia Faculty of Medicine with informed consent obtained from all participants prior to commencement of the study after thorough explanation of the study objectives.

Inclusion criteria

Pregnant women attending the antenatal care clinic at Menoufia University hospital during early first trimester at 6-12 weeks gestation were invited to participate. Gestational age was based on the date of last menstrual period together with early obstetric ultrasound scan.

Exclusion criteria

- Women with preexisting medical disorders as diabetes mellitus and hypertension
- History of congenital fetal malformations or interrupted pregnancy (whether spontaneous or induced abortion)
- Pregnant with drug intake affecting lipid profile.

Methods

After exclusion of non-responders, drop out participants and those with exclusion criteria, 94 women completed the study (this number was considered suitable enough sample for statistical analysis with significant results and correlations).

Blood samples was drawn from all the subjects following a fast of 12 hours at 16-18 weeks' gestation and analyzed for Serum Triglycerides (TG), Total cholesterol (TC), LDL cholesterol and HDL cholesterol (HDL-C) by enzymatic methods auto analyzer. Participants were followed up at the antenatal care clinic throughout pregnancy with frequency of visits of 5 visits and delivered at Menoufia University hospital.

Outcome measures

Maternal outcome development of gestational hypertension (elevation of blood pressure $\geq 140/90$ mmHg after 20 weeks' gestation on two occasions with 4-hours apart after rest). The definition of hypertension in pregnancy is based only on office (or in-hospital) BP values [systolic BP (SBP) ≥ 140 mmHg and/or DBP ≥ 90 mmHg and distinguishes mildly (140-159/90-109 mmHg) or severely ($\geq 160/110$ mmHg) elevated BP.




Figure 2: Correlation between age (years) and total cholesterol (mg/dl).

Also, there is positive significant correlation between total cholesterol (TC) level and systolic blood pressure (Figure 3).




Figure 3: Correlation between systolic blood pressure SBP (mmHg) and total cholesterol (mg/dl).

While, there is positive significant correlation between (LDL) level and (SBP) (Figure 4).




Figure 4: Correlation between systolic blood pressure (mmHg) and low density lipoproteins (LDL, mg/dl).

In addition, there is positive significant correlation between (TG) level and age (Figure 5).




Figure 5: Correlation between age (years) and TG (mg/dl).

Also, there is positive significant correlation between (TG) and BMI (Figure 6).




Figure 6: Correlation between body mass index (BMI, kg/m²) and triglycerides (mg/dl).

Mean percent difference from the reference normal values in patient developed gestational DM were significantly higher than those without GDM: Cholesterol 44% (vs. 23% in those without GDM) and LDL 31% (vs. 3% in those without GDM) (P value 0.044 and 0.049 respectively) (Figures 7 and 8).




Figure 7: Percents of difference of total cholesterol from the normal value between cases who developed gestational DM (GDM) and who did not (P value 0.044).




Figure 8: Percents of difference of LDL from the normal value between cases who developed GDM and those who did not (P value 0.049).

As shown in Figures 9-11, we showed mean percent difference of lipid profile in patient developed gestational HTN were significantly higher than those without gestational HTN: Cholesterol 46% (21% in those without HTN), LDL 33% (vs. 1.1% in those without HTN) and TG 74% (vs. 42% in those without HTN) (P value 0.0001, 0.004 and 0.029 respectively).




Figure 9: Percents of difference of TC from the reference normal value between cases who developed gestational HTN (=pregnancy induced HTN) and who did not (P value 0.000).




Figure 10: Percents of difference of LDL from the reference normal value between cases who developed gestational HTN (=pregnancy induced HTN) and who did not (P value 0.004).




Figure 11: Percents of difference of TGs from the reference normal value between cases who developed pregnancy induced HTN (=gestational HTN) and who did not (P value 0.029).

We speculated mean percent difference of lipid profile in patient developed preeclampsia (PE) were significantly higher than those without: Cholesterol 48% (vs. 23% in those without) and LDL 41% (vs. 2.2% in those without) P value 0.003 and 0.006 respectively (Figures 12 and 13).




Figure 12: Percents of difference of TC from the reference normal Value between cases who developed preeclampsia (PE) and those who did not (P value 0.003).




Figure 13: Percents of difference of LDL from the reference normal value between cases who developed PE and those who did not (P value 0.006).

Discussion

The current study confirmed the previous associations between altered maternal lipid profile components and adverse maternal outcome. Second trimester elevated TC and LDL were found in women developed GDM. TC, TG and LDL were elevated in those developing gestational hypertension while elevated TC, LDL and high BMI were associated with the development of preeclampsia.

Maternal plasma lipids climb substantially above levels seen in normal pregnancies in women who developed preeclampsia or IUGR without maternal systemic manifestations of hypertension and proteinuria [8]. A more recent population-based study from China including 934 pairs of non-diabetic mothers and neonates between 2010 and 2011 with maternal lipid profile measured during the first, second and third trimesters had revealed that maternal high TG in late pregnancy was independently associated with increased risk of GDM [9].

Even though, pre-pregnancy altered maternal lipid profile was associated with subsequent development of GDM as depicted in recent nested case control study [10]. Bharathi et al., tested lipid profile in 100 Indian pregnant showed serum triglyceride, total cholesterol and VLDL level are significantly higher among woman with GDM compared to non GDM pregnant women [11].

A previous larger cohort study included 4008 non-diabetic women with early maternal lipid profile measured during early gestation (12-14 weeks) to detect the association between lipid profile and pregnancy-induced hypertension (PIH), had revealed that every unit increase in TG was linearly associated with an increased risk of PIH [odds ratio (OR)=1.60, P=0.021] while TC was not associated with any of the outcome measures [12]. Our results were concordant to Chandi et al., who recognized an association of maternal hypertriglyceridemia with pregnancy-induced hypertension. Among 59 women, a significant positive correlation was found between serum TG concentration and systolic blood pressure. Also, the mothers with hypertriglyceridemia were found to be at higher risk of developing early-onset gestational HTN [13].

Anaelechi et al. have evaluated the serum lipid profile and C-reactive protein (CRP) in 35 Nigerian women with pre-eclampsia and indicated that the women with pre-eclampsia had a significantly higher mean serum CRP (p=0.001), TG (p=0.001) and total cholesterol (p=0.001) level but a lower mean HDL-C (p=0.001) level than the controls [14].

The difference between our results and others can be explained on the basis of race, maternal age, nutrition, stage of pregnancy, socioeconomic, genetic factors and dietary pattern of individuals. From all these data, lifestyle modifications for women during the reproductive years should focus on lowering maternal triglyceride levels through diet, weight reduction and physical activity that may help in decreasing the maternal complications during pregnancy. Inability to measure maternal lipid profile during the three trimesters secondary to their high cost as well as to include larger cohort was unintended limitations of the current study. Future research should explore the implementation of maternal lipid profile and lifestyle modifications for the prevention of hypertensive disorders during pregnancy and other adverse outcome.

Conclusion

Maternal lipid profile components increase significantly in patients with hypertensive disorders and GDM among Egyptian women. Lifestyle modifications are required to prevent complications. Accordingly, we recommend Lipid profile should be done routinely during gestational course follow. Further studies with larger number of patients are required and, if possible, assessment of lipid profile pre pregnancy, during the three trimesters and after delivery (to compare all together).

References

1. Lippi G, Albiero A, Montagnana M, Salvagno GL, Scevarolli S (2007) Lipid and lipoprotein profile in physiological pregnancy. *Clin Lab* 53: 173–177.
2. Husain F, Latif S, Uddin M, Nessa A (2008) Lipid profile changes in second trimester of pregnancy. *Mymensingh Med J* 17: 17–21.

3. Charlton F, Tooher J, Rye KA (2014) Hennessy A: Cardiovascular risk, lipids and pregnancy: preeclampsia and the risk of later life cardiovascular disease. *Heart Lung Circ* 23: 203-212.
4. Bartels A, Egan N, Broadhurst DI, Khashan AS, Joyce C (2012) Maternal serum cholesterol levels are elevated from the 1st trimester of pregnancy: A cross-sectional study. *J Obstet Gynaecol* 32: 747-52.
5. Koukkou E, Watts GF, Lowy C (1996) Serum lipid, lipoprotein and Apo lipoprotein changes in gestational diabetes mellitus: a cross-sectional and prospective study. *J Clin Pathol* 49: 634-637.
6. Wiznitzer A, Mayer A, Novack V, Sheiner E, Gilutz H, et al. (2009) Association of lipid levels during gestation with preeclampsia and gestational diabetes mellitus: a population-based study. *Am J Obstet Gynecol* 201: 482.
7. Ferriols E, Rueda C, Gamero R, Vidal M, Payá A (2016) Relationship between lipid alterations during pregnancy and adverse pregnancy outcomes. *Clin Investig Arterioscler* 28: 232-244.
8. Sattar N, Greer IA, Galloway PJ, Packard CJ, Shepherd J (1999) Lipid and lipoprotein concentrations in pregnancies complicated by intrauterine growthrestriction. *J Clin Endocrinol Metab* 84: 128-130.
9. Jin WY, Lin SL, Hou RL, Chen XY, Han T (2016) Associations between maternal lipid profile and pregnancy complications and perinatal outcomes: a population-based study from China. *BMC Pregnancy Childbirth* 16: 60.
10. Vrijkotte TG, Krukziener N, Hutten BA, Vollebregt KC, Van EM (2012) Maternal lipid profile during early pregnancy and pregnancy complications and outcomes: the ABCD study. *J Clin Endocrinol Metab* 97: 3917-3925.
11. Bharathi KR, Vijayalakshmi S, Shrunga RP (2017) A study of lipid parameters among GDM and non GDM pregnant women: a hospital based study. *Int J Reprod Contracept Obstet Gynecol* 6: 5488-5490.
12. Han ES, Krauss RM, Xu F, Sridhar SB, Ferrara A (2016) Prepregnancy Adverse Lipid Profile and Subsequent risk of Gestational Diabetes. *J Clin Endocrinol Metab* 101: 2721-2727.
13. Chandi A, Sirohiwal D, Malik R (2015) Association of early maternal hypertriglyceridemia with pregnancy-induced hypertension. *Arch Gynecol Obstet* 292: 1135-1143.
14. Anaelechi JO, Japhet MO, John UU (2015) Evaluation of high-sensitivity c-reactive protein and serum lipid profile in southeastern nigerian women with Pre-Eclampsia, Departments of a Chemical Pathology and Obstetrics and Gynaecology, College of Health Sciences, Nnamdi Azikiwe University, Nnewi, Nigeria. *Med Princ Pract* 24: 276-279.