



Case report

Pisa Syndrome and Neurosyphilis: A Case Report in Taiwan

Ming-Kung Wu¹, Chih-Hsiang Chiu², Chun-Yi Su² and Kuan-Ying Hsieh^{2*}

Abstract

Background: Pisa syndrome, defined as more than 10 degrees tonic lateral flexion in upright spine without any significant associated vertebral rotation resembled the leaning tower of Pisa, is considered to be related to neuroleptics. Unfamiliarity of this disease, many neuro- and orthopedic surgeons can lead to unnecessary diagnostic and interventions.

Method: We report an extremely rare case of neurosyphilis presenting with PISA syndrome in order to assess the clinical presentation and treatment.

Results: A 49-year-old man, who has neurosyphilis and psychotic disorder due to general medical condition, treated by valproic acid 500 mg twice and quetiapine 200 mg once daily for as an unchanged regimen for the past 1 year. Five days after treatment with clozapine for poor controlled psychosis, he was noted leaned to one side. Physical examinations revealed a severe right truncal shift associated with left cervical tilt. We discontinued all medication except adding an anticholinergic drug (2 mg biperiden twice daily). After 7 days, the deformity disappeared with normal body posture. His psychosis flared up and we rechallenged clozapine 25mg daily. He was noticed leaned to one side again.

Conclusions: orthopedic and neurological surgeons visiting patients with abnormal postures of the trunk need to evaluate medications especially with neuropsychiatric disease. Even drugs with a minimum risk of extrapyramidal symptoms, such as clozapine, can cause Pisa syndrome in even short duration and low dosage.

Keywords

Pisa syndrome; Neurosyphilis

Introduction

Pisa syndrome (PS) is defined as more than 10 degrees of tonic lateral flexion in the upright spine without associated vertebral rotation, like the leaning tower of Pisa which is first described by Ekblom et al. in 1972 [1,2]. PS often considered related to longterm treatment with neuroleptics.

We report an extremely rare case of neurosyphilis presenting with PISA syndrome in order to assess the clinical presentation and treatment.

*Corresponding author: Kuan-Ying Hsieh, M.D.; Kaohsiung Municipal Kai-Syuan Psychiatric Hospital, 130, Kai-Syuan 2nd Rd., Ling-Ya District, Kaohsiung 802, Taiwan, Tel: +886-7-751-3171 ext. 2373; Fax: +886-7-16-1843; E-mail: isanirra@gmail.com

Received: October 14, 2017 Accepted: November 06, 2017 Published: November 12, 2017

Case Report

The 49-year-old man has neurosyphilis and psychotic disorder due to general medical condition since the age of 48. He had been admitted three times within a period of 1 years because of exacerbation of mood or psychotic episodes. The main presenting symptoms of each episode were aggression, religious delusion, thought of being controlled, and auditory hallucination. He was treated by valproic acid 500 mg twice and quetiapine 200 mg once daily for as an unchanged regimen for the past 1 year. His psychiatric symptoms were under controlled in this regimen without hospitalization.

However, he was admitted this time for persisted auditory and visual hallucination, physical and verbal aggression. We have titrated quetiapine dosage but muscle rigidity, sialorrhea, tilting to right side were noted. We then shift quetiapine gradually to clozapine 50mg daily for poor controlled psychosis and intolerable side effects.

Five days after treatment with clozapine, he was noticed progressively leaned to one side with a tonic right flexion of the trunk and left flexion of the neck but himself seemed unaware of and unconcerned by his body change. On physical examination, he had no evidence of dynamic (e.g., active spasms, posturing with jerks or tremor, paradoxical muscle groups associated with an abnormal posture or asymmetrical contractions of paraspinal muscles). Neurologic exam was unremarkable for motor and sensory deficits. Under the impression of PISA syndrome, we discontinued all medication except adding an anticholinergic drug (2 mg Biperiden twice daily). After 7 days, the deformity disappeared with normal body posture. His psychosis flared up and we rechallenged clozapine 25mg. He was noticed progressively leaned to one side again.

Discussion

The prevalence rate of PISA syndrome was reported to be 9.3% in women, 6.4% in men [3]. The risk factors were female sex, old age, organic brain disorders and a history of treatment with conventional neuroleptics [4]. The pathophysiology of PS is related to dopaminergic system.

PS has been described in several diseases including Parkinson disease, progressive supranuclear palsy, and Huntington disease [5-7]. PS was also described as a side effect of neuroleptics such as risperidone, quetiapine, ziprasidone, paliperidone, aripiprazole, and cholinesterase inhibitors [8-13]. Clozapine, has lower binding affinity to Dopamine 2 receptors, seems less risk of PS. Furthermore, clozapine is also reported to be a treatment of PS [14].

To our knowledge, this is the first case report of PISA syndrome in neurosyphilis. The patient has the risk factors as organic brain disorders and treated with antipsychotics. PS occurs five days after initiating clozapine treatment in a low dosage of 50mg daily and improved after discontinuation. Relapse of PS after rechallenging clozapine in a lower dosage is also noted.

In summary, orthopedic and neurological surgeons visit patients with abnormal postures of the trunk need to evaluate medications especially with neuropsychiatric disease. Even drugs with a minimum risk of extrapyramidal symptoms, such as clozapine, can cause Pisa syndrome in short duration and low dosage.

References

1. Suzuki T, Matsuzaka H (2002) Drug-induced Pisa syndrome (pleurothotonus): epidemiology and management. *CNS Drugs* 16: 165e74.
2. Ekblom K, Lindholm H, Ljungberg L (1972) New dystonic syndrome associated with butyrophenone therapy. *Z Neurol* 202: 94e103.
3. Yassa R, Nastase C, Cvejic J (1991) The Pisa syndrome (or pleurothotonus): Prevalence in a psychogeriatric population. *Biol Psychiatry* 29: 942–945.
4. Stübner S, Padberg F, Grohmann R (2000) Pisa syndrome (pleurothotonus): report of a multicenter drug safety surveillance project. *J Clin Psychiatry* 61: 569–574.
5. Villarejo A, Camacho A, García-Ramos R (2003) Cholinergic-dopaminergic imbalance in Pisa syndrome. *Clin Neuropharmacol* 26: 119-121.
6. Solla P, Cannas A, Costantino E (2012) Pisa syndrome in a patient with progressive supranuclear palsy. *J Clin Neurosci* 19: 922-923.
7. Salazar Z, Tschopp L, Calandra C (2008) Pisa syndrome and parkinsonism secondary to valproic acid in Huntington's disease. *Mov Disord* 23: 2430-2431.
8. Cordeiro Q, Zung S, Vallada H (2008) Pisa syndrome induced by rapid increase and high dosage of risperidone. *Arq Neuropsiquiatr* 66: 896e7.
9. Walder A, Greil W, Baumann P (2009) Drug-induced Pisa syndrome under quetiapine. *Prog Neuropsychopharmacol Biol Psychiatry* 33: 1286e7.
10. Yeh YM, Chen YC, Chen CK (2009) Chronic Pisa syndrome associated with switching antipsychotics from olanzapine to ziprasidone. *Prog Neuropsychopharmacol Biol Psychiatry* 33: 162e3.
11. Pan PY, Chou HW, Cheng CC (2014) Pisa syndrome induced by paliperidone. *J Neuropsychiatry Clin Neurosci* 26: E07e8.
12. Chen HK, Wu BJ, Shao CH (2010) Drug-induced Pisa syndrome associated with aripiprazole during clozapine treatment. *Prog Neuropsychopharmacol Biol Psychiatry* 34: 707e8.
13. Cossu G, Melis M, Melis G (2004) Reversible Pisa syndrome (pleurothotonus) due to the cholinesterase inhibitor galantamine: case report. *Mov Disord* 19: 1243e4.
14. Hazari N, Kate N, Grover S (2013) Clozapine and tardive movement disorders: a review. *Asian J Psychiatr* 6: 439e51.

Author Affiliation

Top

¹Department of Psychiatry, Kaohsiung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Kaohsiung, Taiwan

²Kaohsiung Municipal Kai-Syuan Psychiatric Hospital, Taiwan

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000  fans
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission