



Presentation and Management of Erosive Pustular Dermatitis Involving the Upper Limb

Zhao Huang*

Introduction

Erosive pustular dermatitis (EPD) is a cutaneous inflammatory condition, characterized by chronically non-healing lesions with sterile pustules and hyperkeratotic crusted erosions. Although uncommon, extreme cases have been alluded by dermatologists to plastic specialists for the board. The point of this paper was to portray the show and the board of broad EPD just as a survey of current writing on this uncommon condition.

An instance of assumed EPD of twelve years length was introduced. Broad nature of the infection in a man with foundation history of constant local agony condition required reference to plastic medical procedure administration from dermatology. This difficult case included a malnourished patient with broad space of his upper appendage skin influenced by hyperkeratotic crusting disintegrations with encompassing atrophic changes and joint contractures. After starting debridement and tissue biopsy to reject danger, patient was saved inpatient for day by day dressing changes and effective intense corticosteroid application with help of sedative assistance. Critical improvement of the condition was clear with sequential photographic documentation records [1-3].

Erosive pustular dermatitis, albeit seldom alluded to plastic medical procedure, ought to be considered in the differential conclusion of a non-mending wound. It should resolve promptly with broadened course of effective powerful corticosteroid treatment. EPD is a cutaneous fiery condition, portrayed by constantly non-mending sores with sterile pustules and hyperkeratotic crusted disintegrations. Broad or contaminated infection might result in huge scarring and contracture. Albeit moderately uncommon, extreme cases have been eluded by dermatologists to plastic specialists for the board.

Under 200 cases have been accounted for in writing since late 1970's, when it was first depicted, with most cases including the scalp with actinic changes or the legs with venous inadequacy. We portray here an instance of assumed EPD including most of an upper appendage of twelve years length, related with upper appendage complex provincial torment condition [4].

Differential History of Melanoma or Multiple Nevi

EPD is a somewhat under-analysed cutaneous incendiary infection of obscure etiology. Since its first depiction in the last part

*Corresponding author: Zhao Huang, Department of Dermatology, Beijing Hospital of Traditional Chinese Medicine, Capital Medical University, Beijing, China, E-mail: huang@126.com

Received: August 03, 2021 Accepted: August 18, 2021 Published: August 25, 2021

of the 1970's, there are just roughly 160 cases detailed in the writing. It has been portrayed to for the most part influence either the scalp or the leg.

EPD of the scalp is a condition announced for the most part in older patients who have had delayed sun openness, set off by a nearby injury like injury, skin malignant growth, medical procedure, and so forth.

EPD of the leg has been depicted as an uncommon structure in patients with longstanding venous inadequacy and balance skin changes.

Clinical components of EPD are erythematous, pustular sores with crusted disintegrations. Histology typically shows vague persistent aggravation with neutrophilic penetration, as is displayed for our situation. The typical misdiagnoses are disease or harm. These EPD skin sores are quite often impervious to neighbourhood or fundamental anti-microbial treatment, however the condition for the most part reacts to strong skin steroid specialists [5]. As of late EPD has been depicted in patients with non-mending consume wounds. It, in any case, has never been accounted for in the setting of atrophic skin changes identifying with CRPS which is certifiably not an exceptional element among patients of the plastic/hand specialists.

Our unique case has shown some trademark clinical and histo-obsessive components reminiscent of EPD, and other dermatological findings have been prohibited by histo-clinical connection. We theorize that the tactile and atrophic skin changes related with CRPS might be inclined to aggravations and consequently bringing about improvement of this ongoing debilitating condition. The historical backdrop of conceivable self-scouring/disturbing the all-around delicate skin may likewise be connected.

Albeit a difficult case at time, our patient's condition had reacted well to a drawn out course of skin corticosteroid treatment while inpatient. It additionally had exhibited the probability of sickness backslide if rebellious to recommended therapy. Multi-disciplinary group inputs are the key in the administration of these patients [6,7].

Conclusion

Erosive pustular dermatitis, although rarely referred to plastic medical procedure, should be considered in the differential determination of a non-recuperating wound. It should resolve promptly with broadened course of effective powerful corticosteroid treatment. Patients ought to be educated in regards to the significance of treatment consistence and the potential for relapse.

References

- Burton JL (1977) Case for diagnosis. Pustular dermatitis of scalp. Br J Dermatol 97 Suppl 15: 67-69.
- Pye RJ, Peachey RD, Burton JL (1979) Erosive pustular dermatitis of the scalp. Br J Dermatol 100: 559-566.
- Semkova K, Tchernev G, Wollina U (2013) Erosive pustular dermatitis (chronic atrophic dermatitis of the scalp and extremities). Clin Cosmet Investig Dermatol 6: 177-182.
- Grattan CE, Peachey RD, Boon A (1988) Evidence for a role of local trauma in the pathogenesis of erosive pustular dermatitis of the scalp. Clin Exp Dermatol 13: 7-10.
- Levakov O, Gajic B (2014) Erosive pustular dermatitis of the scalp-is it really a rare condition? Vojnosanit Pregl 71: 307-310.

6. Brouard MC, Prins C, Chavaz P, Saurat JH, Borradori L (2002) Erosive pustular dermatosis of the leg: report of three cases. *Br J Dermatol* 147: 765-769.
7. Cotterill JA, Lanigan SW (1990) Erosive pustular dermatosis of the leg—a definition. *Br J Dermatol* 123: 548.

Author Affiliations

[Top](#)

Department of Dermatology, Beijing Hospital of Traditional Chinese Medicine, Capital Medical University, Beijing, China