



Quantitative Coronary Angiography for Assessment of Non-Obstructive Coronary Artery Disease: Comparison to Fractional Flow Reserve

Mahmoud Kamel Ahmed*, Ahmed Abdel Azeez Emara, Morad Beshay Mena and Mostafa Abdel Aty Melegy

Abstract

Objective: To assess intermediate coronary artery lesions by 2D and 3D Quantitative Coronary Angiography (QCA) and compare the results with standard fractional flow reserve (FFR).

Background: FFR is the standard for assessment of physiological significance of an intermediate coronary stenosis, but it expensive, sometimes it is unavailable, or limited use by contraindications to adenosine, QCA especially three-dimensional (3D-QCA) can be used as another method to help in assessment of the intermediate coronary lesions.

Patients and methods: Thirty two vessels in 30 patients with intermediate coronary lesions were scheduled for FFR, 2D and 3D QCA measurement and the results of QCA were compared and correlated to the obtained from FFR. Results: The studied group included 32 vessels, 24 diabetic patients (75%). 18 hypertensive (56.3%). 20 smokers (62.5%). Mean FFR value was 0.80 ± 0.13 . FFR ≥ 0.80 was observed in 18 lesions (56.25%). Lesions severity obtained by 3D-QCA was better correlated to FFR than 2D-QCA. Both 3D area stenosis percent and 3D diameter stenosis percent have comparable correlation in term of accuracy but with better sensitivity for the percent diameter stenosis. 2D-QCA measurements were less correlated to FFR.

Conclusion: 3D QCA is better than 2D QCA in assessment of intermediate coronary lesions and it is better correlated with FFR so it may be used in assessment of intermediate coronary lesions, when FFR is unavailable or contraindicated. Key words: Quantitative Coronary Angiography (QCA), Fractional Flow Reserve (FFR), Intermediate coronary lesions, Functional severity of coronary stenosis.

Keywords

Quantitative coronary angiography (QCA); Fractional flow reserve (FFR); Intermediate coronary lesions; Functional severity of coronary stenosis

Introduction

The presence of ischemic myocardium is a significant risk factor for unfavorable clinical outcome. Revascularization of

*Corresponding author: Mahmoud Kamel Ahmed, Department of Cardiology, Menoufia University, Menoufia Governorate, Egypt, Tel: 01006945404; E-Mail: mahmoudkamel35@gmail.com

Received: August 24, 2018 Accepted: September 13, 2018 Published: September 18, 2018

obstructive coronary lesions that induce ischemia can improve a patient's functional condition and outcome [1]. For lesions with stenosis that do not induce ischemia, however, the beneficial result of revascularization is less clear, and pharmacological therapy alone is likely to be effective [2]. With the invention of drug-eluting stents (DES), the number of patients with multi-vessel coronary artery disease in whom invasive percutaneous coronary intervention (PCI) is performed has progressively increased [3]. However, in patients with multi-vessel coronary artery disease (MVD), determining which lesions cause ischemia and deserve stenting can be challenging [9] (Table 1).

Noninvasive stress imaging studies has limited ability to accurately confine ischemia-producing lesions in these patients [4]. Although coronary angiography may underestimate/overestimate a lesion's functional severity, it is still the standard technique to guide PCI in patients with MVD [5]. Fractional flow reserve (FFR) is a guide of the physiological significance of a coronary stenosis and is defined as the ratio of maximal blood flow in a stenotic artery to normal maximal flow [6]. It can be assessed during coronary angiography by calculating the ratio of distal coronary pressure measured with a coronary pressure guide wire to aortic pressure measured simultaneously with the guiding catheter. FFR in a normal coronary artery equals 1.0. An FFR value of 0.80 or less identifies ischemia-causing coronary stenosis with an accuracy of more than 90% [6,7] (Table 2). The data provided by FFR is similar to that gained with myocardial perfusion studies, but it is more specific and has a better spatial resolution, because each artery or segment is separately analyzed [4,8] Deferring PCI in non-significant stenotic lesions as assessed by FFR is associated with an annual rate of death or myocardial infarction of approximately 1% in patients with single-vessel coronary artery disease, which is lower

Table 1: Personal and clinical data of the studied group.

| Personal data | Total studied group N=32 | |
|-------------------|--------------------------|---------|
| Age mean \pm SD | 60.8 \pm 6.9 | |
| Sex | NO | % |
| Male/Female | 24/8 | 75%/25% |
| DM | 24 | 75% |
| HTN | 18 | 56.3% |
| Smoker | 20 | 62.5% |
| HDL | 32.8 \pm 6.3 | |
| LDL | 128.6 \pm 35.2 | |

Table 2: Coronary stenosis assessment by FFR vs. area stenosis% and diameter stenosis% seen by 2D angiography.

| Personal data | FFR assessment | |
|-------------------------------------|----------------|--------------|
| | Diseased | Not diseased |
| 2D assessment of area stenosis% | | |
| Positive | (a) 10 | (b) 10 |
| Negative | (c) 8 | (d) 4 |
| 2D assessment of diameter stenosis% | | |
| Positive | (a) 10 | (b) 6 |
| Negative | (c) 8 | (d) 8 |

a=True positive cases b=False positive c=False negative d=True negative

in comparison to the rate after routine stenting [2] On the other hand, the deferring PCI in lesions with an FFR of less than 0.75 to 0.80 may lead to worse outcomes than those who do revascularization [9] Retrospective studies suggest that in patients with multi-vessel coronary artery disease, FFR-guided PCI is associated with a favorable outcome with respect to event-free survival [10] QCA can be used as another method to help assessment of the intermediate coronary lesions.

Three-dimensional quantitative coronary angiography (3D-QCA) uses standard images acquired during routine coronary angiography to reconstruct a 3D model of a coronary artery by fusing two or more orthogonal angiographic images. 3D-QCA reportedly allows a more accurate representation of true vessel geometry when compared with standard two-dimensional (2D) QCA in phantom models [11] and has been tested against intravascular ultrasound (IVUS) [12] Although IVUS currently provided the most accurate measurements of vessel geometry and lesion severity [13] 3D-QCA measurements can be obtained by using the existing standard coronary angiography images without the need for additional time or equipment during the procedure [14].

The aim of this study is to compare the result of QCA with the results of FFR in assessment of intermediate coronary lesion (Table 3).

Patients and Methods

The study prospectively included 34 patients performing percutaneous coronary angiography with/without revascularization due to known or suspected coronary artery disease in International Medical Center (IMC, Cairo) during the period between May 2015 and December 2016. All patients had intermediate coronary artery disease that can be managed both medically and by revascularization, so the ischemic objective evidence is needed to justify treatment method and help decision making for either PCI or medical treatment [15].

Exclusion criteria

Patient were excluded if they have any of the following: Serial stenosis that may complicate the measurement of FFR, as in diffusely diseased vessels or successive nearby stenosis, to the extent that no adequate healthy area where the pressure wire sensor can be advanced to it, lesions with known infarction in the target vessel territory (because microvascular perturbations from myocardial infarction cause discordance between the FFR and angiographic stenosis

severity), left main coronary stenosis (due to differences between the severity left main stenosis and functional significance).

Some mild insignificant lesions in coronaries may be functionally significant if it occurs with the same percentage of stenosis in left main and myocardial bridge, as this not a true atherosclerotic lesion and medical treatment is the treatment of option. All patients had provided informed consent and all were subjected to full history taken, thorough clinical examination, ECG and routine laboratory testing (CBC, renal function and lipid profile).

Coronary angiography

Diagnostic coronary angiography using the routine angiographic projections, in addition to dedicated projections if needed. Assessment of non-obstructive coronary lesions by 2D-QCA. An image that shows the specified coronary vessel with least lesion foreshortening is chosen. After calibration of the used catheter, drawing a line into the vessel, starting from before the lesion and ending after it including the lesion itself (from healthy to healthy).

Assessment of non-obstructive coronary lesions by 3D quantitative coronary angiography

The machine is set to 3D mode and the patient position adjusted midway between the C-arm ends with insuring that the C-arm has a clear filed of moving with no obstacles. 20 ml of contrast agent in the lower lock syringe should be ready for injection (Table 4).

For both 2D and 3D the reference vessel diameter, minimal luminal diameter, percent of diameter stenosis, minimal luminal area, and percent of area stenosis, and lesion length are calculated automatically. Assessment of the same lesions by fractional flow reserve: The pressure wire (PW) is opened in a sterile technique and its distal end (two types of pressure wires are commercially available; Volcano Corporation (San Diego, Calif) and St Jude Medical, Inc (St.

Paul, Minn) is connected to a set/analyzer that displays the intracoronary pressure, aortic pressure and can measure the pressure distal to the lesion (Pd)/pressure proximal to the lesion (which is same the aortic pressure) (Pa). Once the sensor of the PW is distal to the stenosis, a hyperemic stimulus is administered by injection through the guide catheter, and the FFR is monitored for a significant change. To achieve maximum hyperemia, adenosine is typically used: a 100 µg bolus in the right coronary artery, a 200 µg bolus in the left coronary artery.

Table 3: Validity of 2D angiography to diagnose coronary stenosis according to percent of area stenosis.

| | AUC | P value | Sensitivity | Specificity | Accuracy | PPV | NPP |
|------------------------|-------|---------|-------------|-------------|----------|-----|-----|
| Area by 2D angiography | 0.421 | 0.597 | 56% | 29% | 44% | 55% | 33% |

Table 4: Validity of 2D angiography to diagnose coronary stenosis according to percent of diameter stenosis.

| | AUC | P value | Sensitivity | Specificity | Accuracy | PPV | NPP |
|----------------------------|-------|---------|-------------|-------------|----------|-----|-----|
| Diameter by 2D angiography | 0.492 | 0.958 | 56% | 57% | 56% | 63% | 50% |

Table 5: Coronary stenosis assessment by FFR vs. area stenosis% and diameter stenosis% seen by 3D angiography.

| Personal data | FFR assessment | |
|-------------------------------------|----------------|--------------|
| | Diseased | Not diseased |
| 3D assessment of area stenosis% | | |
| Positive | (a) 14 | (b) 4 |
| Negative | (c) 4 | (d) 10 |
| 3D assessment of diameter stenosis% | | |
| Positive | 1. (a) 16 | 2. (b) 6 |
| Negative | (c) 2 | (d) 8 |

Pd/Pa is displayed on the analyzer; if it <0.8 this means that the lesion is functionally significant and justifies revascularization. PCI can be done, if needed, through the same PW by disconnecting it from the analyzer and using it directly as a usual PTCA guide wire.

Intra- and Inter-observer variation analysis

For intra-observer variability, a single operator repeats analysis of QCA image (2D&3D) in 10 patients (one month apart). For inter-observer variation analysis, two independent operators did the same procedure for QCA measurement separately in 20 patients.

Results

The study included 34 studied patients, 4 were excluded because of the poor image quality and difficult obtaining the data and two patients have two vessels lesions studied, so the total number of studied vessels is 32 vessels.

Of the 32 studied vessels there were 24 vessels in male patients (75%), 8 vessels in female patients (25%) with mean age 60.8 ± 6.9 . Of the studied vessels, left anterior descending (LAD) constitute 50% of the affected vessels, while left circumflex (LCX) and right coronary artery (RCA) constitute the remaining vessels equally (25% for each). For statistical purposes; each vessel will be represented as single case.

The studied group included 24 diabetic patients (75%). 18 hypertensive (56.3%). 20 smokers (62.5%). The mean HDL level was 32.8 ± 6.3 mg/L and the LDL level was 128.6 ± 35.2 mg/L. Mean FFR value was 0.80 ± 0.13 . FFR 0.80 was observed in 18 lesions (56.25%). Lesions severity obtained by 3D-QCA was better correlated to FFR than 2D-QCA.

Both 3D area stenosis percent and 3D diameter stenosis percent have comparable correlation in term of accuracy but with better sensitivity for the percent diameter stenosis. 2D-QCA measurements were less correlated to FFR. Overall, 3D-QCA showed a non-significant trend towards more accurate prediction of FFR than 2D-QCA. Both 3D- and 2D-QCA were less accurate in predicting FFR? 0.80 than in predicting FFR, 0.75 (Table 5).

Intra-observer and inter-observer variation analysis for QCA were assessed and the difference between two measurements was

found to be located within 95% confidence interval so that P value had more than 0.05.

Two dimensional stenosis percent (area stenosis % and diameter stenosis %)

I-Area stenosis percent: The number of truly diagnosed cases by 2D area % is 14 cases (10 true positive, 4 true negative) which represent 43.75% of total cases (Table 6).

While the number of falsely diagnosed cases by 2D area % is 18 (10 false positive, 8 false negative) which represent 56.25% of total cases. The best cut off point to diagnose coronary artery stenosis by 2D area % is 69.3% with 56% sensitivity and 29% specificity, 55% PPV and 33% NPV.

II- Diameter stenosis percent: The number of truly diagnosed cases by 2D diameter stenosis % is 18 cases (10 true positive, 8 true negative) which represent 56.25% of total cases. While the number of falsely diagnosed cases by 2D diameter stenosis% is 14 (6 false positive, 8 false negative) which represent 43.75% of total cases (Table 7). The cutoff point to diagnose coronary artery stenosis by 2D diameter stenosis % is 51.8% with 56% sensitivity and 57% specificity, 63% PPV and 50%.

Three dimensional area stenosis percent (area stenosis % and diameter stenosis %)

Area stenosis percent: The number of truly diagnosed cases by 3D area stenosis % is 24 cases (14 true positive, 10 true negative) which represent 75% of total cases. While the number of falsely diagnosed cases by 2D diameter stenosis% is 8 (4 false positive, 4 false negative) which represent 25% of total cases with 66.4% as a cutoff point to diagnose coronary artery with 78% sensitivity and 71% specificity.

Diameter stenosis percent: The number of truly diagnosed cases by 3D diameter stenosis % is 24 cases (16 true positive, 8 true negative) which represent 75% of total cases. While the number of falsely diagnosed cases by 3D diameter stenosis% is 8 (6 false positive, 2 false negative) which represent 25% of total cases. With 47.4% cut off point to diagnose coronary artery stenosis by 3D diameter stenosis % is 47.4% with 89% sensitivity and 57% specificity.

Table 6: Validity of 3D angiography to diagnose coronary stenosis according to percent of area stenosis.

| | AUC | P value | Sensitivity | Specificity | Accuracy | PPV | NPP |
|-----------------------------|-------|---------|-------------|-------------|----------|-----|-----|
| Area seen by 3D angiography | 0.730 | 0.125 | 78% | 71% | 75% | 78% | 71% |

Table 7: Validity of 3D angiography to diagnose coronary stenosis according to percent of diameter stenosis.

| | AUC | P value | Sensitivity | Specificity | Accuracy | PPV | NPP |
|---------------------------------|-------|---------|-------------|-------------|----------|-----|-----|
| Diameter seen by 3D angiography | 0.611 | 0.459 | 89% | 57% | 75% | 73% | 80% |

Table 8: Correlation between FFR and QCA.

| | | FFR |
|-----------------------|--------------|------------|
| 3D area stenosis% | r (Spearman) | -0.320 |
| | p | 0.074 (NS) |
| 3D diameter stenosis% | r (Spearman) | -0.358 |
| | p | 0.044 (S) |
| 2D area stenosis% | r (Pearson) | -0.032 |
| | p | 0.861 (NS) |
| 2D diameter stenosis% | r (Pearson) | 0.013 |
| | p | 0.943 (NS) |

Discussion

In patients with stable coronary artery disease, it was unclear whether an initial management strategy by percutaneous coronary intervention (PCI) with intensive medical therapy is superior to optimal medical therapy alone in reducing the cardiovascular risk events [15]. A meta-analysis of randomized clinical trials that studied PCI versus OMT in stable CAD showed that PCI, as compared with OMT, did not reduce the mortality risk, cardiovascular death, nonfatal myocardial infarction, or revascularization. PCI, however, provided a greater ischemic pain relief compared with OMT alone [16]. Considering the lack of clear clinical benefit, in addition to cost implication these findings continue to support existing clinical practice guidelines that medical therapy be considered the most appropriate initial clinical management for patients with stable angina [16].

The assessment of lesion severity was firstly depending on visual estimation of the operator, which may be inaccurate with marked difference between assumed lesion severity and physiological significance. Angiographic-physiological mismatch is frequent in patients with moderate coronary stenosis, which suggests the clinical importance of using physiological assessment to guide PCI [17]. A meta-analysis [18] showed that SPECT, CMR, and PET all have a high sensitivity, while a broad range of specificity was observed.

SPECT is widely available also it is extensively validated; PET achieved the highest diagnostic performance; CMR may provide another tool without ionizing radiation and with a similar diagnostic accuracy as PET [19,20]. The FAME study assessed the FFR-guided PCI and concluded the lack of accuracy of angiography in assessing the functional significance of a coronary stenosis when compared with the FFR, not only in the 50% to 70% category but also in the 70% to 90% of angiographic severity category [21], 2 years [22] and 5 years [23] follow up of the study results confirm the long-term safety of FFR-guided PCI in patients with multi-vessel disease. However, despite its accuracy and being helpful in intermediate lesion physiological significance assessment, the use of both techniques is limited by its relatively high cost.

In a study by Naganuma et al. [19] showed that the accuracy of QCA in predicting functionally significant FFR is dependent on FFR cut-off used and lesion severity. It suggested that where FFR is not available or contraindicated; 3D-QCA may help in the evaluation of coronary lesions of intermediate severity. Saad et al. [24] showed 3-D QCA showed a significant correlation with FFR values. A cross-sectional stenosis >57% obtained by 3-D QCA has a high degree of sensitivity and specificity to detect a hemodynamically significant intermediate coronary stenosis [25]. Recently Nishi et al. [26], found that that 3D-QCA is more useful than 2D-QCA and possibly comparable to IVUS in the assessment of functional stenosis severity. The study suggested that when FFR is not available, 3D-QCA MLA and MLD may assist in the assessment of functional severity of intermediate lesions. Another prospective, head-to-head study (ATLANTA I and II) [27] comparing QCA, quantitative CTA, and IVUS for the prediction of hemodynamic significance in intermediate and severe lesions, using fractional flow reserve as reference standard showed in intermediate-to-severe lesions, QCA-, CTA-, and IVUS-derived quantitative anatomic measurements correlated with FFR (Table 8). CTA-derived cut-points were similar to respective measurements on QCA and IVUS and had similar or better diagnostic performance compared with IVUS. In our study we found that the best cut off point to diagnose coronary artery stenosis

is 66.4%, as assessed by 3D-QCA, above which or equal to it there is significant stenosis.

Conclusion

2D and 3D QCA can be used for assessment of intermediate coronary lesions, when FFR is unavailable or contraindicated to assess the need for revascularization. Study limitations: Patients with suboptimal image quality were excluded, who, in practice, constitute a proportion of patients with CAD. Also, the unavailability or relatively expensiveness of the pressure wires and finally the small sample size were the main limitations in our study.

Study Limitations

Patients with suboptimal image quality were excluded, whom, in practice, constitute a proportion of patients with CAD. Also, the unavailability or relatively expensiveness of the pressure wires and finally the small sample size were the main limitations in our study.

References

1. Erne P, Schoenenberger AW, Burckhardt D, Zuber M, Kiowski W, et al. (2007) Effects of percutaneous coronary interventions in silent ischemia after myocardial infarction: The SWISSI II randomized controlled trial. *JAMA* 297: 1985-1991.
2. Pijls NH, Van Schaardenburgh P, Manoharan G, Boersma E, Bech JW, et al. (2007) Percutaneous coronary intervention of functionally nonsignificant stenosis: 5-year follow-up of the DEFER Study. *J Am Coll Cardiol* 49: 2105-2011.
3. Ong AT, van Domburg RT, Aoki J, Sonnenschein K, Lemos PA, et al. (2006) Sirolimus-eluting stents remain superior to bare-metal stents at two years: Medium-term results from the Rapamycin-Eluting Stent Evaluated at Rotterdam Cardiology Hospital (RESEARCH) registry. *J Am Coll Cardiol* 47: 1356-1360.
4. Lima RS, Watson DD, Goode AR, Siadaty MS, Ragosta M, et al. (2003) Incremental value of combined perfusion and function over perfusion alone by gated SPECT myocardial perfusion imaging for detection of severe three-vessel coronary artery disease. *J Am Coll Cardiol* 42: 64-70.
5. Fischer JJ, Samady H, McPherson JA, Sarembock IJ, Powers ER, et al. (2002) Comparison between visual assessment and quantitative angiography versus fractional flow reserve for native coronary narrowing of moderate severity. *Am J Cardiol* 90: 210-205.
6. Pijls NH, De Bruyne B, Peels K (1996) Measurement of fractional flow reserve to assess the functional severity of coronary-artery stenoses. *N Engl J Med* 334: 1703-1708.
7. De Bruyne B, Pijls NH, Bartunek J (2001) Fractional flow reserve in patients with prior myocardial infarction. *Circulation* 104: 157-162.
8. Pijls NH (2004) Optimum guidance of complex PCI by coronary pressure measurement. *Heart* 90: 1085-1093.
9. Legalery P, Schiele F, Seronde MF (2005) One-year outcome of patients submitted to routine fractional flow reserve assessment to determine the need for angioplasty. *Eur Heart J* 26: 2623-2629.
10. Berger A, Botman KJ, MacCarthy PA (2005) Long-term clinical outcome after fractional flow reserve-guided percutaneous coronary intervention in patients with multivessel disease. *J Am Coll Cardiol* 46: 438-442.
11. Ramcharitar S, Daeman J, Patterson M, van Guens RJ (2008) First direct *in vivo* comparison of two commercially available three-dimensional quantitative coronary angiography systems. *Catheter Cardiovasc Interv* 71: 44-50
12. Schuurbijs JC, Lopez NG, Ligthart J, Gijzen FJ (2009) *In vivo* validation of CAAS QCA-3D coronary reconstruction using fusion of angiography and intravascular ultrasound (ANGUS). *Catheter Cardiovasc Interv* 73: 620-626.
13. Takayama T, Hodgson JM (2001) Prediction of the physiologic severity of coronary lesions using 3D IVUS: validation by direct coronary pressure measurements. *Catheter Cardiovasc Interv* 53: 48-55.

14. Gradaus R, Mathies K, Breithardt G, Bocker D (2006) Clinical assessment of a new real time 3D quantitative coronary angiography system: Evaluation in stented vessel segments. *Catheter Cardiovasc Interv* 68: 44-49.
15. Pursnani S, Korley F, Gopaul R, Kanade P (2012) Percutaneous coronary intervention versus optimal medical therapy in stable coronary artery disease: A systematic review and meta-analysis of randomized clinical trials. *Circ Cardiovasc Interv* 5: 476-490.
16. Thomas S, Gokhale R, Boden WE, Devereaux PJ (2013) A meta-analysis of randomized controlled trials comparing percutaneous coronary intervention with medical therapy in stable angina pectoris. *Can J Cardiol* 29: 472-482.
17. Nakamura M, Yamagishi M, Ueno T, Hara K, Ishiwata S (2014) Prevalence of visual-functional mismatch regarding coronary artery stenosis in the CVIT-DEFER registry. *Cardiovasc Interv Ther* 29: 300-308
18. Jaarsma C, Leiner T, Bekkers SC, Crijns HJ (2012) Diagnostic performance of noninvasive myocardial perfusion imaging using single-photon emission computed tomography, cardiac magnetic resonance, and positron emission tomography imaging for the detection of obstructive coronary artery disease: A meta-analysis. *J Am Coll Cardiol* 59: 1719-1728.
19. Naganuma T, Latib A, Costopoulos C, Takagi K, Naim C (2014) The role of intravascular ultrasound and quantitative angiography in the functional assessment of intermediate coronary lesions: correlation with fractional flow reserve. *Cardiovasc Revasc Med* 15: 3-7.
20. Yong AS, Ng AC, Brieger D, Lowe HC (2011) Three-dimensional and two-dimensional quantitative coronary angiography, and their prediction of reduced fractional flow reserve. *Eur Heart J* 32: 345-353.
21. Tonino PA, Fearon WF, De Bruyne B, Oldroyd KG (2010) Angiographic versus functional severity of coronary artery stenoses in the FAME study fractional flow reserve versus angiography in multivessel evaluation. *J Am Coll Cardiol* 55: 2816-2821.
22. Pijls NH, Fearon WF, Tonino PA, Siebert U (2010) FAME Study Investigators Fractional flow reserve versus angiography for guiding percutaneous coronary intervention in patients with multivessel coronary artery disease: 2-year follow-up of the FAME (Fractional Flow Reserve Versus Angiography for Multivessel Evaluation) study. *J Am Coll Cardiol* 56: 177-184.
23. Van Nunen LX, Zimmermann FM, Tonino PA, Barbato E (2015) FAME Study Investigators Fractional flow reserve versus angiography for guidance of PCI in patients with multivessel coronary artery disease (FAME): 5-year follow-up of a randomised controlled trial. *Lancet* 386: 1853-1860.
24. Saad M, Toelg R, Khattab AA, Kassner G (2009) Determination of haemodynamic significance of intermediate coronary lesions using three-dimensional coronary reconstruction. *EuroIntervention* 5: 573-579.
25. Chung WY, Choi BJ, Lim SH, Matsuo Y (2015) Three dimensional quantitative coronary angiography can detect reliably ischemic coronary lesions based on fractional flow reserve. *J Korean Med Sci* 30: 716-724.
26. Nishi T, Kitahara H, Fujimoto Y, Nakayama T (2017) Comparison of 3-dimensional and 2-dimensional quantitative coronary angiography and intravascular ultrasound for functional assessment of coronary lesions. *J Cardiol* 69: 280-286.
27. Voros S, Rinehart S, Vazquez-Figueroa JG, Kalynych A, Kampaliotis D, et al. (2014) Prospective, head-to-head comparison of quantitative coronary angiography, quantitative computed tomography angiography, and intravascular ultrasound for the prediction of hemodynamic significance in intermediate and severe lesions, using fractional flow reserve as reference standard (from the ATLANTA I and II Study). *Am J Cardiol* 113: 23-29.

Author Affiliations

Top

Department of Cardiology, Menoufia University, Menoufia Governorate, Egypt

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission