



Research Article

Sensitivity Analysis on Regime-Switching Models by Wiener-Malliavin Calculus

Yue Liu^{1,2*}, Yongsheng Hang², Hongxing Yao² and Yan Chen²

Abstract

In this paper we present the sensitivity analysis on regime switching models applying the techniques of Malliavin calculus. As is well known, risk management in portfolio pricing and hedging is often achieved by estimating the Greeks, which are price sensitivities relative to variations in the model parameters. By developing this method for sensitivity analysis, we have multiple versions of Greeks expression, optimization by minimizing the variance of weight is available among those alternatives. Although the classical Malliavin calculus approach requires the differentiability of the payoff function, we extend the results for models with non-differentiable payoff function.

Keywords

Wiener-Malliavin calculus; Sensitivity analysis; Regime-switching models

Introduction

Emerging interests focus on sensitivity analysis with its wide application in risk management, especially for hedging strategy. In particular, Greeks are the sensitivities with respect to parameters in the generalized Black-Scholes' models. Starting from Fournié et al. [1], Malliavin calculus is applied for computation of Greeks such like Delta (Δ), Rho (ρ), respectively representing the sensitivities of option value to spot price, risk-free rate. Via this approaches, fast algorithms for Greeks computations are designed. Following the method initiated on the Wiener space in Fournié and El-Khatib et al. [1,2] calculates the Greeks in a model of jump process driven by Poisson jump times. By Debelley et al. [3], sensitivities in a jump diffusion model are computed by using the Malliavin calculus. By Davis et al. [4], Malliavin calculus is applied for Levy processes and integration by parts formula is developed for Greeks calculation, where the computational efficiency is assessed by comparison through Monte Carlo simulation. Using the Malliavin calculus on Poisson space, [5] computes the sensitivities for European and Asian options simulated by jump type diffusion. Applying the Malliavin calculus in time inhomogeneous jump-diffusion models, Denis et al. [6] obtains an expression for the sensitivity Theta of an option price as the expectation of the option payoff multiplied by a stochastic weight. Also, Bayazit et al. [7] presents sensitivities for options when the underlying dynamic follows an exponential Levy process. For other

recent works on computation of Greeks in asset price models, c.f. also Denis and Liu [8,9].

On the other hand, Regime-switching models have been introduced by Hamilton et al. [10] in discrete time and are among the most popular and effective risky asset models. The regime switching property is reflected in the changes of states of a Markov chain, which stands for the influence of external market factors. Intensive researches about asset pricing and trading are modelled with Markovian regime-switching, such as Yao et al. [11], Kim et al. [12], Liu and privault [13,14]. In this paper we compute the Greeks by Malliavin calculus in a generalized Black-Scholes model with regime switching that reflects the underlying changes in the state of the economy and extend the results for models with non-differentiable payoff function.

Applying the Wiener-Malliavin calculus, we compute the sensitivities in the framework of the regime switching model (2) below. For any payoff function $\phi \in C_b^1(\mathbb{R})$ with bounded derivative, value function V given by (3) below, and any $g: \{1, \dots, m\} \rightarrow (0, \infty)$, we show by Proposition 3.1 that

$$\Delta = \frac{\partial V(x_0, \alpha_0)}{\partial x_0} = \frac{e^{-rT}}{x_0} E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \quad (1)$$

Where $\langle \sigma, g \rangle := \int_0^T \sigma(\alpha_s) g(\alpha_s) ds$ note that optimal choice of g for efficient Monte Carlo simulation is achievable, by Section 4, the choice of g such that $g(i) = \sigma(i)$ for all $i \in \{1, \dots, m\}$ minimizes the variance of the weight of Δ . Other Greeks like Γ , vega, Λ are similarly computed in Subsection 3.2-3.4. Moreover we note that,

- i) Given $\phi \in C_b^1(\mathbb{R})$ differentiating the payoff function ϕ inside expectation tends to be easier than our approach applying Malliavin calculus.
- ii) In the right-hand of the expression (1), no derivative of the payoff function ϕ appears any more.

Based on these observations, it is tempted and necessary to relax the restriction of ϕ . Therefore, by Section 5, we extend the results to a class of non-differentiable payoff functions.

Formulation

Consider a standard Brownian motion $(W_t)_{t \in \mathbb{R}_+}$ and a Markov chain $(\alpha_t)_{t \in \mathbb{R}_+}$, assume that they are independent and the Markov chain α_t has a state space $M = \{1, \dots, m\}$. Consider the stochastic process $(X_t)_{t \in \mathbb{R}_+}$ given by the following SDE:

$$dX_t = X_t [\mu(\alpha_t) dt + \sigma(\alpha_t) dW_t], \quad 0 \leq t \leq T, \quad (2)$$

with $X_0 = x_0 > 0$, $\alpha_0 \in M$, and $\mu: M \rightarrow \mathbb{R}$ and $\sigma: M \rightarrow (0, \infty)$, are deterministic functions. Denote the filtration generated by $(B_t)_{t \in \mathbb{R}_+}$ and $(\alpha_t)_{t \in \mathbb{R}_+}$ as $(F_t)_{t \in \mathbb{R}_+}$. Given a payoff function ϕ on \mathbb{R} consider t

he value function $V(x_0, \alpha_0)$ defined as follows:

$$V(x_0, \alpha_0) = e^{-rT} E[\phi(X_T) \mid X_0 = x_0, \alpha_0] \quad (3)$$

Where $(X_t)_{t \in [0, T]}$ follows SDE (2), $r > 0$ denotes the risk-free rate. Then we proceed to compute the sensitivities of $V(x_0, \alpha_0)$ to the changes of coefficients in this model, such like $X_0 = x_0$, μ , σ and even Q-matrix.

*Corresponding author: Yue Liu, School of Physical and Mathematical Sciences, Nanyang Technological University, Finance and Economics Jiangsu University, China, Tel: +86 15905288730; E-mail: h.zheng@imperial.ac.uk

Received: July 27, 2017 Accepted: March 01, 2018 Published: March 08, 2018

To begin with, we recall some Malliavin operators. For any random variable $F \in L^2(\Omega, P)$ of the form:

$$F = f\left(\int_0^T h_1(\alpha_s) dW_s, \dots, \int_0^T h_n(\alpha_s) dW_s, \int_0^T h_0(\alpha_s) ds\right)$$

For $n \in \mathbb{N}$, $f \in C^1(\mathbb{R}^{n+1})$, and deterministic functions $h_i: M \rightarrow \mathbb{R}$, $i \in \{0, 1, \dots, n\}$. Given $F \in L^2(\Omega, P)$ in the form of (3), the gradient of F is defined by

$$D_t F = \sum_{k=1}^n \partial_k f\left(\int_0^T h_1(\alpha_s) dW_s, \dots, \int_0^T h_n(\alpha_s) dW_s, \int_0^T h_0(\alpha_s) ds\right) h_k(\alpha_t), \quad (4)$$

For $t \in [0, T]$. For any deterministic function $g: M \rightarrow \mathbb{R}$ we define

$$W(g) := \int_0^T g(\alpha_t) dW_t \quad (5)$$

$$D_g F := \int_0^T D_t F g(\alpha_t) dt \quad (6)$$

for any random variable $F \in L^2(\Omega, P)$ in form of (3).

Computation of the Greeks based on the Partial Malliavin Calculus

In this section, we proceed the computation of the sensitivities of $V(x_0, \alpha_0)$ to the changes of coefficients in this model, such like X_0, μ, \dots . First in this section, we assume that $\phi \in C_b^1(\mathbb{R})$ and has bounded derivative, extension to non-differentiable payoff function will be shown in Section 5.

Variations in the initial price

The sensitivity of value function (2) to the initial price X_0 is given by the following proposition:

Proposition 3.1 For any payoff function $\phi \in C_b^1(\mathbb{R})$ with bounded derivative $(X_t)_{t \in [0, T]}$, defined in (2) and any $g: M \rightarrow (0, \infty)$, we have

$$\frac{\partial V(x_0, \alpha_0)}{\partial x_0} = \frac{e^{-rt}}{x_0} E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \quad (7)$$

Where $\langle \sigma, g \rangle := \int_0^T \sigma(\alpha_s) g(\alpha_s) ds$

Proof

First we check that the right-hand side of (7) is well defined by the following points,

- i) $\langle \sigma, g \rangle \geq \varepsilon > 0$ a.s. for some $\varepsilon > 0$, since $\sigma(i) > 0$ and $g(i) > 0$ for all $i \in M$
- ii) We show that $\phi(X_T) W(g) / \langle \sigma, g \rangle$ is integrable. Since

$$\begin{aligned} E \left[\frac{W(g)^2}{\langle \sigma, g \rangle^2} \right] &= E \left[\frac{\int_0^T g^2(\alpha_s) ds}{\left(\int_0^T \sigma(\alpha_s) g(\alpha_s) ds \right)^2} \right] \\ &\leq \frac{\max\{g^2(i); i \in M\}}{T \min\{\sigma^2(i) g^2(i); i \in M\}} < \infty \end{aligned}$$

and $E\phi(X_T)^2 < \infty$ for ϕ is a bounded function, we claim the integrability of $\phi(X_T) W(g) / \langle \sigma, g \rangle$.

For the left-hand side, we need to show that $V(x_0, \alpha_0)$ is differentiable with respect to x_0 . Since ϕ has bounded derivative, there exists a $K_0 > 0$ that $|\partial \phi(x) / \partial x| < K_0$ for any $x \in \mathbb{R}$. We have the solution to the SDE (2)

$$X_t = x_0 \exp\left(\int_0^t \mu(\alpha_u) - \frac{1}{2} \sigma^2(\alpha_u) du + \int_0^t \sigma(\alpha_u) dW_u\right), t \in [0, T] \quad (8)$$

it follows that

$$\frac{\partial \phi(X_T)}{\partial x_0} = \phi'(X_T) \frac{X_T}{x_0},$$

which is uniformly bounded by an integrable random variable $X_T K_0 / x_0$.

Therefore by $V(x_0, \alpha_0)$ is differentiable with respect to $X_0 = x_0$ and

$$\begin{aligned} \frac{\partial V(x_0, \alpha_0)}{\partial x_0} &= e^{-rt} \frac{\partial}{\partial x_0} E[\phi(X_T) \mid X_0 = x_0, \alpha_0] \\ &= e^{-rt} E \left[\frac{\partial \phi(X_T)}{\partial x_0} \mid X_0 = x_0, \alpha_0 \right] \end{aligned} \quad (9)$$

Then we continue to prove (7). It follows from (8) that $X_T \in L^2(\Omega, P)$ in form of (2), and we have

$$D_g X_T = X_T \langle \sigma, g \rangle > 0 \quad (10)$$

Since the payoff function $\phi \in C_b^1(\mathbb{R})$, $\phi(X_T) \in L^2(\Omega, P)$ and is in form of (2), we have the following chain rule:

$$D_g \phi(X_T) = \phi'(X_T) D_g X_T \text{ for any } g: M \rightarrow (0, \infty). \quad (11)$$

Hence by (8-10) we see that

$$\begin{aligned} \frac{\partial V(x_0, \alpha_0)}{\partial x_0} &= e^{-rt} E \left[\phi'(X_T) \frac{\partial X_T}{\partial x_0} \mid X_0 = x_0, \alpha_0 \right] \\ &= e^{-rt} E \left[\frac{D_g \phi(X_T) X_T}{D_g X_T x_0} \mid X_0 = x_0, \alpha_0 \right] \\ &= \frac{e^{-rt}}{x_0} E \left[\frac{D_g \phi(X_T)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \end{aligned} \quad (12)$$

Denote by F_t a filtration generated by $\{\alpha_t; t \in [0, T]\}$, it follows from the integration by parts formula by Lemma 1.2.1 in Nualart et al. [15] and the independence between $(W_t)_{t \in [0, T]}$ and $(\alpha_t)_{t \in [0, T]}$ that,

$$\begin{aligned} &E \left[\frac{D_g \phi(X_T)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \\ &= E \left[E \left[\frac{D_g \phi(X_T)}{\langle \sigma, g \rangle} \mid X_0 = x_0, F_t \right] \mid X_0 = x_0, \alpha_0 \right] \\ &= E \left[E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, F_t \right] \mid X_0 = x_0, \alpha_0 \right] \\ &= E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \end{aligned}$$

for any $g: M \rightarrow (0, \infty)$. By (12,13), we obtain,

$$\Delta = \frac{e^{-rt}}{x_0} E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right]$$

Variations in the initial price for second order

As in most context, the second derivative of the value function with respect to the initial price is denoted as Δ . Similarly, we have,

Proposition 3.2 For any payoff function $\phi \in C_b^2(\mathbb{R})$ with bounded derivative, $g: M \rightarrow (0, \infty)$, and $(X_t)_{t \in [0, T]}$ defined in (2), we have

Where $\langle \sigma, g \rangle := \int_0^T \sigma(\alpha_s) g(\alpha_s) ds$ and $\|g\| := \int_0^T g^2(\alpha_s) ds$

Proof.

$$\begin{aligned} \frac{\partial^2 V(x_0, \alpha_0)}{\partial x_0^2} &= \frac{\partial}{\partial x_0} \left(\frac{e^{-rT}}{x_0} E \left[\frac{\phi(X_T)W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \right) \\ &= -\frac{e^{-rT}}{x_0^2} E \left[\frac{\phi(X_T)W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \\ &+ \frac{e^{-rT}}{x_0} \frac{\partial}{\partial x_0} E \left[\frac{\phi(X_T)W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \\ &= -\frac{e^{-rT}}{x_0^2} E \left[\frac{\phi(X_T)W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \\ &+ \frac{e^{-rT}}{x_0} E \left[\frac{W(g)}{\langle \sigma, g \rangle} \phi'(X_T) \frac{X_T}{x_0} \mid X_0 = x_0, \alpha_0 \right] \tag{13} \\ &= -\frac{e^{-rT}}{x_0^2} E \left[\frac{\phi(X_T)W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \\ &+ \frac{e^{-rT}}{x_0} E \left[\frac{W(g)}{\langle \sigma, g \rangle} D_g \phi(X_T) \frac{X_T}{x_0} \mid X_0 = x_0, \alpha_0 \right] \\ &= -\frac{e^{-rT}}{x_0^2} E \left[\frac{\phi(X_T)W(g)}{\langle \sigma, g \rangle} \mid X_0 = x_0, \alpha_0 \right] \\ &+ \frac{e^{-rT}}{x_0^2} E \left[\frac{D_g \phi(X_T)W(g)}{\langle \sigma, g \rangle^2} \mid X_0 = x_0, \alpha_0 \right] \end{aligned}$$

Where $\langle \sigma, g \rangle > 0$ a.s. By the chain rule of derivative and the independence between $(W_t)_{t \in [0, T]}$ and $(\alpha_t)_{t \in [0, T]}$ we have

$$\begin{aligned} &E \left[D_g \phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle^2} \mid X_0 = x_0, \alpha_0 \right] \\ &= E \left[D_g \left(\frac{\phi(X_T)W(g)}{\langle \sigma, g \rangle^2} \right) \mid X_0 = x_0, \alpha_0 \right] \\ &- E \left[\phi(X_T) D_g \left(\frac{W(g)}{\langle \sigma, g \rangle^2} \right) \mid X_0 = x_0, \alpha_0 \right] \tag{14} \\ &= E \left[\frac{\phi(X_T)W(g)^2}{\langle \sigma, g \rangle^2} \mid X_0 = x_0, \alpha_0 \right] - E \left[\frac{\phi(X_T) \int_0^T g^2(\alpha_s) ds}{\langle \sigma, g \rangle^2} \mid X_0 = x_0, \alpha_0 \right] \end{aligned}$$

where in the last line we applied the integration by parts formula. Substituting (14) into (13), we see that

$$\frac{\partial^2 V(x_0, \alpha_0)}{\partial x_0^2} = \frac{e^{-rT}}{x_0^2} E \left[\phi(X_T) \frac{W(g)^2 - \|g\| - W(g)\langle \sigma, g \rangle}{\langle \sigma, g \rangle^2} \right]$$

Where $\langle \sigma, g \rangle = \int_0^T \sigma(\alpha_s)g(\alpha_s)ds > 0$ and $\|g\| = \int_0^T g^2(\alpha_s)ds > 0$

Variations in the diffusion coefficient

The variations in the diffusion coefficient is denoted as Vega, precisely,

$$vega = \frac{\partial V(x_0, \alpha_0)}{\partial \sigma}$$

where σ is a constant parameter. In our model, $\sigma(\cdot)$ is a function on M , so there is no direct way of derivative. Instead, we introduce the perturbed process:

$$dX_t^{\sigma, \varepsilon} = X_t^{\sigma, \varepsilon} [\mu(\alpha_t) dt + (\sigma(\alpha_t) + \varepsilon) dW_t] \tag{15}$$

Where $\varepsilon \in \mathbb{R}$, $X_0^{\sigma, \varepsilon} := X_0 = x_0$ Accordingly, we define

$$V^{\sigma, \varepsilon}(x_0, \alpha_0) := E \left[e^{-rT} \phi(X_T^{\sigma, \varepsilon}) \mid X_0 = x_0, \alpha_0 \right] \tag{16}$$

Then the variation in the diffusion coefficient is given by

$$vega = \frac{\partial V^{\sigma, \varepsilon}(x_0, \alpha_0)}{\partial \varepsilon} \Big|_{\varepsilon=0} \tag{17}$$

Note that the solution to (15) is

$$X_t^{\sigma, \varepsilon} = x_0 \exp \left(\int_0^t \mu(\alpha_u) - \frac{1}{2} (\sigma(\alpha_u) + \varepsilon)^2 du + \int_0^t (\sigma(\alpha_u) + \varepsilon) dW_u \right)$$

For $t \in [0, T]$, we compute Vega by the following proposition.

Proposition 3.3 For any payoff function $\phi \in C_b^1(\mathbb{R})$ with bounded derivative, $g: M \rightarrow (0, \infty)$, and $(X_t)_{t \in [0, T]}$ defined in (2), we have

$$\begin{aligned} &vega = \frac{\partial V^{\sigma, \varepsilon}(x_0, \alpha_0)}{\partial \varepsilon} \Big|_{\varepsilon=0} \\ &= e^{-rT} E \left[\phi(X_T) \left(\frac{W_T W(g) - \int_0^T g(\alpha_s) ds - W(g) \int_0^T \sigma(\alpha_s) ds}{\langle \sigma, g \rangle} \right) \mid X_0 = x_0, \alpha_0 \right] \tag{18} \end{aligned}$$

Where $\langle \sigma, g \rangle := \int_0^T \sigma(\alpha_s)g(\alpha_s)ds$

Proof. By the integration by parts formula and chain rule we have,

$$\begin{aligned} &\frac{\partial V^{\sigma, \varepsilon}(x_0, \alpha_0)}{\partial \varepsilon} = \frac{\partial}{\partial \varepsilon} E \left[e^{-rT} \phi(X_T^{\sigma, \varepsilon}) \mid X_0 = x_0, \alpha_0 \right] \\ &= e^{-rT} E \left[\phi'(X_T^{\sigma, \varepsilon}) \frac{\partial X_T^{\sigma, \varepsilon}}{\partial \varepsilon} \mid X_0 = x_0, \alpha_0 \right] \\ &= e^{-rT} E \left[\frac{D_g \phi(X_T^{\sigma, \varepsilon})}{D_g X_T^{\sigma, \varepsilon}} X_T^{\sigma, \varepsilon} \left(W_T - \int_0^T \sigma(\alpha_u) du - \varepsilon T \right) \mid X_0 = x_0, \alpha_0 \right] \\ &= e^{-rT} E \left[\frac{D_g \phi(X_T^{\sigma, \varepsilon})}{\langle \sigma + \varepsilon, h \rangle} \left(W_T - \int_0^T \sigma(\alpha_u) du - \varepsilon T \right) \mid X_0 = x_0, \alpha_0 \right] \\ &= e^{-rT} E \left[\frac{1}{\langle \sigma + \varepsilon, h \rangle} D_g \left(\phi(X_T^{\sigma, \varepsilon}) \left(W_T - \int_0^T \sigma(\alpha_u) du - \varepsilon T \right) \right) \mid X_0 = x_0, \alpha_0 \right] \tag{19} \\ &= e^{-rT} E \left[\frac{\phi(X_T^{\sigma, \varepsilon})}{\langle \sigma + \varepsilon, h \rangle} D_g \left(W_T - \int_0^T \sigma(\alpha_u) du - \varepsilon T \right) \mid X_0 = x_0, \alpha_0 \right] \\ &= e^{-rT} E \left[\frac{\phi(X_T^{\sigma, \varepsilon}) W(g)}{\langle \sigma + \varepsilon, h \rangle} \left(W_T - \int_0^T \sigma(\alpha_u) du - \varepsilon T \right) \mid X_0 = x_0, \alpha_0 \right] \\ &- e^{-rT} E \left[\frac{\phi(X_T^{\sigma, \varepsilon}) \int_0^T g(\alpha_s) ds}{\langle \sigma + \varepsilon, h \rangle} \mid X_0 = x_0, \alpha_0 \right] \end{aligned}$$

for any deterministic function $g: M \rightarrow \mathbb{R}$, $\varepsilon \in \mathbb{R}$. Note by (18) that

$$\lim_{\varepsilon \rightarrow 0} X_t^{\sigma, \varepsilon} = X_t, t \in [0, T] \text{ a.e.,}$$

the expression Vega in (18) is obtained by passing ε to 0 in (19).

Variations in the drift coefficient

Similarly, we introduce the perturbed process for drift coefficient:

$$dX_t^{\mu, \varepsilon} = X_t^{\mu, \varepsilon} \left[(\mu(\alpha_t) + \varepsilon) dt + \sigma(\alpha_t) dW_t \right] \tag{20}$$

and the value function $V^{\mu, \varepsilon}(x_0, \alpha_0)$ driven by $X_t^{\mu, \varepsilon}$

$$V^{\mu, \varepsilon}(x_0, \alpha_0) := e^{-rT} E \left[\phi(X_T^{\mu, \varepsilon}) \mid X_0 = x_0, \alpha_0 \right] \tag{21}$$

Accordingly, we have the solution to (20) as follows,

$$\begin{aligned} &X_T^{\mu, \varepsilon} = x_0 \exp \left(\int_0^T \left(\mu(\alpha_u) + \varepsilon - \frac{1}{2} \sigma^2(\alpha_u) \right) du + \int_0^T \sigma(\alpha_u) dW_u \right) \\ &= X_{t^{\varepsilon, t}}^{\mu, \varepsilon}, t \in [0, T] \end{aligned} \tag{22}$$

Through a similar computation as that of Vega, we have the derivative of $V^{\mu, \varepsilon}(x_0, \alpha_0)$ with respect to ε :

$$\begin{aligned} & \frac{\partial V^{\mu,\varepsilon}(x_0, \alpha_0)}{\partial \varepsilon} \\ &= e^{-rT} E \left[\frac{Dg\phi(X_T^{\mu,\varepsilon}) \partial X_T^{\mu,\varepsilon}}{DgX_T^{\mu,\varepsilon} \partial \mu} \Big| X_0 = x_0, \alpha_0 \right] \\ &= Te^{-rT} E \left[\frac{Dg\phi(X_T^{\mu,\varepsilon})}{\langle \sigma, g \rangle} \Big| X_0 = x_0, \alpha_0 \right] \\ &= Te^{-rT} E \left[\frac{\phi(X_T^{\mu,\varepsilon}) W(g)}{\langle \sigma, g \rangle} \Big| X_0 = x_0, \alpha_0 \right] \end{aligned}$$

for any deterministic function $g : M \rightarrow R$ and $\varepsilon \in R$. Therefore, we have the following Proposition 3.4 for Lambda.

Proposition 3.4 For any payoff function $\phi \in C_b^1(R)$ with bounded derivative, $g : M \rightarrow (0, \infty)$, and $(X_t)_{t \in [0, T]}$ defined in (2), we have

$$\Lambda = Te^{-rT} E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \Big| X_0 = x_0, \alpha_0 \right]$$

Where $\langle \sigma, g \rangle := \int_0^T \sigma(\alpha_s) g(\alpha_s) ds$

Optimization of Convergence

In this section, we aim at figuring out an optimal choice of g for efficient Monte Carlo simulation. Based on the computation of Theta in a jump diffusion model, Proposition 4.1 of Denis et al. [6] minimizes the variance of the weight of Theta. In our model, there exists such a problem too. We can minimize the weight by choosing a optimal $g \in G$. We gain a similar conclusion to that in Denis et al. [8].

Proposition 4.1 The weight of Delta in (7) is $(g) := W(g) / \langle \sigma, g \rangle$, then the infimum on $\{\text{var}[\Psi(g)]; g : M \rightarrow R\}$ is attained when

$$g(i) = c(i) \quad i \in M;$$

with a constant $c > 0$, and is given by

$$\inf_{g \in G} \text{Var}[\Psi(g)] = E \left[\left(\int_0^T \sigma^2(\alpha_t) dt \right)^{-1} \right]$$

Proof. Since $W(g)$ and $\langle \sigma, g \rangle$, are independent, we have $E[\Psi(g)] = 0$ and $\text{Var}[\Psi(g)] = E[\Psi(g)^2]$ (23)

For any $g : M \rightarrow R$, we see that $\int_0^T \sigma(\alpha_t) g(\alpha_t) dt > 0$, with the isometry property of $W(g)$, we have

$$E[\Psi(g)^2] = E \left[\frac{\int_0^T g^2(\alpha_t) dt}{\left(\int_0^T \sigma(\alpha_t) g(\alpha_t) dt \right)^2} \right] \tag{24}$$

By Cauchy-Schwarz inequality,

$$\left(\int_0^T \sigma(\alpha_t) g(\alpha_t) dt \right)^2 \leq \int_0^T \sigma^2(\alpha_t) dt \int_0^T g^2(\alpha_t) dt$$

Therefore, by (4.1)-(4.3), we obtain that

$$\text{Var}[\Psi(g)] \geq E \left[\left(\int_0^T \sigma^2(\alpha_t) dt \right)^{-1} \right]$$

In the case of Denis et al. [8], it is proved that the best function chosen in the weight is the constant number. However, in our case, $\{g(i) = c\sigma(i); i \in M\}$ obtains a faster convergence than $\{g(i) = 1; i \in M\}$ does, as shown in the following (Figure 1).

Extension to Non-Differentiable Payoff Function

In this section, by a similar arguments as Kawai et al. [16], we show by Proposition 5.1 that the conclusion in Proposition 3.1 below is able to be extended for non-differentiable payoff function in the class $\Lambda(R)$ defined as in Kawai et al. [16],

$$\Lambda(R) := \left\{ f : \mathbb{R} \rightarrow \mathbb{R} \mid f = \sum_{i=1}^n f_i 1_{A_i}, n \geq 1, f_i \in C_L(\mathbb{R}; \mathbb{R}) \right.$$

A_i are intervals of \mathbb{R} },

Where

$$C_L(R) := \{f \in C(R; R) \mid |f(x) - f(y)| \leq K|x - y| \text{ for some } K \geq 0\}$$

Proposition 3.4 For any payoff function $\phi \in C_b^1(R)$ with bounded derivative, $g : M \rightarrow (0, \infty)$, and $(X_t)_{t \in [0, T]}$ defined in (2), we have

$$\Delta = \frac{e^{-rT}}{x_0} E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \Big| X_0 = x_0, \alpha_0 \right] \tag{25}$$




Figure 1: Computation of Delta with respect to the number of samples.

Where $\langle \sigma, g \rangle = \int_0^T \sigma(\alpha_s)g(\alpha_s)ds > 0$

Proof. For any $\varphi \in \Lambda(R)$, there exists $N \geq 1$ and a sequence $\{k_i \in R, i=1, \dots, N\}$ and a list of disjoint sets (A_1, \dots, A_N) such that

$$\phi(x) = \sum_{i=1}^N f_i(x)1_{A_i}(x), \quad x \in R \tag{26}$$

where $f_i(x) \in C_L(R; R)$ with

$$|f(x) - f(y)| \leq k_i |x - y|, \quad x, y \in A_i, \quad i=1, \dots, N, \tag{27}$$

And $A_i = (a_{i-1}, a_i], \quad i=1, \dots, N,$

With $a_0 = -\infty$ and $a_N = \infty$

(i) First, we prove that (25) holds for $\varphi \in \Lambda(R)C^1(R)$ Recalling the definition (2.1) of $(X_t)_{t \in [0, T]}$, we let $X_0 = x_0 > 0$, and for any $\varepsilon \in R$ we define $(X_t^\varepsilon)_{t \in [0, T]}$ by

$$X_t^\varepsilon := \frac{x_0 + \varepsilon}{x_0} X_t, \quad t \in [0, T] \tag{28}$$

Without letting φ bounded, we can also show that $\varphi(X_T)$ is integrable. Since φ is continuous, by (27) we see that

$$\begin{aligned} & |\varphi(X_T) - \varphi(X_0)| \\ & \leq \max_{i \in \{1, \dots, N\}} k_i \max_{i \in \{1, \dots, N\}} |X_T - a_i| + \max_{i \in \{1, \dots, N\}} k_i \max_{i \in \{1, \dots, N\}} |x_0 - a_i| \\ & + \sum_{i, j \in \{1, \dots, N\}} |\phi(a_i) - \phi(a_j)| \\ & \leq \max_{i \in \{1, \dots, N\}} k_i \left(X_T + \max_{i \in \{1, \dots, N\}} |a_i| \right) + \max_{i \in \{1, \dots, N\}} k_i \max_{i \in \{1, \dots, N\}} |x_0 - a_i| \\ & + \sum_{i, j \in \{1, \dots, N\}} |\phi(a_i) - \phi(a_j)| \end{aligned}$$

which is integrable, hence the integrability of $\varphi(X_T)$ is proved. On the other hand, without the bounded derivative of φ , we also show that (10) holds, namely,

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \left[\frac{\phi(X_T^\varepsilon) - \phi(X_T)}{\varepsilon} \middle| X_0 = x_0, \alpha_0 \right] \\ & = E \left[\lim_{\varepsilon \rightarrow 0} \frac{\phi(X_T^\varepsilon) - \phi(X_T)}{\varepsilon} \middle| X_0 = x_0, \alpha_0 \right] \end{aligned} \tag{29}$$

By (27) and definition (28) of X_T^ε we see that

$$\lim_{\varepsilon \rightarrow 0} \left| \frac{\phi(X_T^\varepsilon) - \phi(X_T)}{\varepsilon} \right| \leq \lim_{\varepsilon \rightarrow 0} \left(\max_{1 \leq i \leq N} k_i \right) \left| \frac{X_T^\varepsilon - X_T}{\varepsilon} \right| = \frac{1}{x_0} \max_{1 \leq i \leq N} k_i$$

which is uniformly bounded. Therefore, (29) is proved by Lebesgue's dominated convergence theorem and we have

$$\begin{aligned} & \frac{\partial}{\partial x_0} V(x_0, \beta_0) = \lim_{\varepsilon \rightarrow 0} e^{-rT} E \left[\frac{\phi(X_T^\varepsilon) - \phi(X_T)}{\varepsilon} \middle| X_0 = x_0, \alpha_0 \right] \\ & = e^{-rT} E \left[\lim_{\varepsilon \rightarrow 0} \frac{\phi(X_T^\varepsilon) - \phi(X_T)}{\varepsilon} \middle| X_0 = x_0, \alpha_0 \right] \end{aligned} \tag{30}$$

Relation (26) follows from (31) in the case $\phi \in \Lambda(R) \cap C^1(R; R)$ by repeating the arguments from (10) until the end of Subsection 3.1.

(ii) Finally, we extend from $\phi \in \Lambda(R) \cap C^1(R)$ to the class $\Lambda(R)$. We will structure a sequence $(\phi_n)_{n \in N} \in \Lambda(R) \cap C^1(R)$ approaching to $\phi \in \Lambda(R)$. By e.g. Theorem 7.17 of Rudin et al. [17] or (3:6)-(3:7) in Kawai et al. [16], it suffices to show that for all compact set $K(0, \infty)$ we have

$$\lim_{n \rightarrow \infty} E[\phi_n(X_T) | X_0 = x_0, \alpha_0] = E[\phi(X_T) | X_0 = x_0, \alpha_0] \tag{31}$$

for any $x_0 \in K$ and

$$\limsup_{n \rightarrow \infty, x_0 \in K} \left| \frac{\partial}{\partial x_0} E[\phi_n(X_T) | X_0 = x_0, \alpha_0] - E \left[\phi_n(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \middle| X_0 = x_0, \alpha_0 \right] \right| = 0 \tag{32}$$

Since $\varphi \in \Lambda(R)$ is continuous on every interval $(a_{i-1}, a_i), i=1, \dots, N,$ there exists a point wise increasing sequence $(\varphi_n)_{n \in N} \in \Lambda(R) \cap C^1(R)$ such that

$$\lim_{n \rightarrow \infty} \phi_n(x) = \phi(x), \quad x \in R \setminus \{a_0, a_1, \dots, a_N\}. \tag{33}$$

For any $x \in R \setminus \{a_0, a_1, \dots, a_N\},$ there exist a $N_x > 0$ such that for $n \geq N_x$ we have $|\varphi_n(x)| \leq |\varphi(x)| + 1.$ Since $(\varphi_n)_{n \in N} \in \Lambda(R) \cap C^1(R)$ by (29) we see that, for any $n \geq N_x$

$$\begin{aligned} & |\phi(X_T)| \leq |\phi_n(X_T)| + \max_{i \in \{1, \dots, N\}} k_i (X_T + \max_{i \in \{1, \dots, N\}} |a_i|) \\ & + \max_{i \in \{1, \dots, N\}} k_i \max_{i \in \{1, \dots, N\}} |x - a_i| + \sum_{i, j \in \{1, \dots, N\}} |\phi(a_i) - \phi(a_j)| \\ & \leq |\phi_n(X_T)| + 1 + \max_{i \in \{1, \dots, N\}} k_i (X_T + \max_{i \in \{1, \dots, N\}} |a_i|) \\ & + \max_{i \in \{1, \dots, N\}} k_i \max_{i \in \{1, \dots, N\}} |x - a_i| + \sum_{i, j \in \{1, \dots, N\}} |\phi(a_i) - \phi(a_j)| \end{aligned}$$

which is integrable, hence we can apply the Lebesgue's dominated convergence theorem and by (34) we obtain (32). Regarding (33), it follows from (i) that (25) holds for any $(\varphi_n)_{n \in N} \in \Lambda(R) \cap C^1(R),$ therefore

$$\begin{aligned} & \left| \frac{\partial}{\partial x_0} E[\phi_n(X_T) | X_0 = x_0, \alpha_0] - E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \middle| X_0 = x_0, \alpha_0 \right] \right|^2 \\ & = \left| E \left[\phi_n(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \middle| X_0 = x_0, \alpha_0 \right] - E \left[\phi(X_T) \frac{W(g)}{\langle \sigma, g \rangle} \middle| X_0 = x_0, \alpha_0 \right] \right|^2 \\ & \leq \left(E \left[(\phi(X_T) - \phi_n(X_T)) \frac{W(g)}{\langle \sigma, g \rangle} \middle| X_0 = x_0, \alpha_0 \right] \right)^2 \\ & \leq E[(\phi(X_T) - \phi_n(X_T))^2 | X_0 = x_0, \alpha_0] E \left[\left(\frac{W(g)}{\langle \sigma, g \rangle} \right)^2 \middle| X_0 = x_0, \alpha_0 \right] \end{aligned} \tag{34}$$

where by (8) we have

$$E \left[\left(\frac{W(g)}{\langle \sigma, g \rangle} \right)^2 \middle| X_0 = x_0, \alpha_0 \right] < \infty$$

Define a decreasing sequence $(f_n)_{n \in N}$ of continuous functions:

$$F_n(x) := E[(\varphi(X_T) - \varphi_n(X_T))^2 | X_0 = x, \alpha_0], \quad n \in N, x \in K.$$

By Dini's theorem, $f_n(x)$ pointwise decreases to 0 on $K.$ Therefore, by (34), we complete the proof for (32).

References

1. Fournié E, Lasry JM, Lebuchoux J, Lions PL, Touzi N (1999) Applications of Malliavin calculus to Monte Carlo methods in finance. *Fin Stochastics* 3: 391-412.
2. El-Khatib Y, Privault N (2004) Computations of Greeks in a market with jumps via the Malliavin calculus. *Finance and Stochastics* 8: 161-179.
3. Debelley V, Privault N (2018) Sensitivity analysis of European options in jump discussion models via the Malliavin calculus on the Wiener space Preprint. UK.
4. Davis MH, Johansson MP (2006) Malliavin Monte Carlo greeks for jump diffusions. *Stoch Process Appl* 116: 101-129.
5. Bavouzet MP, Messaoud M (2006) Computation of Greeks using Malliavin's calculus in jump type market models. *Electron J Probab* 11: 276-300.
6. Denis L, Nguyen TM (2016) Malliavin calculus for Markov chains using perturbations of time. *Stochastics* 88: 813-840.
7. Bayazit D, Nolder CA (2013) Sensitivities of options via Malliavin calculus:

- applications to markets of exponential Variance Gamma and Normal Inverse Gaussian processes. Quant Fin 13: 1257-1287.
8. Denis L, Nguyen TM (2016) Sensitivities of options via Malliavin calculus: applications to markets of exponential variance Gamma and normal inverse Gaussian processes. Stochastics 88: 813840.
 9. Liu Y, Privault N (2017) An integration by parts formula in a Markovian regime switching model and application to sensitivity analysis. Stoch Anal Appl 35: 919-940.
 10. Hamilton JD (1989) A new approach to the economic analysis of nonstationary time series and the business cycle. Econometrica 1: 357-384.
 11. Yao DD, Zhang Q, Zhou XY (2006) A regime-switching model for European options. In: Stochastic processes, optimization, and control theory: applications in financial engineering, queueing networks, and manufacturing systems. Springer, Boston, USA.
 12. Kim J, Jeong D, Shin DH (2014) A regime-switching model with the volatility smile for two-asset European options. Automatica 50: 747-755.
 13. Liu Y, Privault N (2017) A recursive algorithm for optimal selling in regime switching models. Methodol Comput Appl 1702: 1-16.
 14. Liu Y, Privault N (2017) Selling at the ultimate maximum in a regime-switching model. Int J Theoretical Appl Finance 20: 1750018.
 15. Nualart D (1995) The Malliavin Calculus and Related Topics. Probability and its Application. Springer, Verlag, USA.
 16. Kawai R, Takeuchi A (2011) Greeks formulas for an asset price model with Gamma processes. Mathematic Fin 21: 723-742.
 17. Rudin W (1964) Principles of mathematical analysis. McGraw-hill, New York, USA.

Author Affiliation

Top

¹School of Physical and Mathematical Sciences, Nanyang Technological University, China

²School of Finance and Economics Jiangsu University, China

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission