



Research Article

A SCITECHNOL JOURNAL

Synergistic Potential of Dillapiole Combined With Pyrethroids against Mosquitoes

Erika Oliveira Gomes^{1*}, Sergio Massayoshi Nunomura², Osvaldo Marinotti³ and Wanderli Pedro Tadei¹

Abstract

The synergistic potential of dillapiole, a phenylpropanoid isolated from *Piper aduncum*, in a combination with pyrethroids was evaluated. Although dillapiole has already been proposed as a synergist for a range of insecticides, it has not been tested for controlling disease vector mosquitoes. *Aedes aegypti* and *Anopheles albittarsis* mosquitoes were exposed to Cypermethrin, α -cypermethrin and dillapiole. Tests were conducted in the bottles were impregnated with pyrethroid or pyrethroid plus dillapiole. Bottles with dillapiole-insecticide combinations were prepared in the same way as the insecticide-only impregnated bottles. The TL50 and TL90 values of dillapiole/pyrethroid combinations were significantly different from those obtained with pyrethroid or dillapiole applied separately. The data indicates potential for dillapiole to be used in public health programs directed to mosquito control. The potential advantages of using dillapiole as a synergist and the consequent reduction in the quantity of applied insecticide include lowered cost of applications, decreased environmental impact and improved insecticide resistance management.

Keywords

Anopheles; *Aedes*; Dillapiole; Pyrethroid; Synergism

Introduction

Vectors of human diseases include anopheline mosquitoes, which are vectors of parasites causing malaria and culicine mosquitoes such as *Aedes aegypti* which transmit the viruses that cause dengue, chikungunya, zika and yellow fever. The high rates of morbidity and mortality associated with these diseases encourage studies to develop and test novel means of controlling vector mosquito populations. Biologically active compounds extracted from plants indicate the possibility of replacing or reducing the use of chemical insecticides [1-8].

Some plant extracts, for example, acting as insecticide synergists, containing compounds that are inhibitors of insect P450 cytochromes responsible for the metabolism of xenobiotics, including insecticides. These include phytochemicals containing methylene dioxy rings such as dillapiole, piperamides, and furanocoumarins [1,9]. Since the 1940s, many natural and synthetic compounds with insecticide-synergistic action have been used commercially [2,10-16].

*Corresponding author: Erika Oliveira Gomes, Laboratory of Malaria and Dengue, National Research Institute of Amazon — INPA, Avenida Andre Aratijo, 2936, Aleixo, Manaus, Amazonas, Brasil, Tel: + 55-92 3643 3364; E-mail: erika.gomes@inpa.gov.br

Received: November 24, 2016 Accepted: December 08, 2016 Published: December 15, 2016

Plants of the genus *Piper* have been investigated for their antibacterial, antifungal and insecticidal properties [17-19]. One compound in particular, Dillapiole, extracted from *Piper aduncum* and its derivatives have been investigated for their insecticide-synergist action [2,12-14,20-25]. In this work, we evaluated synergistic potential of dillapiole, combined with synthetic insecticides (cypermethrin and α -cypermethrin), against *Aedes aegypti* and *Anopheles albittarsis* adult mosquitoes.

Material and Methods

Extraction of essential oil and isolation of dillapiole

Aerial parts (leaves and twigs) of *Piper aduncum* were collected in the Bosque da Ciência, Instituto Nacional de Pesquisas da Amazônia - INPA, Manaus -AM, Brazil. *Piper aduncum* L. 1753, was identified by the INPA Herbarium personnel, according to the exsiccates of the species already deposited in the Institution. Collected plant materials were kept for 5 days at about 23°C and 45% relative humidity. The dried plant material was milled and volatile oil extracted by hydro distillation using a Clevenger-type apparatus [26]. A total of 6.1g of essential oil were obtained from 490 g of starting dry plant, yielding 1.25%. Dillapiole, was isolated from the oil by flash column chromatography over silica gel, eluted with hexane/ethyl acetate (98:2). Dillapiole was identified by high resolution gas chromatography with flame ionization detection (GC-FID) and gas chromatography coupled to a mass spectrometer (GC-MS).

Insecticides: The commercially available insecticides used in this study were: Cypermethrin: Cipermetrina CE 200, registered by Fersol Ltda, composed of (R,S)- α -cyano-(3-phenoxyphenyl)methyl 3-(2,2-dichlorovinyl)-2,2-dimethyl- cyclopropanecarboxylate; and α -Cypermethrin: AlphaGold SC 200, registered by Bayer CropScience Ltda, composed of cis (R) and (S) enantiomeric isomers of α -cyano-3-phenoxybenzyl-2, 2-dimethyl-3-(2, 2-dichlorovinyl)-cyclopropanecarboxylate.

Mosquitoes

***Anopheles albittarsis* Lynch-Arribálzaga, 1878:** The specimens of *A. albittarsis* used in this work refer to the sensu lato species [27,28]. Collected in the town of Cacaú Pirera, municipality of Iranduba, Amazonas, Brazil. Mosquitoes were collected at night with the aid of a flashlight and captured using a suction apparatus (Castro device). The captured specimens were transferred to paraffin waxed paper cups, stored in a polystyrene (Styrofoam) box and transported to the Malaria and Dengue Laboratory, National Institute of Amazonian Research (INPA) in Manaus, where they were identified and allowed to rest and feed on sugar: water (1:1). *Anopheles albittarsis* mosquitoes were kept in the laboratory for at least 24 hours prior to the bioassays.

***Aedes (Stegomyia) aegypti* Linnaeus, 1762:** *Aedes aegypti* were obtained from a permanent colony, maintained at the temperature of 26 ± 2 °C with relative humidity of 80% in the Laboratório de Vetores de Malária e Dengue of the Instituto Nacional de Pesquisas da Amazônia. Eggs were placed in plastic trays with approximately one liter of water and hatched larvae fed a 1: 1 mixture of fish meal and liver powder [29]. Three day old mosquitoes were used in the tests.

Bioassays: CDC bottle bioassays [27] were performed in 250 mL bottles with screw cap (Schott Inc.). Bottles were impregnated with 1 mL of acetone containing pyrethroid and/or dillapiole to disperse the products homogeneously, and dried for 24 h to ensure that the solvent evaporated completely. Negative control bottles were impregnated with only solvent (acetone). Mortality was never observed in the negative control tests. The concentrations of commercial chemical insecticides were calculated according to the manufacturer's specifications and the internal area for the bottles (mean value of 275.5 cm²).

Tests were performed in triplicate; forty-five mosquitoes were used for each sample, with 15 specimens per bottle, the survival rate and mortality were assessed every 15 min for 90 min. First, pyrethroids and dillapiole were evaluated individually. Cypermethrin was tested at 0.0015, 0.003, 0.004 and 0.006 µg/cm², whereas α-cypermethrin at 0.0001, 0.0006, 0.001 and 0.002 µg/cm². Dillapiole was tested at 0.007, 0.07 and 0.73 µg/cm² for *Aedes aegypti* and 0.004, 0.007, 0.02 and 0.04 µg/cm² for *A. albiparvus*. Test of synergism was conducted combining the commercial insecticides with dillapiole. The registered mortality of mosquitoes *Aedes* and *Anopheles* was subjected to analysis with non-linear model with the following equation:

$$y = 1 / (1 + \exp(a * (\log(x) + b)))$$

Where, y is the proportion of dead mosquitoes (deaths/45) and x is the time in minutes. Statistical analyses were performed with the software R 3.0.2. TL50 is defined as the time that causes 50 % of mortality in the mosquito population and accordingly, TL90 is the time necessary to cause 90 % of mortality.

Results

Initial tests with *A. aegypti* and cypermethrin, in quantities ranging between 0.0015 and 0.006 µg/cm², showed mortality varying from 40% to 100% in 45 min. Dillapiol at 0.007 to 0.730 µg / cm², resulted in *A. aegypti* mortality between 24% and 100% in 45 min. Based on these results, synergism tests were performed with cypermethrin at 0.0015 mg/cm² and dillapiole at 0.07 mg/cm².

A. aegypti mortality following exposure to only cypermethrin (0.0015 mg/cm²), only dillapiole (0.07 mg/cm²), or cypermethrin plus dillapiole ((0.0015 mg/cm² and 0.07 mg/cm² respectively) was determined (Figure 1). At 15 and 30 minutes the mortality values were observed in the assays with mixed compounds (Cypermethrin/Dillapiole) was more than double the sum of the mortalities caused by exposure to Cypermethrin and dillapiole separately (Cypermethrin + Dillapiole). The association Cypermethrin/dillapiole caused 100% mortality in 45 min, while living mosquitoes were still present at 90 minutes of exposure in the assays performed with each separate compound. The cypermethrin/dillapiole association modulated the time these compounds act upon mosquitoes, shortening in approximately half the time necessary to kill, as shown in lethal time causing 50 % (TL50) and 90% of mortality (TL90) (Table 1).

The susceptibility of *A. albiparvus* to α-cypermethrin at concentrations ranging from 0.0001 to 0.002 µg/cm² showed mortalities from 13% to 44% at 45 min. Dillapiole in amounts of 0.04 to 0.004 µg/cm² resulted in percent mortality ranging between 36% and 22% at 45 min. For the study of synergism, the amounts of 0.0006 µg/cm² for α-cypermethrin and 0.0400 µg/cm² for dillapiole were selected.

Similar to the observations with *Aedes aegypti*, the association α-cypermethrin/dillapiole increased mortality of *A. albiparvus* at all times between 30 minutes and 90 minutes of exposure (Figure 2). No statistically significant differences were noted at 15 minutes. The association α-cypermethrin/dillapiole caused 100% mortality between 60 and 75 min, while living mosquitoes were still present at 90 minutes of exposure in the assays performed with each separate compound. The association of these two compounds shortened the time necessary to kill mosquitoes, estimated as TL50 and TL90 (Table 1). These results further support our conclusion that dillapiole is useful in targeting mosquitoes when used together with pyrethroids.




Figure 1: Mortality of *Aedes aegypti* following exposure to Cypermethrin, Dillapiole, or a combination of Cypermethrin plus dillapiole. Error bars represent the 95% confidence interval (CI).




Figure 2: Mortality of *Anopheles albiparvus* following exposure to Cypermethrin, Dillapiole, or a combination of Cypermethrin plus dillapiole. Error bars represent the 95% confidence interval (CI).

Table 1: TL50 and TL90 in *Aedes aegypti* L. and *Anopheles albiparvus* exposed to dillapiole plus pyrethroids. Lethal time causing 50 % and 90% of mortality (TL50 and TL90). Cypermethrin/Dillapiole and a-Cypermethrin/Dillapiole indicate values calculated for mosquitoes exposed to both compounds simultaneously. Cypermethrin + Dillapiole and a-Cypermethrin/Dillapiol indicate values calculated for the sum of mortalities observed for exposure to dillapiol and pyrethroid separately. 95% confidence interval (CI) were calculated for each experimental condition.

Biological Model	Compounds	TL50			TL90		
		Estimate	Lower IC 50	Upper IC 50	Estimate	Lower IC 95	Upper IC 95
<i>Aedes aegypti</i>	Cypermethrin + Dillapiole	36,0	32,9	39,0	55,0	48,8	61,3
	Cypermethrin/Dillapiole	15,6	13,6	17,5	25,2	20,7	29,6
<i>Anopheles albiparvus</i>	a-Cypermethrin + Dillapiole	42,3	33,2	51,4	267,1	82,3	451,9
	a-Cypermethrin/Dillapiole	24,6	21,8	27,4	40,7	35,2	46,3

Discussion

Mosquitoes are important vectors of parasitic, viral pathogens and serious nuisance pests to humans and animals. Therefore the management of mosquito populations and control of mosquito-borne disease transmission are of prime importance to public health and welfare. The most widely-used mosquito control strategies at present use various formulations of chemical insecticides to kill larvae and adults [28,29]. However, these tools are not always efficient, applicable, or available. Increased resistance of mosquitoes to chemical insecticides and the cost of these interventions make them prohibitive in many settings [30-33].

Considering these drawbacks, pursuing informed, efficient ways of using chemical insecticides are essential. In the formulation process it is important to choose the best compounds, including synergists in order to achieve safety, environmental and economic benefits. Our results support the conclusion that dillapiole is useful for public health applications targeting mosquitoes when used together with pyrethroids [34-36].

Acknowledgments

We thank CNPq for awarding a fellowship (Process No. 553235/2005-2), CTPetro, PIATAM project (Potenciais Impactos e Riscos Ambientais na Indústria de Petróleo e Gás na Amazônia), FAPEAM (Fundação de Amparo à Pesquisa do Amazonas) and CAPES/Pró-Amazônia for financial support. We are also grateful to Fundação de Vigilância em Saúde do Amazonas – FVS/Secretaria de Estado da Saúde do Amazonas – SUSAM, for helping with the testing and mosquito collections

References

- Regnault-Roger C, Vincent C, Arnason JT (2012) Essential oils in insect control: low-risk products in a high-stakes world. *Annu Rev Entomol* 57: 405-424.
- Lichtenstein EP, Liang TT, Schulz KR, Schnoes HK, Carter GT (1974) Insecticidal and synergistic components isolated from dill plants. *J Agric Food Chem* 22: 658-664.
- Consoli RA, de- Oliveira RL (1994) Major health importance of mosquitoes in Brazil. *Fiocruz, Rio de Janeiro, Brasil*.
- Alavanja MCR, Hoppin JA, Kamel F (2004) Health effects of chronic pesticide Exposure: Cancer and Neurotoxicity. *Annu Rev Public Health* 25:155-97.
- Boulogne I, Petit P, Ozier-Lafontaine H, Desfontaines L, Loranger-Merciris G (2012) Insecticidal and antifungal chemicals produced by plants: a review. *Environ Chem Let* 10:325-347
- Govindarajan M Sivakumar R, Rajeswary M, Yogalakshmi K (2013) Chemical composition and larvicidal activity of essential oil from *Ocimum basilicum* (L.) against *Culex tritaeniorhynchus*, *Aedes albopictus* and *Anopheles subpictus* (Diptera: Culicidae). *Exp Parasitol* 134: 7-11.
- Briët OJ, Amerasinghe PH, Vounatsou P(2013) Generalized seasonal autoregressive integrated moving average models for count data with application to malaria time series with low case numbers. *PLoS One* 13:8.
- George DR, Finn RD, Graham KM, Sparagano OA (2014) Present and future potential of plant-derived products to control arthropods of veterinary and medical significance. *Parasit Vectors* 7:28.

- Casida JE (1970) Mixed-function oxidase involvement in the biochemistry of insecticide synergists. *J Agric Food Chem* 18: 753-772.
- Haller HL, Mcgovran ER, Goodhue LD, Sullivan WN (1942)The Synergistic Action of Sesamin with Pyrethrum Insecticides. *J Org Chem* 07: 183-184.
- Metcalf RL (1992) The synergistic activity of various selected chemicals on the insecticides carbofuran and malathion against brine shrimp larvae. Department of Agricultural and Environmental Sciences, University of Newcastleupon-Tyne, UK.
- Mukerjee, SK, Saxena VS, Tomar SS(1979) New methylenedioxyphenyl synergists for pyrethrins. *J Agric. Food Chem* 27: 1209-1211.
- Corbel V, Chandre F, Darriet F, Lardeux F, Hougard JM (2003) Synergism between permethrin and propoxur against *Culex quinquefasciatus* mosquito larvae. *Med Vet Entomol* 17: 158-164.
- Waliwitiya R, Nicholson RA, Kennedy CJ, Lowenberger CA (2012) The synergistic effects of insecticidal essential oils and piperonyl butoxide on biotransformational enzyme activities in *Aedes aegypti* (Diptera: Culicidae). *J. Med. Entomol* 49: 614-623.
- Faulde MK, Nehring O (2012) Synergistic insecticidal and repellent effects of combined pyrethroid and repellent-impregnated bed nets using a novel long-lasting polymer-coating multi-layer technique. *Parasitol Res* 111:755-765.
- Chouaïbou M, Zivanovic GB, Knox TB, Jamet HP, Bonfob B (2014) Synergist bioassays: A simple method for initial metabolic resistance investigation of field *Anopheles gambiae* s.l. populations. *Acta Tropica*. 130: 108-111.
- Pimentel FA, Pereira JBM, Oliveira MN (1998) Zoning and characterization of natural habitats of long pepper (*Piper hispidinervum*) in Acre. *Rio Branco: Embrapa. Research Bulletin*.
- Navickiene HMD, Morandim AA, Marques MOM, Young MCM, Kato MJ (2006) Composition and antifungal activity of essential oils from *Piper aduncum*, *Piper arboreum* e *Piper tuberculatum*. *Quimica Nova* 29: 467-470.
- Duarte MR, Siebenrock MCN (2010) Anatomical Characters of Leaf and Stem *Piper mikanianum* (Kunth) Steud, Piperaceae. *Latin American J Pharm* 29: 45-51.
- Tomar SS, Maheshwari ML. Mukerjee SK (1979) Synthesis and synergistic activity of dillapiole based pyrethrum synergists. *J Agri Food Chem* 27:547-550.
- Belzile AS, Majerus SL, Podeszinski C, Guillet G, Durst T, Arnason JT (2000) Dillapiol Derivatives as Synergists: Structure-Activity Relationship Analysis. *Pesticide Biochem Physiol* 66: 33-40.
- Pohlit AM, Quignard ELJ, Nunomura SM, Tadei WP, Hidalgo AF, et al. (2004) Screening of plants found in the State of Amazonas Brazil for larvicidal activity against *Aedes aegypti* larvae. *Acta Amazonica* 34: 97-105.
- Mohan L, Sharma P, Srivastava CN (2007) Comparative efficacy of *Solanum xanthocarpum* extracts alone and in combination with a synthetic pyrethroid, cypermethrin, against malaria vector, *Anopheles stephensi*. *Southeast Asian J Trop Med Public Health* 38: 256-260.
- Scott IM, Jensen HR, Philogène BJR, Arnason JT (2008) A review of *Piper* spp. (Piperaceae) phytochemistry, insecticidal activity and mode of action. *Phytochem Rev* 7: 65-75.
- Rafael MS, Hereira-Rojas WJ, Roper JJ, Nunomura SM, Tadei WP (2008) Potential control of *Aedes aegypti* (Diptera: Culicidae) with *Piper aduncum* L. (Piperaceae) extracts demonstrated by chromosomal biomarkers and toxic effects on interphase nuclei. *Genet Mol Res* 7: 772-781.
- Maia JGS, Zoghbi MGS, Andrade EHA (2001) Aromatic plants in the Amazon and its essential oils. Belém: Museu Goeldi Paraense Emílio, Brasil.

27. Galvão ALA, Damasceno RG (1942) On a new anopheles of Marajo n sp. *Annais Paulista Medica* 44: 424-427.
28. Conn JE, Wilkerson RC, Segura MN, de Souza RT, Schlichting CD, et al. (2002) Emergence of a new Neotropical malaria vector facilitated by human migration and changes in land use. *American J Trop Med Hyg* 66: 18-22.
29. Mulla MS, Chaudhury MFB (1968) Influence of some environmental factors on the viability and hatching of *Aedes aegypti* (L.) eggs. *Mosquito news* 28: 217-221
30. Brogdon WG, Mac Allister JC (1998) Simplification of adult mosquito bioassays through use of time-mortality determinations in glass bottles. *J Am Mosq Control Assoc* 14: 159-164.
31. Lacey LA (2007) *Bacillus thuringiensis* serovariety israelensis and *Bacillus sphaericus* for mosquito control. *J Am Mosq Control Assoc* 23:133-63
32. Walton WE (2007) Larvivorous fishes including *Gambusia*. *J Am Mosq Control Assoc* 23: 184- 220.
33. Hemingway J, Beaty BJ, Rowland M, Scott TW, Sharp BL (2006) The Innovative Vector Control Consortium: improved control of mosquito-borne diseases. *Trends Parasitol* 22:308-312.
34. Federici G, Shaw BJ, Handy RD (2007) Toxicity of titanium dioxide nanoparticles to rainbow trout (*Oncorhynchus mykiss*): gill injury, oxidative stress, and other physiological effects. *Aquat Toxicol*. 84:415-430.
35. Nauen R (2007) Insecticide resistance in disease vectors of public health importance. *Pest Manag Sci* 63: 628-633.
36. Hemingway J (2014) The role of vector control in stopping the transmission of malaria: threats and opportunities. *Philos Trans R Soc Lond B Biol Sci* 369: 20130431.

Author Affiliations

[Top](#)

¹Laboratory of Malaria and Dengue, National Research Institute of the Amazon – INPA, Brasil

²Coordination of Technology and Innovation, National Research Institute of the Amazon – INPA, Brasil

³MTEKPrime, 132 Woodcest Lane, Aliso Viejo, CA 92656, USA

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission