



# The Concept of Personalized Nutrition in Benin as Example for Developing Countries

Silvia Kolossa and Kurt Gedrich\*

### Abstract

Personalized nutrition (PN) collects and analyses individual data iteratively to generate dietary advice tailored to the characteristics of a person. While PN is mainly discussed in the context of Nutrigenetics and non-communicable diseases in developed countries with excessive food supply, this paper contrasts this scenario to one of unequal or short food supply, as found in developing countries. To overcome under-nutrition and avoid untargeted countrywide distributions of supplements, PN could identify individual deficiencies and target these by recommending specific locally available food items. As PN aims at a balanced diet, it simultaneously targets deficient as well as excessive intakes. This might additionally decelerate the prevalence of double burden of malnutrition in developing countries in the course of nutrition transition.

### Keywords

Personalized nutrition; Developing countries; Nutrition transition; Double burden

**Abbreviations:** NCD: Non-Communicable Diseases; PN: Personalized Nutrition; DBS: Dried Blood Spots.

### Introduction

For the last decades, agricultural production, processing and storage of food items, as well as infrastructure, mechanization and urbanization have been enhanced significantly around the world [1]. Consequently, a growing part of the world population faces a continuous abundant food supply, not only in high-income countries, but also in urbanized areas of middle- and low-income countries. This includes many regions in Africa, where the so-called nutrition transition takes place; a shift in the composition of diets towards increased intakes of saturated fat and sugar, and decreased intakes of fiber [2-5]. Unfortunately, such circumstances not only resulted in a decrease in under-nutrition and related diseases. Worldwide, the prevalence rates of formerly rare conditions such as overweight, elevated blood pressure, hyperglycemia and dyslipidemia have been rising contemporaneously, with currently 1.9 billion adults and 41 million children under 5 years being overweight [6,7]. These parameters may lead to chronic non-communicable diseases (NCD) which have been related to 71% of global deaths. It was estimated

that each year 17.7 million people die of cardiovascular diseases, i.e. 43% of the total NCD deaths, 8.8 million of cancer (22%), 3.9 million of respiratory diseases (10%), and 1.6 million of diabetes mellitus (4%) [8]. 75% of the global deaths occur in low- and middle-income countries [7], and it is discussed that nutrition transition as well as social status influence the incidence of cardiovascular diseases and cancer [9,10]. In the West African Republic of Benin the adults' prevalence of obesity rose from 6% in 2008 to 9.6% in 2016 [11]. The prevalence of diabetes was about 5.1% in 2016 [12], and there are already projects to reduce the incidence of diabetes, as well as to reduce the prevalence of its risk factors [13]. Additionally, Benin established a population-based cancer registry in 2014 to monitor cancer incidences [14]. The global burden of NCDs affects not only the subjective well-being; it also contributes substantially to the rise in health care costs. In 2010 for instance, they amounted to 500 billion US\$ for the treatment of diabetes and to 863 billion US\$ for cardiovascular diseases. By the year 2030, these costs were estimated to make 745 and 1,044 billion US\$, respectively [15].

Nevertheless, 462 million adults are still underweight, 155 million children are stunted, and 45% of deaths among children below the age of 6 years are associated with under-nutrition worldwide [6,16]. In Benin, food insecurity still affected about 11% of the population and 34% have limited or poor food consumption in 2013 [17]. Additionally, in 2013, Benin had the third highest hidden hunger index out of 149 countries after Niger and Kenya [18].

This relatively new phenomenon of a "coexistence of under-nutrition along with overweight, obesity or diet-related NCDs, within individuals, households and populations, and across the life-course" was defined by the World Health Organization as "Double burden of malnutrition" [16]. An obese individual with nutritional anemia serves as example for an individual double burden of disease, while an overweight mother with a stunted child is a typical example of a household double burden. Thus in 2016, the United Nations General Assembly declared the years 2016 to 2025 as the "United Nations Decade of Action on Nutrition". Among others, it called for providing nutrition-related education for everyone, to enhance the worldwide knowledge of a healthy lifestyle and prevent the increase of diet-related diseases and costs [6].

### General Advice for a Healthy Diet

The idea to provide information on nutrition and give dietary advice to prevent nutrition related diseases already started centuries ago. In the 15<sup>th</sup> century, for example, the adventurer Jacques Cartier described "scurvy", a disease due to ascorbic acid deficiency. He discovered that consuming "Anned tree extract" cured and prevented scurvy and developed a general recommendation for ships' crews to consume Anneda tree beverages [19]. In the following centuries, further deficiencies of food residues were discovered to cause specific diseases and over time, dietary recommendations were extended and specified.

Nowadays, dietary guidelines are omnipresent and provide easy-to-understand advice for consumers on how to achieve and maintain a healthy lifestyle. They are distributed by transnational organizations, such as the "10 tips for healthy eating" of the European

\*Corresponding author: Kurt Gedrich, Lehrstuhl für Ernährungsphysiologie, Technische Universität München, Munich, Germany, Tel: +49-8161-71-2483, E-mail: kgedrich@tum.de

Received: March 20, 2019 Accepted: April 03, 2019 Published: April 15, 2019

Food Information Council [20], but also on country-level, such as the “10 Regeln der DGE” published by the German Nutrition Society in Europe [21], or the “Guide alimentaire” by the nutritional council of Benin in West Africa [22]. The guidelines are quite similar; however, country specific guidelines put general recommendations into the local context to increase their populations’ comprehension and compliance. Benin for instance provides a visualization of food groups to be consumed as traditional roundhouse with a thatch roof. The food items within the five groups are largely local products, like yam, cassava or Amiwo. The glass and a water bottle at the entrance symbolize Benin’s hospitality and serves as reminder to drink water throughout the whole day (Figure 1).

The burden of NCDs has been rising continuously, despite the widespread availability of plain and locally adapted dietary advice. This raises the question: why do people not follow the guidelines to improve their own health? The food industry follows the innate human demands, e.g. for sweet and fatty food items [2], and lifestyle magazines sell by various claims on healthy nutrition. Additionally to this cacophony of partly conflicting information, general guidelines on a healthy lifestyle do not necessarily stimulate changes. In a meta-analysis of 29 studies, for example, only a weak positive association was found on the relationship of nutritional knowledge and dietary intake [23]. Increasing efforts to promote generic dietary advice might thus only have minor effects on dietary transition and a subsequent decrease in NCDs.

### Personalized Nutrition

Personalized nutrition (PN) tailors dietary advice to the individual requirements and preferences of a person and is supposed

to improve the perceived relevance of a given advice and consequently leads to an increased motivation for changing the dietary behavior towards a healthier lifestyle compared to generic guidelines [24,25]. PN in form of one-on-one attendance by dietitians or physicians has already been existing for several decades [26]. By means of dietary assessment, medical history, and/or phenotypic markers, as blood metabolite levels, an expert develops a personalized dietary plan with specific goals.

Through the worldwide usage of smartphones and applications, the possibility of home-based blood sampling as well as the development of high-through-put technologies to analyze characteristic biomarkers in blood or urine, PN has the opportunity to achieve a new level. PN does not necessarily require expert-conducted manual assessments and analyses anymore, but can use smartphone applications. Here, the analysis of data and feedback development is performed time- and cost-efficiently by implemented algorithms, and the recommendations are available at anytime and anywhere [27]. It is thus applicable for large parts of the population to act as preventative tool, achieving dietary behavior change, before nutrition related-diseases may become manifest. The opportunity for active participation and recognition in social media may also account for a higher effectiveness of personalized advice [28,29]. Additionally, individual feedback to a person’s nutritional behavior might also overcome the proposed inability to evaluate one’s own diet, as most Europeans believe their diet to be healthy enough [30].

Generally, we determined three consecutive steps to PN: data collection, data evaluation, and feedback generation. Data collection is conducted to determine the current characteristics of an individual.




Figure 1: Country specific nutritional guidelines using local food items for visualization: Guide alimentaire, Benin [22].

During data evaluation, the status quo is compared to optimal reference values of the individual. Deviations are rated for example by magnitude and or by predefined rankings. For feedback generation, characteristics to be changed are described and suggestions for behavior changes are given in form of recommendations concerning nutrients, food items or recipes. These steps are iteratively accomplished, and with each cycle, the individual can further approach his/her individual optimal pattern (Figure 2).

### Data collection

We divided the data collection for PN into two parts; i.e. data based on phenotype and genotype, respectively. The phenotype describes observable characteristics, for example the behavior of a person; which includes individual dietary intake, physical activity, and anatomy including anthropometry, as well as biomedical or physiological properties. The genotype is the genetic constitution of a person. Based on these data, the risk for certain diseases can be estimated and opportunities on how to change which characteristics towards a healthier phenotype can be indicated. The data or sample collection might require particular expertise (e.g. bioelectrical impedance analysis for estimating body composition), but often studies aim at measurement approaches that are independent of any professional assistance.

To estimate dietary intake, pro- and retrospective assessment tools have been developed. Widely accepted assessment methods are for example the retrospective 24-hour recalls and food frequency questionnaires. Modern technology facilitates dietary assessment and offers the opportunity for immediate data analysis. Besides the digitalization of formally handwritten questionnaires, more objective assessment tools like photos [27,31] or micro-cameras [32] might increase accuracy, especially in terms of portion size estimation and minimize costs and time. Such tools, however, need additional development and testing concerning validity and usability [33]. Most of the dietary assessment tools aim eventually at assessing not only the intake of food items, but of nutrients. The daily nutrient intake can be calculated using a food composition database, which lists relevant food items of the questionnaires or photos, and additionally comprise precise values of the nutrient content.

Next to dietary behavior, physical activity plays a crucial role in developing NCDs. It can be estimated by questionnaires, e.g. Baecke's questionnaire [34], but also by wearable devices, such as physical activity trackers [35]. However, in only a few cases, the validity of accelerometers was scientifically accredited [36,37]. Anthropometric data offer valuable clues on the body composition and nutritional status. Measurements of the height, weight and circumferences using simple scales and tapes are sufficient to determine e.g. the Body Mass Index, a marker to classify under- and overweight [38,39], and the waist circumference, indicating fat distribution [40]. More sophisticated tools exist for home-based self-assessment of further physiological parameters, such as blood pressure, pulse rate or blood oxygen saturation [41,42]. Again, their validity was rarely confirmed, e.g. for certain sphygmomanometers [41,43]. Phenotyping also includes the analysis of blood metabolites. Several parameters reflect various aspects of nutritional status, e.g. plasma glucose, cholesterol, triglycerides, vitamins etc. [44]. Next to the classical venipuncture sampling, the usage of capillary blood sampling, so-called dried blood spots (DBS), becomes more important in epidemiological studies, as sampling does not require medically trained staff [45].


Nutritional advice can also be personalized based on findings of Nutrigenetics, which examines the interaction of diet and genome [46-48]. Buccal cells, home-based collected with commercially available kits, serve as samples for determining the genetic constitution of a person. DNA samples will then be sequenced using e.g. micro arrays or Next Generation Sequencing [48].

### Data evaluation

Data evaluation aims at comparing the status quo to the individuals' requirements and preferences. However, it is often not feasible to precisely determine the requirements of an individual; they rather have to be estimated. Thus, the assessed daily nutrient intake is compared to respective dietary recommendations, which are estimations for a nutrient intake covering the requirements of 97.5% of the population. They usually only distinguish between sex and age, as well as breastfeeding and pregnancy [49,50]. The recommendations for nutrients vary from country to country, depending on the scientific evidence and the references taken into account, as well as the updating-processes, there are different recommendations [51-53]. The careful choice of the recommendation database is not only evident for the nutrient intake, but for every marker (anthropometrics, blood values etc.). There exists for example a substantial ethnic variation in the distribution of visceral and subcutaneous fat in the body [54]. A study on blood reference values of laboratories in the U.S. and Africa suggested that each laboratory should determine locally derived reference values for each parameter, due to major differences in blood values due to ethnicity [55].

The evaluation of genetical data is based on the findings of nutrigenetic studies. One example is the protective effect of cruciferous vegetables against lung cancer for null allele carrier of GSTM1 and GSTT1, genes coding for glutathione-S-transferases [56]. Thus, if the genotyping of a person's DNA reveals a null allele for GSTM1, the evaluation of the data would indicate a higher need for cruciferous vegetables.

Depending on the feedback, the deviation between status quo and recommendation of each nutrient or phenotypic value is calculated and weighted. The deviation can be expressed by percent of recommended intake or by classifying into certain groups, such as small, medium or high deviation.



### Feedback generation

Feedback on the data evaluation can be based on foods, recipes or broader lifestyle advice. A human dietary advisor would mentally combine the gathered information and address the most important aspects to be changed. Computerized feedback uses the information of the data evaluation and produces semi-quantitative or quantitative recommendations. Semi-quantitative recommendations mainly aim at reducing or increasing the intake of certain food items or groups. If the intake of a nutrient is higher than the recommended intake, food items with a high content of this nutrient are to be consumed less, and vice versa, e.g. “low carotenoids: eat more carrots; high saturated fat and genetic risk allele: eat less chocolate cream”. Quantitative feedback on certain nutrients includes specific quantities of the food items to be consumed e.g. “eat 100 g carrots, pepper or tomatoes per day”. A proof-of-principle study on PN showed that focusing on three to four specific markers, e.g. nutrients, anthropometric or blood markers seem to have a substantial effect on behavior change towards a healthier nutrition. During the study they used the semi-quantitative approach, and recommended several food items to be consumed more or less for each target nutrient [24]. For a holistic quantitative feedback, recipes and menu plans can be considered. Mathematical tools like optimizations are able to calculate the optimal combination of recipes to fulfill not only three or four, but all nutritional requirements and can include economic factors or food preferences [57,58].

If more information is requested on why and how to change the diet, texts on all possible target markers, including recommendations on nutrients, food items and lifestyle can be generated using decision trees [59,60]. Pictures or other visualizations might serve as tools to communicate recommendations without the need of a high education level or even literacy, as simple photos of food items with checkmarks or crossed out already point out the desired dietary change. This approach does however not include any kind of explanation or further education.

On a broader view, recommendations based on genetic variants might raise ethical concerns. Certain genetic constitutions were not only linked to diet-related diseases, but also to elevated risks for severe non-dietary related diseases. An example is a single nucleotide polymorphism in ApoE. On the one hand, it is associated with increased LDL-cholesterol and triglyceride levels [61], on the other, also with an elevated risk for developing Alzheimer’s disease [62]. Before including genetic-based personalized advice, the associated

gene-diet-interaction needs sound scientific evidence for health effects, iterative evaluation of potential psychological effects and the promotion of autonomous choice of each person [63].


### Hypotheses on the Application of PN


#### Excessive food supply

PN was mainly developed for countries with an excessive food supply, both in amounts and variety, as in Europe or North America. It was introduced as a concept to contribute to the reduction of the pandemic dimension of obesity and thus the prevalence of NCDs and their associated health care costs. As the human is evolutionary tailored to save energy, a paradisiac environment where palatable food is omnipresent, also has to be considered as an obesogenic environment. Combined with a sedentary lifestyle and absence of famines, excessive food supply leads to an excessive food intake and promotes obesity. In case of additionally imbalanced food choices, micronutrient deficiencies might be caused, as well (Figure 3, upper part). Even if one intends to follow a healthy diet, a vast quantity of articles, TV magazines or social media entries might also lead to confusion about the “healthiness” of particular food items. The search engine google.com gives over 70 million entries on “healthy eating” (5<sup>th</sup> Nov. 2018). Additionally, a meta-analysis compared the price difference between healthier and less healthy diet patterns, and concluded that the healthier patterns were more expensive [64]. PN might be able to guide the individuals’ food choices towards a healthy but palatable diet, which fulfills the individual requirements (Figure 3, lower part) and increases the awareness of nutrition-related diseases by estimating the personal risks.

#### Unequal or short food supply

An unequal distribution and/or short supply of food leads to under-nutrition of macro- and/or micronutrients (Figure 4, upper part). The challenge is therefore to adapt the regional food production with available resources to meet the nutritional requirements. Individual, household or community based PN advice might be able to identify specific deficiencies or excesses in macro- and micronutrient intake (Figure 4, middle part). The understanding of these deficiencies and/or excessive intakes as well as personalized recommendations enables individuals and families to react proactively by increasing or decreasing the consumption of locally produced foods. PN may also prevent the prevalence of the double burden of disease, as it generally aims at a balanced diet and is not strictly focusing on nutrient





deficiencies. The recognition of the nutritional value of food items, the knowledge about healthy nutrition, and the benefits caused by a large food variety may thus lead to a shift in the intake or at least the demand of certain food items or groups. In rural areas, PN grants additional opportunities for improving the nutritional situation, when food choices are linked to decisions relating the individual agricultural production system causing changes in food production towards nutrient-sensitive agriculture (Figure 4, lower part), which aims, among other, at diversification and sustainable as well as nutritious food production [65].

Using PN via smartphones in rural areas of developing countries is currently limited by availability of suitable devices as well as mobile internet access. However, statistics show an increase in households internet access at home in developing countries from 9% in 2005 to 43% in 2017, which assumes higher coverage of internet every year [66]. In the meantime, PN approaches can be put into practice via suitably equipped advisors (e.g. nurses or community workers).

### Effectiveness of PN

Several studies compared personalized dietary advice to untargeted generic advice. Besides proof of principle studies with healthy participants, other studies examined PN in the context of NCDs. To address the question, whether online-based PN is advantageous over generic advice to induce a life style change, the European Food4Me Study was designed as online-based randomized control trial in seven European counties with about 1200 participants. The aim was to compare the conventional one-size-fits-all to a PN approach, involving phenotypic and genotypic data [67]. In the study, feedback was provided as semi-quantitative recommendations, advising the participants to increase or decrease the intake of certain nutrients and food items. The Food4Me Study confirmed that PN based on participants' dietary data is more effective than a general conventional advice. However, advice on phenotypic or phenotypic plus genotypic information did not lead to an enhanced effectiveness of PN [24].

Zeevi et al. incorporated phenotypic parameters including gut microbiota in a machine learning algorithm to predict individual postprandial glucose response. In their sub sequential randomized

controlled intervention study, participants with PN based on this algorithm had a significantly lower postprandial response [68]. A study by Wright et al. demonstrated that PN was more effective in increasing the fruit intake in participant with cardiovascular risk factors compared to generalized nutritional advice [69]. Shiao et al. identified significant genetic and nutritional predictors for colorectal cancer. They developed a mathematical model, which can be used for personalized advice in colorectal cancer prevention [70].

PN interventions are mainly conducted and analyzed in developed countries to target obesity and over-nutrition in individuals. In contrast, tailored nutritional advice in developing countries was mainly discussed for certain cohort and sub topics, for example infectious diseases or micronutrient deficiencies. HIV-positive children in Tanzania, for example, have a high rate of under-nutrition, which was mainly due to a low frequency of meals and a poor dietary diversity. It was suggested that PN is helpful to educate households to efficiently use available foods and take care of the special nutritional needs of HIV-positive children [71]. A PN program which adapted to local conditions and particular requirements in South Africa successfully improved the weight status of children less than 5 years, but did not catch-up growth in stunted children within two years [72]. A long-term study within the Middle East and North Africa revealed a nutritional shift towards a "Westernized diet" and it was recommended to tailor nutritional education in order to support healthy diets to decrease the rising burden of chronic diseases [73]. Until now however, and to the best of our knowledge, the concept of PN as a holistic approach to improve behavior change generally towards a balanced diet was theoretically approved, but has not yet been tested in developing countries.

### Limitations

Implementing PN in the setting of scarcity is limited with regards to individual, social and technological aspects. Individual aspects include the ability to read and write, as an autonomous usage of PN e.g. as a smartphone application might need written input, e.g. for dietary assessment. However, only about 33% of Benin's population aged 15 years and older is literate, twice as many men than women

[74]. Even if the users could read and write, more than 50 different local languages are spoken in the country, while the statutory national language French is only spoken by about 35% in Benin [75]. Audio systems, pictures or else would enable also illiterate and non-French speakers to take part. The social and religious dietary habits are as diverse as the different languages, thus PN in Benin should include the different food taboos to ensure compliance and reduce the risk for diseases, which are connected to the restrictions [76,77].

## Conclusions

In the last years, PN gained momentum, in science but also in public. The science on PN provides use-oriented research, involving members from a variety of disciplines, such as life and behavioral scientists, psychologists, informatics and data analysts as well as sensor technology experts, and agro-economic specialists.

For future applications of PN, the development of valid, affordable, and easy-to-use data collection methods, especially for an objective measurement of food intake as well as blood analyses, can help to better display the individual behavior and metabolism to find relevant parameters for behavior change. Genetic research may reveal new interaction of genes and nutrition and produce possible ways to affect the health outcome. However, dietary behavior is driven by various aspects; thus, further studies should implement advice also involving the respective environmental, social and demographic context. The inclusion of factors as market price analyses, soil potential and mechanization can contribute to a personalized recommendation on agricultural level, for example in rural Africa. Additionally, the inclusion of personality and education, i.e. grade of self-confidence or stamina, and of the ability to read and understand explanations will pave the way to better analyses of the current status, and improved modes of communication of personalized advice to the recipient. A larger involvement of general health care may be implemented, including guidance to take medical check-ups or indicate nearest health centers for specific tests.

While mainly discussed in the context of Nutrigenetics and obesity, a broader view on the PN concept reveals enormous possibilities in various environments to educate the public and contribute to a nutritional-balanced environment worldwide. It provides the opportunity to individually take responsibility for the own health in a most comprehensive way.

## Author Contributions

Conceptualization, S.K, K.G.; writing-original draft preparation, S.K.; writing-review and editing, K.G.; supervision, K.G.

## Funding

This research was funded by the Bundesministerium für Wirtschaftliche Zusammenarbeit und Entwicklung, Germany, as part of the Program of Accompanying Research for Agricultural Innovation-PARI.

## Acknowledgments

The development of these hypotheses was part of a larger project, which could not have happened without the contributions of Getachew Abate-Kassa and Lena Schickramm. We also thank Joachim von Braun and Heike Baumüller for their engagement in the overall project.

## Conflicts of Interest

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## Reference

1. Food and Agriculture Organization (2017) Crop Prospects and Food Situation No. 4, December.
2. Popkin BM, Adair LS, Ng SW (2012) Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev* 70: 3-21.
3. Vorster HH, Kruger A, Margetts BM (2011) The nutrition transition in Africa: Can it be steered into a more positive direction? *Nutrients* 3: 429-441.
4. Abrahams Z, McHiza Z, Steyn NP (2011) Diet and mortality rates in Sub-Saharan Africa: Stages in the nutrition transition. *BMC Public Health* 11:801.
5. Popkin BM (1993) Nutritional Patterns and Transitions. *Population and Development Review* 19: 138.
6. World Health Organization (2018) Malnutrition.
7. World Health Organization (2018) Noncommunicable diseases - Fact sheet.
8. World Health Organization (2017) Non communicable diseases.
9. Romieu I, Margetts B, Barquera S, Gomes FdS, Gunter M, et al. (2016) Strengthening the evidence base for nutrition and cancer in low and middle income countries. *J Glob Health* 6: 20306.
10. Williams J, Allen L, Wickramasinghe K, Mikkelsen B, Roberts N, et al. (2018) A systematic review of associations between non-communicable diseases and socioeconomic status within low- and lower-middle-income countries. *J Glob Health* 8: 20409.
11. Index mundi Benin - Obesity - adult prevalence rate - Historical Data Graphs per Year.
12. World Health Organization Country Profiles - Benin.
13. World diabetes foundation Prevention of Diabetes in Benin WDF14-872.
14. Egue M, Gnangnon FHR, Akele-Akpo M-T, Maxwell Parkin D (2019) Cancer incidence in Cotonou (Benin), 2014-2016: First results from the cancer Registry of Cotonou. *Cancer Epi* 59: 46-50.
15. Bloom DE, Cafiero ET, Jané-Llopis E, Abrahams-Gessel S, Bloom LR, et al. (2011) The Global Economic Burden of Noncommunicable Diseases. Geneva: World Economic Forum.
16. World Health Organization (2017) The double burden of malnutrition. Policy brief.
17. World Food Program (2018) Benin Country Brief.
18. Muthayya S, Rah JH, Sugimoto JD, Roos FF, Kraemer K, et al. (2013) The global hidden hunger indices and maps: an advocacy tool for action. *PLoS One* 8: e67860.
19. Martini E (2002) Jacques Cartier witnesses a treatment for scurvy. *Vesalius* 8: 2-6.
20. European Food Information Council (2006) 10 tips to healthy eating.
21. Deutsche Gesellschaft für Ernährung e.V. Guidelines of the DGE.
22. Conseil national de l'alimentation et de la nutrition (CAN) Guide alimentaire du Bénin.
23. Spronk I, Kullen C, Burdon C, O'Connor H (2014) Relationship between nutrition knowledge and dietary intake. *Br J Nutr* 111: 1713-1726.
24. Celis-Morales C, Livingstone KM, Marsaux CFM, Macready AL, Fallaize R, et al. (2017) Effect of personalized nutrition on health-related behaviour change: Evidence from the Food4me European randomized controlled trial. *Int J Epidemiol* 46: 578-588.
25. Joost HG, Gibney MJ, Cashman KD, Görman U, Hesketh JE, et al. (2007) Personalised nutrition: Status and perspectives. *Br J Nutr* 98: 26-31.
26. Nizel AE (1972) Personalized Nutrition Counseling. *ASDC J Dent Child* 39: 353-360.
27. Ngo J, Engelen A, Molag M, Roesle J, Garcia-Segovia P, et al. (2009) A review of the use of information and communication technologies for dietary assessment. *Br J Nutr* 101 Suppl 2: S102-12.
28. Brug J, Campbell M, van Assema P (1999) The application and impact of computer-generated personalized nutrition education: A review of the literature. *Patient Educ Couns* 36: 145-156.

29. Brug J, Oenema A, Campbell M (2003) Past, present, and future of computer-tailored nutrition education. *Am J Clin Nutr* 77: 1028S-1034S.
30. Lappalainen R, Kearney J, Gibney M (1998) A pan EU survey of consumer attitudes to food, nutrition and health. An overview. *Food Quality Prefer* 9: 467-478.
31. Boushey CJ, Delp EJ, Ahmad Z, Wang Y, Roberts SM, et al. (2016) Dietary assessment of domoic acid exposure. What can be learned from traditional methods and new applications for a technology assisted device. *Harmful Algae* 57:51-55.
32. Pettitt C, Liu J, Kwasnicki RM, Yang G-Z, Preston T, et al. (2016) A pilot study to determine whether using a lightweight, wearable micro-camera improves dietary assessment accuracy and offers information on macronutrients and eating rate. *Br J Nutr* 115: 160-167.
33. Forster H, Walsh MC, Gibney MJ, Brennan L, Gibney ER (2016) Personalised nutrition: The role of new dietary assessment methods. *Proc Nutr Soc* 75: 96-105.
34. Baecke JA, Burema J, Frijters JE (1982) A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 36: 936-942.
35. Fitbit Aktivitätstracker.
36. Sijtsma A, Schierbeek H, Goris AHC, Joosten KFM, van Kessel I, et al. (2013) Validation of the TracmorD triaxial accelerometer to assess physical activity in preschool children. *Obesity (Silver Spring, Md.)* 21: 1877-1883.
37. Valenti G, Camps SGJA, Verhoef SPM, Bonomi AG, Westerterp KR (2014) Validating measures of free-living physical activity in overweight and obese subjects using an accelerometer. *Int J Obes (Lond)* (2005) 38: 1011-1014.
38. World Health Organization (2018) Obesity and overweight.
39. World Health Organization (2018) Body mass index-BMI.
40. World Health Organization (2011) Waist circumference and waist-hip ratio. Report of a WHO expert consultation, Geneva, 8-11 December 2008.
41. O'Brien E, Mee F, Atkins N, Thomas M (1996) Evaluation of three devices for self-measurement of blood pressure according to the revised British Hypertension Society Protocol: The Omron HEM-705CP, Philips HP5332, and Nissei DS-175. *Blood Press Monit* 1: 55-61.
42. Medvedev O, Kobelev A, Schookin S, Jatskovsky M, Markarian G, et al. (2007) Smartphone-based Approach for Monitoring Vital Physiological Parameters in Humans. In: Magjarevic R, Nagel JH (eds) *World Congress on Medical Physics and Biomedical Engineering 2006*, Springer Berlin Heidelberg, Berlin, Heidelberg 14: 4020-4022.
43. Mengden T, Chamontin B, Phong CN, Luis PGJ, Chanudet X (2000) User procedure for self-measurement of blood pressure. *First International Consensus Conference on Self Blood Pressure Measurement. Blood Press Monit* 5: 111-129.
44. Herbold N, Edelstein S (2008) *Rapid Reference for Nurses: Nutrition*. Jones & Bartlett Learning.
45. Sakhi AK, Bastani NE, Ellingjord-Dale M, Gundersen TE, Blomhoff R, et al. (2015) Feasibility of self-sampled dried blood spot and saliva samples sent by mail in a population-based study. *BMC Cancer* 15: 265.
46. Müller M, Kersten S (2003) Nutrigenomics: Goals and strategies. *Nat Rev Genet* 4: 315-322.
47. Palou A (2007) From nutrigenomics to personalised nutrition. *Genes Nutr* 2: 5-7.
48. Madenach M, Kolossa S, Gedrich K (2018) Nutrigenetik in Theorie und Praxis. *Aktuel Ernährungsmed* 43: 410-424.
49. IOM IoM Health and Medicine Division (HMD): Health and Medicine Division.
50. Deutsche Gesellschaft für Ernährung e.V. Referenzwerte für die Nährstoffzufuhr.
51. Deutsche Gesellschaft für Ernährung e.V (2012) New reference values for vitamin D. *Ann Nutr Metab* 60: 241-246.
52. Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, et al. (2011) The 2011 report on dietary reference intakes for Calcium and Vitamin D from the Institute of Medicine: What Clinicians Need to Know. *J Clin Endocrinol Meta* 96: 53-58.
53. Finnish Food Safety Authority - Evira (2016) Vitamin D.
54. Rønn PF, Andersen GS, Lauritzen T, Christensen DL, Aadahl M, et al. (2017) Ethnic differences in anthropometric measures and abdominal fat distribution. A cross-sectional pooled study in Inuit, Africans and Europeans. *J Epidemiol Community Health* 71: 536-543.
55. Zeh CE, Odhiambo CO, Mills LA (2012) *Laboratory Reference Intervals in Africa*. IntechOpen.
56. Brennan P, Hsu CC, Moullan N, Szeszenia-Dabrowska N, Lissowska J, et al. (2005) Effect of cruciferous vegetables on lung cancer in patients stratified by genetic status: A mendelian randomisation approach. *The Lancet* 366: 1558-1560.
57. Lv Y (2009) *Multi-objective Nutritional Diet Optimization Based on Quantum Genetic Algorithm*. Tianjin, China.
58. Kolossa S (2018) Effectiveness of personalized nutrition - the German experience. Dissertation. Technische Universität München, Germany.
59. Forster H, Walsh MC, O'Donovan CB, Woolhead C, McGirr C, et al. (2016) A dietary feedback system for the delivery of consistent personalized dietary advice in the web-based multicenter food4me study. *J Med Internet Res* 18: 150.
60. Food4Me (2015) White paper on personalised nutrition-paving a way to better population health.
61. Jofre-Monseny L, Minihane A-M, Rimbach G (2008) Impact of apoE genotype on oxidative stress, inflammation and disease risk. *Mol Nutr Food Res* 52: 131-145.
62. Corder EH, Saunders AM, Strittmatter WJ, Schmechel DE, Gaskell PC, et al. (1993) Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 261: 921-923.
63. Görman U, Mathers JC, Grimaldi KA, Ahlgren J, Nordström K (2013) Do we know enough? A scientific and ethical analysis of the basis for genetic-based personalized nutrition. *Genes Nutr* 8: 373-381.
64. Rao M, Afshin A, Singh G, Mozaffarian D (2013) Do healthier foods and diet patterns cost more than less healthy options? A systematic review and meta-analysis. *BMJ Open* 3: 004277.
65. Food and Agriculture Organization (2014) *Nutrition-sensitive agriculture*.
66. International Telecommunication Union (2017) *Global and regional ICT data*.
67. Celis-Morales C, Livingstone KM, Marsaux CFM, Forster H, O'Donovan CB, et al. (2015) Design and baseline characteristics of the Food4Me study: A web-based randomised controlled trial of personalised nutrition in seven European countries. *Genes Nutr* 10: 450
68. Zeevi D, Korem T, Zmora N, Israeli D, Rothschild D, et al. (2015) Personalized Nutrition by Prediction of Glycemic Responses. *Cell* 163: 1079-1094.
69. Wright JL, Sherriff JL, Dhaliwal SS, Mamo JCL (2011) Tailored, iterative, printed dietary feedback is as effective as group education in improving dietary behaviours: results from a randomised control trial in middle-aged adults with cardiovascular risk factors. *Int J Behav Nutr Phys Act* 8:43.
70. Shiao SPK, Grayson J, Lie A, Yu CH (2018) Personalized Nutrition-Genes, Diet, and Related Interactive Parameters as Predictors of Cancer in Multiethnic Colorectal Cancer Families. *Nutrients* 10: 795.
71. Sunguya BF, Poudel KC, Mlunde LB, Urassa DP, Yasuoka J, Jimba M (2014) Poor nutrition status and associated feeding practices among HIV-positive children in a food secure region in Tanzania: a call for tailored nutrition training. *PLoS One* 9: 98308.
72. Walsh CM, Dannhauser A, Joubert G (2002) The impact of a nutrition education programme on the anthropometric nutritional status of low-income children in South Africa. *Public Health Nutrition* 5: 3-9.
73. Golzarand M, Mirmiran P, Jessri M, Toolabi K, Mojarrad M, Azizi F (2012) Dietary trends in the Middle East and North Africa: an ecological study (1961 to 2007). *Public Health Nutr* 15: 1835-1844.
74. UNESCO (2019) Benin.
75. Worldatlas (2017) What Languages are Spoken in Benin?

76. Crepin S, Houinato D, Nawana B, Avode GD, Preux PM, et al. (2007) Link between epilepsy and malnutrition in a rural area of Benin. Epilepsia 48: 1926-1933.

77. Meyer-Rochow VB (2009) Food taboos: their origins and purposes. J Ethnobiol Ethnomed 5: 18.

### Author Affiliations

[Top](#)

Lehrstuhl für Ernährungsphysiologie, Technische Universität München, Munich, Germany

### Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • [www.scitechnol.com/submission](http://www.scitechnol.com/submission)