



The Effect of Inexpensive and Safe Interventions in People with Dementia

Qingcheng Du*

Abstract

An increasing number of inexpensive and safe interventions, such as music therapy, dance therapy, yoga, massage therapy, storytelling and Tai Chi Chuan are utilized to treat dementia. A large number of clinical trials indicate that these inexpensive interventions show benefits in the treatment of dementia. Due to current evidence, these inexpensive and safe interventions will become a promising therapy for the treatment of dementia. In present, no published reviews summarize the effect of inexpensive and safe interventions in the treatment of dementia. The aim of this study is to provide a review of the scientific evidence on the effect of inexpensive and safe interventions in patients with dementia.

Keywords

Dementia; Interventions; Effect; Review

Introduction

Today, a total of nearly 47 million people worldwide suffer from dementia, approximately 10 million new cases diagnosed each year; by 2050 that figure is expected to nearly three times, to 130 million [1]. Dementia is a syndrome that affects memory, thinking and social abilities, which seriously is detrimental to normal activities of daily life [2]. In the elderly population, it is the most common mental disorders as well as the leading cause of death and disability among elderly individuals in ordinary people [3]. Thus, dementia has a serious threat to the health of the elderly.

The most common strategy of dementia patients is drug therapy. However, pharmacological treatment always results in severe complications because of hypersensitivity of brain in patients with dementia, and on account of their venerable age reducing hepatic and kidney drug metabolism [3,4]. Currently, there are an increasing number of investigations about non-pharmacological treatments for patients with dementia [5]. For example, cognitive stimulation therapy, and physical exercise can lead to modest benefits on cognition in the treatment of people with dementia [5-7]. More and more inexpensive and safe interventions, such as music therapy, dance therapy, yoga, massage therapy, storytelling and Tai Chi Chuan are used to treat dementia. Most of clinical trials report that these inexpensive and safe interventions show benefits in the treatment of dementia. Thus, these inexpensive and safe interventions will become a promising therapy in the treatment of dementia.

In present, there is a lack of published reviews which summarize the effect of inexpensive and safe interventions in the treatment of dementia. Therefore, it is necessary to write a review of this area. The aim of this study is to present a review of scientific evidence on the effect of inexpensive and safe interventions in patients with dementia.

Music therapy

Music is a form of art, which can bring pleasure to people. Music therapy can provide patients with healing of encouragement and well-being of promotion. It may offer benefits by patient listens or patient participates. In addition, according to the selected music and the instruments used, music can be adjusted by the patient's preference.

In the treatment of dementia patients, music therapy can significantly improve next-morning cognitive functioning [8,9], reduce depression [9-11], anxiety [11-13], psychological symptoms (i.e., delusions, agitation, anxiety, apathy, irritability, aberrant motor activity, and night-time disturbances) [14,15] and stress [16], lower the occurrence of aggressive behaviors [17], alleviate agitated behaviors [12,18-21], improve psychological well-being [13], mood [22], self-esteem [10] and orientation [22], and remote episodic memory [22]. Additionally, live interactive music can show positive effects for dementia individuals with apathy [23]. However, a randomized controlled trial shows that the addition of active music therapy or individualized listening to music to standard care is not with a significant effect on behavioral and psychological symptoms of dementia patients [24]. Moreover, another randomized controlled trial shows that older people with dementia take part in the music programme without a significantly effect on both agitation and anxiety [25]. Furthermore, pre-recorded music has non-harmful effect in people with dementia but it shows less clearly benefits in dementia subjects [23].

In conclusion, music therapy shows benefits in dementia subjects. Different music may have diverse effects in people with dementia and dementia subjects' ways of participation may also affect the effect. In order to obtain the greatest therapeutic benefits, clinicians or caregivers should select suitable music and participation ways for dementia patients.

Dance therapy

Dance is a kind of art, moulding brain circuits, which may improve cognition and patterns of emotion and behaviour [26]. It can provide the elderly with a chance for greater social interaction, which substantially contributes to healthy ageing [27]. In addition, older adults with regular participation in set dancing are related to health benefits (physical activity, balance, functional capacity and quality of life) [28].

A randomized controlled trial shows that dementia patients in nursing homes can improve visuospatial abilities and prevent the deterioration of self-care after 9 weeks of dance movement therapy [29].

Nowadays, no adequate clinical trials evaluate the effects of dance in people with dementia. Thus, further studies should need to assess the effect of dance in dementia subjects.

*Corresponding author: Qingcheng Du, School of Pharmacy, Guangdong Medical University, Dongguan, Guangdong, 523808, China, Tel: +86 15625554996; E-mail: dqc20160415@126.com

Received: May 10, 2017 Accepted: June 16, 2017 Published: June 23, 2017

Massage therapy

Massage, with about 2500 years history, is documented in the oldest surviving medical book (Huangdi Neijing) in China [30]. It includes different types, such as Thai massage, Swedish massage, traditional Chinese massage, Shiatsu, myofascial trigger point release and reflexology [31,32]. Massage therapy has immediate or short-term pain relief, increases range of motion and improves neck disability for those who are with neck pain [33-35].

Massage therapy can reduce the levels of anxiety and agitation as well as have an improvement of somnolence in both children and the aged individuals with dementia [36,37]. Additionally, tactile massage can decrease the level of both aggressiveness and stress in dementia subjects [38]. Moreover, massage therapy can improve depression in patients with dementia [39]. For dementia patients, massage therapy has an improvement in behavior as well as sleep disturbances, and increases the participation in both eating and rehabilitation [40]. However, a clinical trial indicates that aroma hand massage therapy cannot significantly improve behavioral and psychological symptoms of dementia individuals [41].

Overall, massage therapy can bring many benefits to patients with dementia. The operator of massage or caregivers should select the suitable locations and ways of massage for dementia subjects because of different locations and ways of massage with different effects.

Yoga

Yoga, a mindful exercise intervention, its origination is the Buddhist tradition with approximately 5000 years history. Yoga has 13 types, of these types, Kundalini yoga with more meditative, Hatha yoga, Iyengar yoga, Ashtanga yoga, and Vinyasa yoga with more physical [42]. All these types of yoga need to have a union of mind, body, and spirit. A survey shows that those who do yoga exercise show benefits in anxiety, stress, asthma, high blood pressure, and depression [43].

Fan JT et al shows that yoga can lower blood pressure, decrease respiration rate, enhance cardiopulmonary fitness and body flexibility, improve muscle strength, endurance and balance, have an increase of joints motion, and significantly reduce depression state and problem behaviors in dementia subjects [44]. Besides, as an intervention for dementia patients, yoga can significantly improve the immediate and delayed recall of verbal, visual memory, attention and working memory, verbal fluency, executive function and processing speed [45]. Thus, yoga is with beneficial effects on both the physical and mental health in those with dementia.

Yoga, an inexpensive and safe intervention, has benefits effects in dementia subjects, with recommendations of clinical application, and it should be as one of the routine activities [46].

Storytelling

Storytelling, a natural as well as universal kind of communication exists in daily life [47]. More and more people have realized that patients and service users can influence, change and benefit clinical practice from a wealth of health-related story [48]. Storytellers, psychologists and social workers have researched the therapeutic effect of storytelling. Current evidence indicates that storytelling is good for treating disease, such as hypertension [49], chronic disease [50].

For dementia patients, storytelling is with the aid of remembering as well as making associations with situations that they are with the

previous experience in their lives [51]. Moreover, compared with control group, patients with dementia of storytelling group exhibited significantly higher pleasure at Week 3, Week 6, and Week 7 [52].

Storytelling as a caring intervention is with advantages for dementia patients. Further studies are required to verify the effect of storytelling in patients with dementia.

Tai Chi Chuan

Tai Chi Chuan is a popular exercise in China. Tai Chi Chuan as a movement therapy, an increasing number of studies have explored its therapeutic effects in various diseases. For example, it has beneficial effects in obese patients with depression symptoms [53], rheumatoid arthritis [54], low back pain [55].

For dementia patients, Tai Chi Chuan can keep certain areas of cognitive function or delay recession, even those people with significant cognitive impairment as well [56]. In addition, Tai Chi Chuan may have specific benefits in cognition in the treatment of dementia patients [57].

Tai Chi Chuan is used as a way of health and it is also as an intervention for patients with dementia [58]. More potential clinical interests of Tai Chi Chuan in people with dementia should be explored by further clinical trials.

Summary

Music therapy, dance therapy, yoga, massage therapy, storytelling and Tai Chi Chuan are carried out through a variety of approaches, which can improve the clinical efficacy of dementia. In addition, these interventions are inexpensive and have the advantage of non-toxic side effects, so they have important promotion significance and application prospects. However, the clinical advantages have not yet fully played, there are still many aspects of urgent need to improve. Large and multicenter randomized controlled trials are required to explore more potential interests of the inexpensive and safe interventions in dementia patients. Furthermore, high quality systematic review should assess the effect of the inexpensive and safe interventions in dementia patients in order to provide clinicians or patients with comprehensive evidence.

References

1. Terry Pratchett (2015) A global assessment of dementia, now and in the future. *Lancet* 386: 931.
2. <http://www.who.int/mediacentre/factsheets/fs362/en/>
3. Brookmeyer R, Gray S, Kawas C (1998) Projections of Alzheimer's disease in the United States and the public health impact of delaying disease onset. *Am J Public Health* 88: 1337-1342.
4. Kim Y, Wilkins KM, Tampi RR (2008) Use of gabapentin in the treatment of behavioural and psychological symptoms of dementia: a review of the evidence. *Drugs Aging* 25: 187-196.
5. Yiannopoulou KG, Papageorgiou SG (2013) Current and future treatments for Alzheimer's disease. *Ther Adv Neurol Disord* 6: 19-33.
6. Venturelli M, Scarsini R, Schena F (2011) Six-month walking program changes cognitive and ADL performance in patients with Alzheimer. *Am J Alzheimers Dis Other Demen* 26: 381-388.
7. Yaguez L, Shaw KN, Morris R, Matthews D (2011) The effects on cognitive functions of a movement-based intervention in patients with Alzheimer's type dementia: a pilot study. *Int J Geriatr Psychiatry* 26: 173-181.
8. Bruer RA, Spitznagel E, Cloninger CR (2007) The temporal limits of cognitive change from music therapy in elderly persons with dementia or dementia-like cognitive impairment: a randomized controlled trial. *J Music Ther* 44: 308-328.

9. Chu H, Yang CY, Lin Y, Ou KL, Lee TY, et al. (2014) The impact of group music therapy on depression and cognition in elderly persons with dementia: a randomized controlled study. *Biol Res Nurs* 16: 209-217.
10. Cooke M, Moyle W, Shum D, Harrison S, Murfield J (2010) A randomized controlled trial exploring the effect of music on quality of life and depression in older people with dementia. *J Health Psychol* 15: 765-776.
11. Guétin S, Portet F, Picot MC, Pommié C, Messaoudi M, et al. (2009) Effect of music therapy on anxiety and depression in patients with Alzheimer's type dementia: randomised, controlled study. *Dement Geriatr Cogn Disord* 28: 36-46.
12. Svansdottir HB, Snaedal J (2006) Music therapy in moderate and severe dementia of Alzheimer's type: a case-control study. *Int Psychogeriatr* 18: 613-621.
13. Sung HC, Lee WL, Li TL, Watson R (2012) A group music intervention using percussion instruments with familiar music to reduce anxiety and agitation of institutionalized older adults with dementia. *Int J Geriatr Psychiatry* 27: 621-627.
14. Raglio A, Bellelli G, Traficante D, Gianotti M, Ubezio MC, et al. (2008) Efficacy of music therapy in the treatment of behavioral and psychiatric symptoms of dementia. *Alzheimer Dis Assoc Disord* 22: 158-162.
15. Raglio A, Oasi O, Gianotti M, Manzoni V, Bolis S, et al. (2010) Effects of music therapy on psychological symptoms and heart rate variability in patients with dementia. A pilot study. *Curr Aging Sci* 3: 242-246.
16. Sakamoto M, Ando H, Tsutou A (2013) Comparing the effects of different individualized music interventions for elderly individuals with severe dementia. *Int Psychogeriatr* 25: 775-784.
17. Clark ME, Lipe AW, Bilbrey M (1998) Use of music to decrease aggressive behaviors in people with dementia. *J Gerontol Nurs* 24: 10-17.
18. Vink AC, Zuidersma M, Boersma F, de Jonge P, Zuidema SU, et al. (2013) The effect of music therapy compared with general recreational activities in reducing agitation in people with dementia: a randomised controlled trial. *Int J Geriatr Psychiatry* 28: 1031-1038.
19. Sung HC, Chang SM, Lee WL, Lee MS (2006) The effects of group music with movement intervention on agitated behaviours of institutionalized elders with dementia in Taiwan. *Complement Ther Med* 14: 113-119.
20. Ridder HM, Stige B, Qvale LG, Gold C (2013) Individual music therapy for agitation in dementia: an exploratory randomized controlled trial. *Aging Ment Health* 17: 667-678.
21. Lin Y, Chu H, Yang CY, Chen CH, Chen SG, et al. (2011) Effectiveness of group music intervention against agitated behavior in elderly persons with dementia. *Int J Geriatr Psychiatry* 26: 670-678.
22. Särkämö T, Tervaniemi M, Laitinen S, Numminen A, Kurki M, et al. (2014) Cognitive, emotional, and social benefits of regular musical activities in early dementia: randomized controlled study. *Gerontologist* 54: 634-650.
23. Holmes C, Knights A, Dean C, Hodkinson S, Hopkins V (2006) Keep music live: music and the alleviation of apathy in dementia subjects. *Int Psychogeriatr* 18: 623-630.
24. Raglio A, Bellandi D, Baiardi P, Gianotti M, Ubezio MC, et al. (2015) Effect of Active Music Therapy and Individualized Listening to Music on Dementia: A Multicenter Randomized Controlled Trial. *J Am Geriatr Soc* 63: 1534-1539.
25. Cooke ML, Moyle W, Shum DH, Harrison SD, Murfield JE (2010) A randomized controlled trial exploring the effect of music on agitated behaviours and anxiety in older people with dementia. *Aging Ment Health* 14: 905-916.
26. Foster PP (2013) How does dancing promote brain reconditioning in the elderly?. *Front Aging Neurosci* 5: 4.
27. Merom D, Cumming R, Mathieu E, Anstey KJ, Rissel C, et al. (2013) Can social dancing prevent falls in older adults? a protocol of the Dance, Aging, Cognition, Economics (DAnCE) fall prevention randomised controlled trial. *BMC Public Health* 13: 477.
28. Shanahan J, Coman L, Ryan F, Saunders J, O'Sullivan K, et al. (2016) To dance or not to dance? A comparison of balance, physical fitness and quality of life in older Irish set dancers and age-matched controls. *Public Health* 141: 56-62.
29. Hokkanen L, Rantala L, Remes AM, Härkönen B, Viramo P, et al. (2008) Dance and movement therapeutic methods in management of dementia: a randomized, controlled study. *J Am Geriatr Soc* 56: 771-772.
30. Goats GC (1994) Massage--the scientific basis of an ancient art: Part 1. The techniques. *Br J Sports Med* 28: 149-152.
31. Patel KC, Gross A, Graham N, Goldsmith CH, Ezzo J, et al. (2012) Massage for mechanical neck disorders. *Cochrane Database Syst Rev* 12: Cd004871.
32. Kumar S, Beaton K, Hughes T (2013) The effectiveness of massage therapy for the treatment of nonspecific low back pain: a systematic review of systematic reviews. *Int J Gen Med* 6: 733-741.
33. Brosseau L, Wells GA, Tugwell P, Casimiro L, Novikov M, et al. (2012) Ottawa Panel evidence-based clinical practice guidelines on therapeutic massage for neck pain. *J Bodyw Mov Ther* 16: 300-325.
34. Topolska M, Chrzan S, Sapula R, Kowerski M, Sobon M, et al. (2012) Evaluation of the effectiveness of therapeutic massage in patients with neck pain. *Ortop Traumatol Rehabil* 14: 115-124.
35. Sherman KJ, Cook AJ, Wellman RD, Hawkes RJ, Kahn JR, et al. (2014) Five-week outcomes from a dosing trial of therapeutic massage for chronic neck pain. *Ann Fam Med* 12: 112-120.
36. Williams TI (2006) Evaluating effects of aromatherapy massage on sleep in children with autism: a pilot study. *Evid Based Complement Alternat Med* 3: 373-377.
37. Malaquin-Pavan E (1997) Therapeutic benefit of touch-massage in the overall management of demented elderly. *Rech Soins Infirm* 49: 11-66.
38. Suzuki M, Tatsumi A, Otsuka T, Kikuchi K, Mizuta A, et al. (2010) Physical and psychological effects of 6-week tactile massage on elderly patients with severe dementia. *Am J Alzheimers Dis Other Demen* 25: 680-686.
39. Rodriguez-Mansilla J, Gonzalez Lopez-Arza MV, Varela-Donoso E, Montanero-Fernandez J, Gonzalez Sanchez B, et al. (2015) The effects of ear acupuncture, massage therapy and no therapy on symptoms of dementia: a randomized controlled trial. *Clin Rehabil* 29: 683-693.
40. Rodriguez-Mansilla J, Gonzalez-Lopez-Arza MV, Varela-Donoso E, Montanero-Fernandez J, Jimenez-Palomares M, et al. (2013) Ear therapy and massage therapy in the elderly with dementia: a pilot study. *J Tradit Chin Med* 33: 461-467.
41. Yoshiyama K, Arita H, Suzuki J (2015) The effect of aroma hand massage therapy for people with dementia. *J Altern Complement Med* 21: 759-765.
42. Broad WJ (2012) The science of yoga: The risks and the rewards. Simon and Schuster, New York.
43. Medicine NcCaA. Yoga for Health: An Introduction.
44. Fan JT, Chen KM (2011) Using silver yoga exercises to promote physical and mental health of elders with dementia in long-term care facilities. *Int Psychogeriatr* 23: 1222-1230.
45. Hariprasad VR, Koparde V, Sivakumar PT, Varambally S, Thirthalli J, et al. (2013) Randomized clinical trial of yoga-based intervention in residents from elderly homes: Effects on cognitive function. *Indian J Psychiatry* 55: S357-S363.
46. Du Q, Wei Z (2017) The therapeutic effects of yoga in people with dementia: a systematic review. *Int J Geriatr Psychiatry* 32: 118.
47. Randall W, Josselson R (2014). *Front Matter. In the stories we are: An essay on self-creation.* (2nd edn), University of Toronto Press, Toronto, I-IV.
48. Haigh C, Hardy P (2011) Tell me a story--a conceptual exploration of storytelling in healthcare education. *Nurse Educ Today* 31: 408-411.
49. Houston TK, Allison JJ, Sussman M, Horn W, Holt CL, et al. (2011) Culturally appropriate storytelling to improve blood pressure: a randomized trial. *Ann Intern Med* 154: 77-84.
50. Gucciardi E, Jean-Pierre N, Karam G, Sidani S (2016) Designing and delivering facilitated storytelling interventions for chronic disease self-management: a scoping review. *BMC Health Serv Res* 16: 249.
51. Holm AK, Lepp M, Ringsberg KC (2005) Dementia: involving patients in storytelling--a caring intervention. A pilot study. *J Clin Nurs* 14: 256-263.
52. Phillips LJ, Reid-Armdt SA, Pak Y (2010) Effects of a creative expression intervention on emotions, communication, and quality of life in persons with dementia. *Nurs Res* 59: 417-425.
53. Liu X, Vitetta L (2015) The effects of tai chi in centrally obese adults with depression symptoms 879712.

54. Han A, Robinson V, Judd M, Taixiang W, Wells G, et al. (2004) Tai chi for treating rheumatoid arthritis. Cochrane Database Syst Rev 3: Cd004849.

55. Hall AM, Maher CG, Lam P, Ferreira M, Latimer J (2011) Tai chi exercise for treatment of pain and disability in people with persistent low back pain: a randomized controlled trial. Arthritis Care Res (Hoboken) 63: 1576-1583.

56. Cheng ST, Chow PK, Song YQ, Yu EC, Chan AC, et al. (2014) Mental and physical activities delay cognitive decline in older persons with dementia. Am J Geriatr Psychiatry 22: 63-74.

57. Lam LC, Chau RC, Wong BM, Fung AW, Lui VW, et al. (2011) Interim follow-up of a randomized controlled trial comparing Chinese style mind body (Tai Chi) and stretching exercises on cognitive function in subjects at risk of progressive cognitive decline. Int J Geriatr Psychiatry 26: 733-740.

58. Chen KM, Lin JN, Lin HS, Wu HC, Chen WT, et al. (2008) The effects of a Simplified Tai-Chi Exercise Program (STEP) on the physical health of older adults living in long-term care facilities: a single group design with multiple time points. Int J Nurs Stud 45: 501-507.

Author Affiliations

[Top](#)

School of Pharmacy, Guangdong Medical University, Dongguan, Guangdong, 523808, China

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission