



Thyroid Carcinoma after the Chernobyl Accident: Diagnosis, Treatment and Overtreatment

Sergei V. Jargin*

Letter to the Editor

On the basis of the linear no-threshold theory (LNT), Chernobyl accident (hereafter accident) was predicted to result in a considerable increase in radiation-induced cancer. In fact, there has been no cancer increase proven to be a consequence of the radiation exposure except for thyroid carcinoma in people exposed at a young age [1-3]. Appearance of radiogenic thyroid cancers after the accident is not denied, but their number has been overestimated due to the following mechanisms. Prior to the accident, the registered incidence of pediatric thyroid cancers was lower in the former Soviet Union (SU) than in other developed countries probably due to differences in diagnostic quality and coverage of the population by medical checkups [4,5]. The mass screening in contaminated territories after the accident detected not only small tumors but also advanced neglected cancers accumulated in the population, misclassified after the accident as aggressive radiogenic cancers. Besides, there was a pressure to be registered as Chernobyl victims to get access to benefits and health provisions [6]. There was no regular screening outside the contaminated areas, so that the cases brought from outside must have been averagely more advanced. Accordingly, the “first wave” thyroid cancers after the accident were on average larger and less differentiated than those detected later [7,8], as the pool of neglected cancers was gradually exhausted by the screening and reliability of the registration tended to improve with time. Admixture of old neglected cases explains the fact that Chernobyl-associated thyroid cancers “behaved in an aggressive fashion with poor clinical outcomes characterized by high frequency of extra-thyroidal extension, lymph node and distant metastases” etc. [9] The following citation is illustrative: “The tumors were randomly selected (successive cases) from the laboratories of Kiev and Valencia... [The cancers were] clearly more aggressive in the Ukrainian population in comparison with the Valencian cases” [10]. There is an explanation: more efficient cancer diagnostics in Western Europe. Further details and references are in [5,11].

The misclassification of neglected advanced cases as aggressive radiogenic cancers gave rise to the concept that supposedly radiogenic thyroid cancers, at least those from the “first wave” after the accident, were more aggressive than sporadic ones [8,12-15]. It had consequences for the practice: the surgical treatment of radiogenic thyroid carcinoma was recommended to be “more radical” [16]. After 1998-1999, the surgery in some institutions switched to a

more aggressive approach [14,17]. The following was recommended for Chernobyl-related pediatric thyroid carcinoma: “Radical thyroid surgery including total thyroidectomy combined with neck dissection followed by radioiodine ablation” [18] or high-dose external radiotherapy (40 Gy) [19]. Some experts regarded subtotal thyroidectomy to be “oncologically not justified” and advocated total thyroidectomy with prophylactic neck dissection [16,20-22]. Lesser resections were regarded to be “only acceptable in exceptional cases of very small solitary intrathyroidal carcinomas without evidence of neck lymph node involvement on surgical revision” [17]. It was written in a recent instructive publication that bilateral neck dissection must be performed in all cases independently of the tumor size, histology and lymph node status [23]. This approach is at variance with a more conservative treatment of thyroid carcinoma applied also in the settings of the Fukushima Daiichi nuclear accident [24]. The articles [25,26] were misquoted by Demidchik and Konratovich [22] advocating total thyroidectomy with bilateral neck dissection for all cases of pediatric thyroid cancer. The sources [27-29] were cited in support of the statement: “The most prevailing opinion calls for total thyroidectomy regardless of tumor size and histopathology” [17]. In fact, subtotal thyroidectomy was used or recommended in these articles, in some of them along with the total thyroidectomy [27-29]. Note that total thyroidectomy with neck dissection is associated with complications such as hypoparathyroidism and recurrent laryngeal nerve palsy [20,30-32]. Moreover, the majority of post-Chernobyl thyroid patients were young females potentially concerned with the cosmetic aspect.

Mechanisms of false-positivity have been discussed previously; among others, the misinterpretation of nuclear pleomorphism as a malignancy criterion of thyroid nodules was not unusual in the former SU of the 1990s due to the shortage of modern literature [33]. On the basis of contemporary morphological descriptions and images from the Russian-language literature on tumor pathology, in some cases no reliable differential diagnosis could be made; potentially misleading histological images from handbooks were reproduced in [11,34]. If a thyroid nodule is found by the screening, a fine needle aspiration biopsy (FNAB) is normally performed. Thyroid cytology is accompanied by some percentage of inconclusive results, when histological examination is indicated. In the former SU during the 1990s, this percentage was relatively high due to lacking experience with pediatric thyroid material, suboptimal quality of specimens, etc. [33] The surgical specimen was sent to a pathologist, who could be sometimes prone, after a complete removal of the nodule, to confirm malignancy even in case of uncertainty, which was not infrequent due to instable quality of specimens [11]. Frozen sections were sometimes used, being suboptimal for histological diagnostics of thyroid nodules. Radiophobia contributed to the overdiagnosis of cancer, which can be illustrated by the following citation: “Practically all nodular thyroid lesions, independently of their size, were regarded at that time in children as potentially malignant tumors, requiring an urgent surgical operation” [35]. Ultrasound devices were introduced into practice earlier than FNAB [4], which, considering the attitude to the thyroid nodules cited above, must have contributed to the overdiagnosis and overtreatment early in the 1990s. Furthermore, iodine deficiency on the contaminated territories and goiter associated with it [15] was a contributing factor because more thyroid abnormalities were found

*Corresponding author: Sergei V. Jargin, Peoples' Friendship University of Russia, Clementovski per 6-82, Moscow, Russian Federation, Tel: 74959516788; E-mail: sjargin@mail.ru

Received: December 12, 2017 Accepted: January 13, 2018 Published: January 19, 2018

by the screening, providing more opportunities for the overdiagnosis of malignancy; more details are in [5,11]. In conclusion, among factors contributing to the persistence of suboptimal methods are lack of scientific discussion and insufficient use of the international literature [36].

References

1. Anspaugh LR, Catlin RJ, Goldman M (1988) The global impact of the Chernobyl reactor accident. *Science* 242: 1513-1519.
2. Raabe OG (2011) Toward improved ionizing radiation safety standards. *Health Phys* 101: 84-93.
3. UNSCEAR (2008) Sources and Effects of Ionizing Radiation. Report to the General Assembly. Annex D: Health effects due to radiation from the Chernobyl accident. United Nations, New York.
4. Lushnikov EF, Tsyb AF, Yamashita S (2006) Thyroid cancer in Russia after the Chernobyl. *Meditsina*, Moscow.
5. Jargin SV (2014) Chernobyl-related cancer and precancerous lesions: Incidence increase vs. late diagnostics. *Dose Response* 12: 404-414.
6. Bay IA, Oughton DH (2005) Social and economic effects. In: Smith J, Beresford NA (eds) *Chernobyl - Catastrophe and Consequences*. Springer, Chichester 239-266.
7. Nikiforov Y, Gnepp DR (1994) Pediatric thyroid cancer after the Chernobyl disaster. Pathomorphologic study of 84 cases (1991-1992) from the Republic of Belarus. *Cancer* 74: 748-766.
8. Williams ED, Abrosimov A, Bogdanova T, Demidchik EP, Ito M, et al. (2004) Thyroid carcinoma after Chernobyl latent period, morphology and aggressiveness. *Br J Cancer* 90: 2219-2224.
9. Baloch ZW, Livolsi VA (2016) Pathologic effects of radiation on the thyroid gland. In: Del Pino J, Diaz MJ, Frejo MT (eds) *Thyroid Toxicity*. Bentham Science 141-163.
10. Romanenko A, Morell-Quadreny L, Ramos D, Nepomnyaschiy V, Voizianov A, et al. (2007) Author reply to: overestimation of radiation-induced malignancy after the Chernobyl accident. *Virchows Arch* 451: 107-108.
11. Jargin SV (2016) Back to Chernobyl: Some aspects of cancer diagnostics. *J Environ Stud* 2(1): 8
12. Zablotska LB, Nadyrov EA, Rozhko AV, Gong Z, Polyanskaya ON, et al. (2015) Analysis of thyroid malignant pathologic findings identified during 3 rounds of screening (1997-2008) of a cohort of children and adolescents from Belarus exposed to radioiodines after the Chernobyl accident. *Cancer* 121: 457-466.
13. Fridman M, Lam AK, Krasko O, Schmid KW, Branovan DI, Demidchik Y. (2015) Morphological and clinical presentation of papillary thyroid carcinoma in children and adolescents of Belarus: the influence of radiation exposure and the source of irradiation. *Exp Mol Pathol* 98: 527-531.
14. Iakovleva IN, Shishkov RV, Poliakov VG, Pankova PA (2008) Clinicomorphological peculiarities of thyroid cancer among children exposed to the Chernobyl disaster radiation. *Vopr Onkol* 54: 315-320.
15. IARC (2001) Thyroid cancer in young people. In: *Ionizing radiation, part 2: some internally deposited radionuclides. Views and expert opinions of an IARC Working Group on the Evaluation of Carcinogenic Risks to Humans*. Lyon, 14-21 June 2000. *IARC Monogr Eval Carcinog Risks Hum* 78 (Pt 2): 233-236.
16. Rumiantsev PO (2009) Thyroid cancer: modern approaches to diagnostics and treatment. *Geotar-Media*, Moscow.
17. Demidchik YE, Demidchik EP, Reiners C, Biko J, Mine M, et al. (2006) Comprehensive clinical assessment of 740 cases of surgically treated thyroid cancer in children of Belarus. *Ann Surg* 243: 525-532.
18. Demidchik YE, Saenko VA, Yamashita S (2007) Childhood thyroid cancer in Belarus, Russia, and Ukraine after Chernobyl and at present. *Arq Bras Endocrinol Metabol* 51: 748-762.
19. Mamchich VI, Pogorelov AV (1992) Surgical treatment of nodular goiter after the accident at the Chernobyl nuclear power station. *Klin Khir* (12): 38-40.
20. Demidchik EP, Tsyb AF, Lushnikov EF (1996) Thyroid carcinoma in children. Consequences of Chernobyl accident. *Meditsina*, Moscow.
21. Lushnikov EF, Vtiurin BM, Tsyb AF (2003) Thyroid microcarcinoma. *Moscow. Meditsina*.
22. Demidchik IuE, Kontratovich VA (2003) Repeat surgery for recurrent thyroid cancer in children. *Vopr Onkol* 49: 366-369.
23. Demidchik IuE, Shelkovich SE (2016) Thyroid tumors. *BelMAPO*, Minsk.
24. Sugitani I (2017) Management of papillary thyroid carcinoma in Japan. In: Yamashita S, Thomas G (eds) *Thyroid Cancer and Nuclear Accidents. Long-Term Aftereffects of Chernobyl and Fukushima*. Elsevier, London 185-194.
25. Segal K, Arad-Cohen A, Mechlis S, Lubin E, Feinmesser R (1997) Cancer of the thyroid in children and adolescents. *Clin Otolaryngol Allied Sci* 22: 525-528.
26. La Quaglia MP, Corbally MT, Heller G, Exelby PR, Brennan MF (1988) Recurrence and morbidity in differentiated thyroid carcinoma in children. *Surgery* 104: 1149-1156.
27. Danese D, Gardini A, Farsetti A, Sciacchitano S, Andreoli M, Pontecorvi A (1997) Thyroid carcinoma in children and adolescents. *Eur J Pediatr* 156: 190-194.
28. Arici C, Erdogan O, Altunbas H, Boz A, Melikoglu M, et al. (2002) Differentiated thyroid carcinoma in children and adolescents: clinical characteristics, treatment and outcome of 15 patients. *Horm Res* 57: 153-156.
29. Giuffrida D, Scollo C, Pellegriti G, Lavenia G, Iurato MP, et al. (2002) Differentiated thyroid cancer in children and adolescents. *J Endocrinol Invest* 25: 18-24.
30. Bohrer T, Pasteur I, Lyutkevych O, Fleischmann P, Tronko M (2005) Permanent hypoparathyroidism due to thyroid cancer surgical procedures in patients exposed to radiation in the Chernobyl, Ukraine, nuclear reactor accident. *Dtsch Med Wochenschr* 130: 2501-2506.
31. Henry JF, Gramatica L, Denizot A, Kvachenyuk A, Puccini M, Defechereux T (1998) Morbidity of prophylactic lymph node dissection in the central neck area in patients with papillary thyroid carcinoma. *Langenbecks Arch Surg* 383: 167-169.
32. Rybakov SJ, Komissarenko IV, Tronko ND, Kvachenyuk AN, Bogdanova TI, et al. (2000) Thyroid cancer in children of Ukraine after the Chernobyl accident. *World J Surg* 24: 1446-1449.
33. Jargin SV (2012) On the RET rearrangements in Chernobyl-related thyroid cancer. *J Thyroid Res* 2012: 373879.
34. Jargin SV (2011) Pathology in the former Soviet Union: scientific misconduct and related phenomena. *Dermatol Pract Concept* 1: 75-81
35. Lushnikov EF, Tsyb AF, Yamashita S (2006) Thyroid cancer in Russia after the Chernobyl. *Meditsina*, Moscow.
36. Jargin SV (2017) Invasive procedures with questionable indications: Prevention of a negligent custom. *J Surg Open Access* 3: 158.

Author Affiliation

Top

Peoples' Friendship University of Russia, Clementovski per 6-82, Moscow, Russian Federation

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission