



Clinical Image

## Tracheal Stenosis Due to an Abscess from Thyroid Tumor

Takeshi Kusunoki<sup>1\*</sup>, Hiroto Homma<sup>1</sup>, Yoshinobu Kidokoro<sup>1</sup>, Aya Yanai<sup>1</sup>, Kenji Sonoda<sup>1</sup>, Yuichiro Saikawa<sup>1</sup>, Ryo Wada<sup>2</sup> and Katsuhisa Ikeda<sup>3</sup>

An 88-year-old Japanese man underwent cervical spondylotic surgery under general anesthesia at a referring hospital. He soon fell into dyspnea after extubation. He underwent tracheotomy to keep the airway patent and was transferred to our hospital.

In the preoperative CT scan, tumors of the bilateral lobes of the thyroid were diffusely swollen and had grown into the space between the posterior wall of the trachea and esophagus, leading to the tracheal stenosis by compressing the posterior wall of the trachea (Figure 1).

Moreover, an upper region of right thyroid lobe showed a low density area 2 cm in the long diameter. We performed total thyroidectomy with preservation of the bilateral recurrent nerve. The postoperative histopathology was mostly adenomatous goiter in both lobes. By preoperative CT an upper region of the right thyroid lobe showed a low density area with the long diameter 2 cm adhering to the surrounding tissues and showed abscess findings without malignancy (Figure 2). Two regions (long diameter 4 mm and 12 mm) in the left lobe showed papillary carcinoma without capsular invasion.

The progress after surgery was good, and there was no dyspnea or vocal cord paralysis. In the postoperative CT, the trachea appeared to be free from stenosis and normal without tracheomalacia (Figure 3).




Figure 1: Tracheal stenosis by compressing the posterior wall of the trachea (\*).



Figure 2: Long diameter 2 cm adhering to the surrounding tissues and showed abscess findings without malignancy (\*).



Figure 3: The trachea appeared to be free from stenosis and normal without tracheomalacia.

Therefore, the stoma from the tracheotomy could be closed. Not all huge benign thyroid tumors with tracheal compression cause dyspnea. We reviewed the CT findings of tumors with dyspnea and found that tumors that occupied the space between the posterior wall of the trachea and esophagus lead to tracheal stenosis, as in our case [1,2].

### References

1. Sato K, Hanzawa H, Watanabe J, Takahashi S (2005) Differential diagnosis and management of air way obstruction in Riedel's thyroiditis: A case report. *Auris Nasus Larynx* 32: 439-443.
2. Ayabe H, Kawahara K, Tagawa Y, Tomita M (1992) Upper airway obstruction from a benign goiter. *Jpn J Surg* 22: 88-90.

**Citation:** Kusunoki T, Homma H, Kidokoro Y, Yanai A, Sonoda K, et al. (2019) *Tracheal Stenosis Due to an Abscess from Thyroid Tumor. J Otol Rhinol* 8:2.

\*Corresponding author: Takeshi Kusunoki, Department of Otorhinolaryngology, Juntendo University of Medicine, Shizuoka Hospital-1129, Nagaoka Izunokuni-shi, Shizuoka 410-2295, Japan, Fax: +81-55-948-5088; E-mail: ttksunoki001@aol.com

Received: April 22, 2019 Accepted: April 29, 2019 Published: May 06, 2019

### Author Affiliations

Top

<sup>1</sup>Department of Otorhinolaryngology, Juntendo University of Medicine, Shizuoka Hospital, Shizuoka, Japan

<sup>2</sup>Department of Pathology, Juntendo University of Medicine, Shizuoka Hospital, Shizuoka, Japan

<sup>3</sup>Department of Otorhinolaryngology, Juntendo University, Faculty of Medicine, Tokyo, Japan