



Whole Exome Sequencing found a Novel Truncating Mutation within *CNTNAP2* Gene in an Iranian Patient with Mental Retardation

Mehrnoosh Shokoohi¹, Mohammadreza Hajjari^{2*}, Javad Mohammadiasl J³ and Maryam Tahmasebi Birgani MT³

¹Noor Genetics Lab, Ahvaz, Iran

²Department of Genetics, Faculty of Science, Shahid Chamran University of Ahvaz, Ahvaz, Iran

³Department of Medical Genetics, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

*Corresponding author: Hajjari M, Department of Genetics, Faculty of Science, Shahid Chamran University of Ahvaz, Ahvaz, Iran, Tel: +98-6133330010; E-mail: Mohamad.hajari@gmail.com

Received date: June 30, 2018; Accepted date: August 21, 2018; Published date: August 28, 2018

Abstract

Intellectual disability (ID) is a major health problem mostly with an unknown etiology, affecting 1–3% of the general population. Discovering the genetic cause of these cases is dramatically challenging. Nowadays, the next-generation sequencing (NGS) technology provides advantages for the genetic diagnosis of ID. Here, by the help of Whole exome sequencing, we report an Iranian family with a child having non-syndromic ID (NS-ID) with autistic behaviors and seizure. The results revealed a novel nonsense mutation (c.3283CGA>TGA) located in an autozygous region on chromosome 7, leading to an amino acid change to terminal codon (p.Arg1095*) in *CNTNAP2*. The affected child was homozygous for the mutation and his parents were heterozygous as expected in autosomal recessive inheritance. In conclusion, our study identified a novel mutation in *CNTNAP2* gene, and showed that WES provides advantages for detecting novel ID associated variants and can greatly facilitate the genetic diagnosis of the disease.

Keywords: Intellectual disability; Whole exome sequencing; *CNTNAP2* gene; Novel nonsense mutation

Introduction

Intellectual disability (ID), a neurodevelopmental disorder, is demonstrated by considerable limitations not only in intellectual functioning but also in adaptive behavior that appears before the age of 18 [1-3]. However, although there is a higher ratio of males to females (Approximately 30%) among milder cases of ID, the ratio decreases followed by intelligence quotient (IQ) reduction [4-6]. Despite the classification of ID by moderate to severe /IQ level, ID can also be divided into two parts; syndromic intellectual disability (SID) and non-syndromic intellectual disability (NS-ID) [7]. On the one hand, In SID, patients suffer from clinical features or co-morbidities in addition to ID like for instance autism, Fragile X syndrome, and so on. On the other hand, there is discussion and challenge about the categorization of NS-ID [8-11].

In all of these cases, non-genetic factors (e.g., infection, trauma and brain damage, prenatal exposure to alcohol and other drugs, environmental exposure to toxins, psychiatric conditions and so on) cannot lead to cognitive impairment as severe as forms of ID with genetic factors and fundamental mutations ranging from large cytogenetic abnormalities to point mutations and even including epigenetic alterations [12,13]. In the past few years, hundreds of genes have been identified which is related to both SID and NS-ID [7,14,15]. However, it has been a challenge to discover genetic cause of individuals with intellectual disability without associated phenotypes [16,17].

Nowadays, exome sequencing which is a variant of next-generation sequencing (NGS) technology focuses on the coding regions of the genome, improves the efficiency of molecular diagnosis and contributes to discover novel mutations present in either sporadic or familial cases with non-specific phenotypes [18-20].

In summary, in this study, we introduced the genetic diagnosis of an Iranian family with one affected child who suffers from Mental retardation by Whole exome sequencing (WES). To our knowledge, this family might be another case of mental retardation caused by mutations located in the Contactin Associated Protein-Like 2 (*CNTNAP2*) gene (OMIM*604569). The data can provide the evidence of contribution for this mutation in further diagnosis.

Case Presentation

Clinical information of the patient

The family has two daughters that seem to be normal individuals without any clinical signs, and one son with features of mental retardation. These children were born of a consanguineous marriage (Figure 1). The patient had an abnormal behavior with below-normal IQ and impairment of adaptive skills. He was born after normal pregnancy, and his growth parameters were within the normal ranges. He had seizure several times since he was 2 years old. Afterward, he had language developmental delay and his language skill was limited to single words. Another serious problem included autistic behaviors which were displayed in this case. He started to walk at the age of 18 months and his muscle tonus was within normal.




Figure 1: The pedigree of Iranian family with mental retardation patient.

DNA isolation

Genomic DNA (gDNA) was isolated from peripheral blood of the proband and family members by Magcore HF16/HF48/Compact Nucleic Acid Extraction Kit.

Whole exome sequencing

DNA extracted from case 1 was sent to Macrogen company (South Korea) for whole exome sequencing to reveal if pathogenic variants exist.

Data analysis and validation

The Mutation Taster (<http://www.mutationtaster.org>), CADD, and DANN were applied to predict if the variants are pathogenic. Then, the variants were traced through the family members.

Results

Our patient had motor delay and limitations in motor skills as well as limited language comprehension. There were no distinguishing physical features. Severe intractable seizures began after 2 years old. He showed aberrant social interactions, mental retardation as well as autistic features.

The in-silico analysis gave information about a couple of candidate variants which were identified by WES data. Eventually, Sanger validation followed by co-segregation analysis were successfully utilized to detect a novel nonsense mutation in *CNTNAP2* gene (p.Arg1095*), which mapped to exon 20 of the *CNTNAP2* gene in the

patient (Figure 2). It was found that the parents of the patient were obligate carriers of this mutation (Table 1).




Figure 2: PCR products and its Sanger sequencings.

As can be seen from figures, fragments with 400 base pair in length were extended by PCR method. Afterwards, they were sequenced by sanger sequencing method. Regarding the sequences, a homozygote mutation in *CNTNAP2* gene (p.Arg1095*) was detected in the patient, and the parents of the patient were obligate carriers of this mutation. Additionally, the patient's sister and her husband were normal for this mutation.

Case No.	Sex	Age	Amino acid change	Nucleotide change	Zygoty
1	Male	-	Arg1095*	c.3283CGA>TGA	Het
2	Female	-	Arg1095*	c.3283CGA>TGA	Het
3	Male	17 yr	Arg1095*	c.3283CGA>TGA	Homo

In silico analysis of the variant:

Different *in silico* analysis softwares including SIFT, PROVEAN, CADD, and mutation taster predicted that the variant is pathogenic. The novel variant was neither found in EXAC nor 1000 genome browser (Table 2).

Prediction software	Prediction
Mutation taster	Disease causing
CADD	Deleterious
DANN	Deleterious

Discussion and Conclusion

Intellectual disability (ID) is a widespread neurodevelopmental disorder characterized by low IQ (below 70). Prevalence of ID varies from 1–3% [21–23]. To identify genetic causes of ID in the last decade, it has been suggested that Whole Exome Sequencing (WES) is a single experiment which can check nearly all the coding content of the

genome and the region likely to contain most of the disease-causing mutations [24–26]. Apart from this, there is a controversial issue whether functional studies are necessary to validate the causative role of the newly identified mutations [27,28].

In this study, we present the results of exome sequencing in a consanguineous Iranian ID patient. He had intellectual disability, speech problems and seizure. However, he did not have any Brain imaging. Candidate gene sequencing followed by exome and sanger sequencing identified a novel homozygous nonsense mutation (Chr.7: p.Arg1095*), in *CNTNAP2* gene that causes mental retardation, autistic behaviors and seizure, as the two unaffected parents share this rare heterozygous mutation in heterozygous form.

Recent studies have reported: 1) a deletion mutation related to cortical dysplasia, focal epilepsy, relative macrocephaly, and diminished deep-tendon reflexes [29], 2) homozygous or compound heterozygous mutations with clinical features like mental retardation, seizures, and hyperbreathing patterns, reminiscent of Pitt-Hopkins syndrome [30], 3) and homozygous or compound heterozygous truncating mutations and/or intragenic deletions [31] in the *CNTNAP2* gene. Further, it was identified a common variant in the *CNTNAP2* gene that was associated with increased risk for autism [32–34]. Pitt-Hopkins-like syndrome-1 (PTSL1) is an autosomal

recessive neurodevelopmental disorder characterized by delayed psychomotor development, intellectual disability, severe speech impairment or regression, and behavioral abnormalities. Most patients have onset of seizures within the first years of life. Some patients may have cortical dysplasia on brain imaging [31]. Autism, the prototypic pervasive developmental disorder (PDD), is usually apparent by 3 years of age. It is characterized by a triad of limited or absent verbal communication, a lack of reciprocal social interaction or responsiveness, and restricted, stereotypic, and ritualized patterns of interests and behavior [35,36]. As previously mentioned, although the biological function of the gene has not been well studied, knowledge about this novel mutation will provide insights that will increase our understanding of ID development. All in all, there is no doubt that finding the broad range of new mutations can help genetic counseling in families with affected individuals, particularly with ID patients.

References

1. Agha Z, Iqbal Z, Azam M, Ayub H, Vissers LE, et al. (2014) Exome sequencing identifies three novel candidate genes implicated in intellectual disability. *PLoS One* 9: e112687.
2. Lamichhane D (2013) New AP4B1 mutation in an African-American child associated with intellectual disability. *J Pediatr Genet* 2: 191-195.
3. Larti F, Kahrizi K, Musante L, Hu H, Papari E, et al. (2015) A defect in the CLIP1 gene (CLIP-170) can cause autosomal recessive intellectual disability. *Eur J Hum Genet* 23: 331-336.
4. Redin C, Gérard B, Lauer J, Herenger Y, Muller J, et al. (2014) Efficient strategy for the molecular diagnosis of intellectual disability using targeted high-throughput sequencing. *J Med Genet* 51:724-736.
5. Hu H, Haas SA, Chelly J, Van Esch H, Raynaud M, et al. (2016) X-exome sequencing of 405 unresolved families identifies seven novel intellectual disability genes. *Mol Psychiatry* 21: 133-148.
6. Bryant L, Lozynska O, Marsh A, Papp TE, van Gorder L, et al. (2018) Identification of a novel pathogenic missense mutation in PRPF31 using whole exome sequencing: A case report. *Br J Ophthalmol* 0: 1-7.
7. Kaufman L, Ayub M, Vincent JB (2010) The genetic basis of non-syndromic intellectual disability: A review. *J Neurodev Disord* 2: 182.
8. Mefford HC, Batshaw ML, Hoffman EP (2012) Genomics, intellectual disability, and autism. *N Engl J Med* 366: 733-743.
9. Lubs HA, Stevenson RE, Schwartz CE (2012) Fragile X and X-linked intellectual disability: Four decades of discovery. *Am J Hum Genet* 90: 579-590.
10. Lemke JR, Hendrickx R, Geider K, Laube B, Schwake M, et al. (2014) GRIN2B mutations in West syndrome and intellectual disability with focal epilepsy. *Ann Neurol* 75: 147-154.
11. Srivastava AK, Schwartz CE (2014) Intellectual disability and autism spectrum disorders: Causal genes and molecular mechanisms. *Neurosci Biobehav Rev* 46: 161-174.
12. Philips AK, Sirén A, Avela K, Somer M, Peippo M, et al. (2014) X-exome sequencing in Finnish families with Intellectual Disability-four novel mutations and two novel syndromic phenotypes. *Orphanet J Rare Dis* 9: 49.
13. Hamdan FF, Saitu H, Nishiyama K, Gauthier J, Dobrzyniecka S, et al. (2012) Identification of a novel in-frame de novo mutation in SPTAN1 in intellectual disability and pontocerebellar atrophy. *Eur J Hum Genet* 20: 796-800.
14. Duan Y, Lin S, Xie L, Zheng K, Chen S, et al. (2017) Exome sequencing identifies a novel mutation of the GDI1 gene in a Chinese non-syndromic X-linked intellectual disability family. *Genet Mol Biol* 40: 591-596.
15. Riazuddin S, Hussain M, Razzaq A, Iqbal Z, Shahzad M, et al. (2017) Exome sequencing of Pakistani consanguineous families identifies 30 novel candidate genes for recessive intellectual disability. *Mol Psychiatr* 22: 1604.
16. Zahir FR, Mwenifumbo JC, Chun HJ, Lim EL, Van Karnebeek CD, et al. (2017) Comprehensive whole genome sequence analyses yields novel genetic and structural insights for Intellectual Disability. *BMC Genom* 18: 403.
17. Birk E, Har-Zahav A, Manzini CM, Pasmanik-Chor M, Kornreich L, et al. (2010) SOBP is mutated in syndromic and nonsyndromic intellectual disability and is highly expressed in the brain limbic system. *Am J Hum Genet* 87: 694-700.
18. Carvill GL, Mefford HC (2015) Next-generation sequencing in intellectual disability. *J Pediatr Genet* 4: 128-135.
19. Grozeva D, Carss K, Spasic-Boskovic O, Tejada MI, Gecz J, et al. (2015) Targeted next-generation sequencing analysis of 1,000 individuals with intellectual disability. *Hum Mutat* 36: 1197-1204.
20. Aretini P, Mazzanti CM, La Ferla M, Franceschi S, Lessi F, et al. (2018) Next generation sequencing technologies for a successful diagnosis in a cold case of Leigh syndrome. *BMC Neurol* 18: 99.
21. Beaulieu CL, Huang L, Innes AM, Akimenko MA, Puffenberger EG, et al. (2013) Intellectual disability associated with a homozygous missense mutation in THOC6. *Orphanet J Rare Dis* 8: 62.
22. Davarniya B, Hu H, Kahrizi K, Musante L, Fattahi Z, et al. (2015) The role of a novel TRMT1 gene mutation and rare GRM1 gene defect in intellectual disability in two Azeri families. *PLoS One* 10: e0129631.
23. Luco SM, Pohl D, Sell E, Wagner JD, Dyment DA, et al. (2016) Case report of novel DYRK1A mutations in 2 individuals with syndromic intellectual disability and a review of the literature. *BMC Med Genet* 17: 15.
24. Topper S, Ober C, Das S (2011) Exome sequencing and the genetics of intellectual disability. *Clin Genet* 80: 117-126.
25. Rump P, Jazayeri O, van Dijk-Bos KK, Johansson LF, van Essen AJ, et al. (2015) Whole-exome sequencing is a powerful approach for establishing the etiological diagnosis in patients with intellectual disability and microcephaly. *BMC Med Genet* 9: 7.
26. Mohammadi-asl J, Hajjari M, Tahmasebi Birgani M, Riahi K, Nasiri H, et al. (2018) Exome sequencing revealed a novel deletion in the ERCC8 gene in an Iranian family with Cockayne syndrome. *Ann Hum Genet* 82: 304-308.
27. Chiurazzi P, Pirozzi F (2016) Advances in understanding-genetic basis of intellectual disability. *F1000Res* 5: 599.
28. Wou K, Weitz T, McCormack C, Wynn J, Spiegel E, et al. (2018) Parental Perceptions of Prenatal Whole Exome Sequencing (PPPWES) Study. *Prenat Diagn* 2018.
29. Strauss KA, Puffenberger EG, Huentelman MJ, Gottlieb S, Dobrin SE, et al. (2006) Recessive symptomatic focal epilepsy and mutant contactin-associated protein-like 2. *N Engl J Med* 354: 1370-1377.

30. Zweier C, de Jong EK, Zweier M, Orrico A, Ousager LB, et al. (2009) CNTNAP2 and NRXN1 are mutated in autosomal-recessive Pitt-Hopkins-like mental retardation and determine the level of a common synaptic protein in *Drosophila*. *Am J Hum Genet* 85: 655-666.
31. Smogavec M, Cleall A, Hoyer J, Lederer D, Nassogne MC, et al. (2016) Eight further individuals with intellectual disability and epilepsy carrying bi-allelic CNTNAP2 aberrations allow delineation of the mutational and phenotypic spectrum. *J Med Genet* 2016: 103880.
32. Bakaloglu B, O'Roak BJ, Louvi A, Gupta AR, Abelson JF, et al. (2008) Molecular cytogenetic analysis and resequencing of contactin associated protein-like 2 in autism spectrum disorders. *Am J Hum Genet* 82: 165-173.
33. Arking DE, Cutler DJ, Brune CW, Teslovich TM, West K, et al. (2008) A common genetic variant in the neurexin superfamily member CNTNAP2 increases familial risk of autism. *Am J Hum Genet* 82: 160-164.
34. Alarcón M, Abrahams BS, Stone JL, Duvall JA, Perederiy JV, et al. (2008) Linkage, association, and gene-expression analyses identify CNTNAP2 as an autism-susceptibility gene. *Am J Hum Genet* 82: 150-159.
35. Bailey A, Phillips W, Rutter M (1996) Autism: Towards an integration of clinical, genetic, neuropsychological, and neurobiological perspectives. *J Child Psychol Psychiatry* 37: 89-126.
36. Risch N, Spiker D, Lotspeich L, Nouri N, Hinds D, et al. (1999) A genomic screen of autism: evidence for a multilocus etiology. *T Am J Hum Genet* 65: 493-507.