



## Diabetic Retinopathy Study *via* Metabolism Improvements and Chronic Diseases Medical Conditions Control of Hba1c, SBP, and Triglycerides (GH-Method: Math-Physical Medicine)

Janet Kwan\*

Department of Diabetes, Eclair Md Foundation, Eufaula, USA

\*Corresponding author: Kwan J, Department of Diabetes, Eclair Md Foundation, Eufaula, USA, E-mail: j.kwan@eclairemd.com

Received Date: 23 February, 2021; Accepted Date: 09 March 2021; Published Date: 16 March 2021

### Abstract

The author used his own medical data to investigate the impact on his diabetic retinopathy condition and its risk probability percentage or hazard ratio (HR) over a period of 7-years where they are deciphered and interpreted into recognizable images. Although the retina is easily visible, it is, ironically, the only major tissue affected by diabetes that cannot be biopsied in human.

**Keywords:** Diabetic retinopathy; Metabolism; Chronic diseases medical conditions

### Introduction

The diabetic retinopathy remains the leading cause of vision impairment and blindness. The risk of developing vision loss from diabetes is predicted to double over the next three decades.

The fundamental functions of retina are to capture photons, convert the photochemical energy into electric energy, integrate the resulting action potentials, and transmit them to the occipital lobe of the brain, where they are deciphered and interpreted into recognizable images. Although the retina is easily visible, it is, ironically, the only major tissue affected by diabetes that cannot be biopsied in human [1].

### Methods

The author suffered from type 2 diabetes (T2D) since 1996. By year 2010, he experienced cardiovascular disease, renal complications, bladder infection, foot ulcer, thyroid disorder, and vision problems. In July of 2010, three of his physicians warned him that he would have three to five remaining years to live. Therefore, he decided to study and research diabetes and its complications in order to save his own life [2,3]. For the past 10 years, he has spent 30,000 hours on endocrinology with a specialty in diabetes and food nutrition.

The following timetable illustrates the focused area of each stage:

- 2000-2013:Self-study diabetes and food nutrition, developing a data collection and analysis software

- 2014:Develop a mathematical model of metabolism, using engineering modeling and advanced mathematics

- 2015:Weight & FPG prediction models, using neuroscience

- 2016:PPG & HbA1C prediction models, using optical physics, artificial intelligence (AI), and neuroscience

- 2017:CVD and Stroke research, using segmentation analysis and pattern analysis

- 2018:Complications due to micro-vascular research, for example, renal, bladder, and foot

- 2019:CGM glucose big data analysis, using wave theory, energy theory, frequency domain analysis, quantum mechanics, and AI

- 2020:Geriatrics, Longevity, Diabetic Retinopathy, Diabetic Hyperthyroidism, linkage between Metabolism and Immunity

To date, he has collected ~ 2 million data regarding his medical conditions and lifestyle details. He has written 246 medical papers and made ~ 120 presentations at ~ 70 international medical conferences [4].

Under the leadership of American Diabetes Association (ADA), a group of 14 authors quoted 137 references to publish a long editorial article, "Perspectives in Diabetes: Diabetic Retinopathy, Seeing Beyond Glucose-Induced Microvascular Disease" [5].

Established neurobiological principles can inform us how diabetes impairs vision, and metabolism knowledge may lead to new treatments. Retina physiology may underlie its vulnerability to diabetes. The combination of high metabolic demand and minimal vascular supply may limit the inner retina's ability to adapt to the metabolic stress of diabetes. The pathogenesis of diabetic retinopathy includes glucose-mediated microvascular damage. Although microvascular changes are undeniably integral to retinopathy, the retina is actually a vascularized neural tissue, not a network of blood vessels. Diabetic retinopathy involves more than elevated glucose and microvascular lesions. Plenty of evidences for neural retinal involvement in diabetic retinopathy have already been presented [6,7].

To the best of our knowledge, there is no evidence that a primary, selective defect in vascular cells is sufficient to cause diabetic retinopathy. Clearly, it is essential to treat both the vascular and neural elements of the retina to preserve vision [3].

Excess glucose (elevated HbA1C) is the primary culprit in the development and progression of diabetic retinopathy. However, disordered lipid (especially triglycerides) and protein metabolism are also linked to the central biochemical abnormality in all forms of impaired insulin action.

This study was conducted in a dynamic cohort of 2,770 type 2 diabetes patients, recruited between April 2005 and July 2013 (~ 8 years) in Ireland. In this diabetic retinopathy paper, the authors demonstrated that Higher current values of HbA1C, systolic blood pressure (SBP), and triglycerides were associated with increased risk of referral diabetic retinopathy [2].

Based on the findings from the long editorial article, the author conducted a study on his own diabetic retinopathy development and progression for the past 7-years (2013-2019). He has collected and further calculated the following data categories associated with his own medical conditions:

Weight: 2 body weight, BMI

Glucose: 1 FPG, 3 PPG, 1 daily A1C

Blood Pressure: SBP, DBP, pulse

Lipid: Triglycerides, HDL, LDL

The data for the top three categories were collected (weight, glucose, and BP) or calculated (i.e. HbA1C, BMI and predicted glucoses) on a daily basis. However, his lipid results were obtained from 22 hospital lab-tests with an average testing period of every four months [8].

Initially, he listed his lab-tested Triglycerides data, and then extracted data from both measured HbA1C and SBP from his stored database to match the actual dates of his lab-test for triglycerides (a total of 22 dates). In summary, 38,385 data of these 2,555 days (7-years) were used in this study.

Next, he selected the following medical conditions as his baseline conditions (i.e. “normal conditions”) for his risk probability analysis.

Triglycerides: 150 mg/dL

SBP: 120

HbA1C: 6.0%

Finally, he applied a linear regression analysis model with 7 different cases to conduct his numerical analysis [9].


In addition, he also applied Cox proportional-hazards regression model to conduct one more set of calculation. The Cox Hazard Model can be expressed as follows:

## Results

Figure 1 shows his records of Triglyceride, SBP, and HbA1C, where his SBP curve is expressed with 90-days moving average data.

Established neurobiological principles can inform us how diabetes impairs vision, and metabolism knowledge may lead to new treatments. Retina physiology may underlie its vulnerability to diabetes. The combination of high metabolic demand and minimal vascular supply may limit the inner retina’s ability to adapt to the metabolic stress of diabetes. The pathogenesis of diabetic retinopathy includes glucose-mediated microvascular damage. Although micro vascular changes are undeniably integral to retinopathy, the retina is actually a vascularized neural tissue, not a network of blood vessels. Diabetic retinopathy involves more than elevated glucose and micro vascular lesions. Plenty of evidences for neural retinal involvement in diabetic retinopathy have already been presented [6,7].

This study was conducted in a dynamic cohort of 2,770 type 2 diabetes patients, recruited between April 2005 and July 2013 (~ 8 years) in Ireland. In this diabetic retinopathy paper, the authors demonstrated that Higher current values of HbA1C, systolic blood pressure (SBP), and triglycerides were associated with increased risk of referral diabetic retinopathy as shown in Figure 1 [2].




**Figure 1:** HbA1C, SBP, Triglycerides during 2013 – 2019.

After he matches the dates of both HbA1C and SBP with the actual lab-test dates of Triglycerides, he then normalized those daily medical values by using baseline or normal conditions. These original data and normalized results.


The author could not find the contribution margins (i.e. weighting factors) related to these three primary factors of diabetic retinopathy study (i.e. A1C, SBP, and Triglycerides) from the external references; therefore, he decided to conduct a sensitivity analysis by using a range of possible weighting factors. Figures 2, 3 and 4 reflect the operational results and his seven different cases of weighting factors [10-12].



**Figure 2:** Time series Line Chart of diabetic retinopathy risk probability % of 7 cases.



**Figure 3:** Bar Chart of diabetic retinopathy risk probability % of 7 cases.



**Figure 4:** Comparison of Case 6 between weighted risk probability (Sum) % and HR Model results.

Here is the list of these seven different weighting factors:

Case 1: Trig 33%, SBP 33%, A1C 33% (even distribution at 33% each)

Case 2: Trig 30%, SBP 30%, A1C 40% (A1C is slightly heavier, 40%)

Case 3: Trig 60%, SBP 20%, A1C 20% (Trig is the heaviest, 60%)

Case 4: Trig 20%, SBP 60%, A1C 20% (SBP is the heaviest, 60%)

Case 5: Trig 20%, SBP 20%, A1C 60% (A1C is the heaviest, 60%)

Case 6: Trig 20%, SBP 30%, A1C 50% (A1C is heavier, 50%)

Case 7: Trig 30%, SBP 36%, A1C 34% (using Ref. 2's HR findings as a clue or source)

From Figures 2, 3 and 4 we can observe the following conclusive phenomena:

(1) Upper bound: Both Case 5 (A1C 60%) and Case 6 (A1C 50%) have the highest risk probability % (106%-108%).

(2) Middle Ground: All of Case 7 (Ireland HR case), Case 1 (even distribution of 33% each), and Case 2 (A1C 40%) are within a range of moderate risk probability % (100-102%).

(3) Lower Bound: Both Case 3 (Trig 60%) and Case 4 (SBP 60%) have the lowest risk probability % (94-98%).

Based on the author's medical data and personal feelings of his diabetic retinopathy conditions at different progression stages, the upper bound curves, Case 5 and Case 6, seem to be closely matched with the situation of his real conditions. On the opposite, the lower bound, Case 3 and Case 4, seem to be further apart from his real conditions.

Although lacking available medical test data, in general, he believes his overall diabetic retinopathy progression has slowed down for the past 7-years. His belief matched the trend of this set of curves.

Although both blood pressure and lipids share their responsibility of damaging the retina, based on this specific study, it seems that HbA1C plays a major role of his diabetic retinopathy conditions and developments.

In summary, he applied both linear regression analysis and Cox proportional-hazard model to calculate his "relative" risk probability % and his hazard ratio (HR) using 0.366 as its baseline. The comparison of these two approaches can be seen. Although the absolute values are slightly different on certain test dates, these two curves have an extremely high correlation coefficient (99.991%). These slight numerical differences are a result from HR Model utilizing exponential operation of the risk % value. The most important thing is that they demonstrate the relative risk probability % and relative hazard ratio (HR) of an upper bound Case 6 with an extremely high correlation [13,14].

## Conclusion

From the diabetic retinopathy sensitivity regression analysis results, it appears that diabetes, in particular HbA1C, plays a more dominating role as the murderer, while both hypertension (SBP) and hyperlipidemia (Triglycerides) play supporting roles as the accomplices. It is no wonder that the medical community calls it diabetic retinopathy! Nevertheless, the combination of these three chronic diseases may definitely cause severe damage to the retina. Furthermore, as stated in Reference 1, diabetic retinopathy is not only a metabolic microvascular blood vessel issue, it is also a serious neuroscientific problem. The author strongly agrees with this viewpoint. In his recent research work of glucose, he has identified the amounts for both fasting and postprandial glucoses' production and timing are controlled by the brain and neuro-system. He has already proven previously that diabetes itself is also closely related to our brain and nervous system.

As pointed out in Reference 1, this particular study may shed some light on using an approach to strengthen metabolic conditions, combination of HbA1C, SBP, and triglycerides in order to improve existing conditions of diabetic retinopathy in the patient.

## References

1. Kondo M, Kiyomizu K, Goto F, Kitahara T, Imai T, et al. (2015) Analysis of vestibular-balance symptoms according to symptom duration: dimensionality of the Vertigo Symptom Scale-short form. *Health Qual Life Outcomes* 13: 4.
2. Zigmond AS, Snaith RP (1983) The hospital anxiety and depression scale. *Acta Psychiatr Scand* 67: 361-370.
3. Matsudaira T, Igarashi H, Kikuchi H, Kano R, Mitoma H, et al. (2009) Factor structure of the hospital anxiety and depression scale in Japanese psychiatric outpatient and student populations. *Health Qual Life Outcomes* 7: 42.

4. Parnes LS, Agrawal SK, Atlas J (2003) Diagnosis and management of benign paroxysmal positional vertigo (BPPV). *CMAJ* 169: 681-693.
5. Cawthone TE, Hallpike CS (1957) A study of the clinical features and pathological changes within the temporal bones, brain stem and cerebellum of an early case of positional nystagmus of the so-called benign paroxysmal type. *Acta Otolaryngol* 48: 89-103.
6. Fujimoto C, Kawahara T, Kinoshita M, Kikkawa YS, Sugawara K, et al. (2018) Aging Is a Risk Factor for Utricular Dysfunction in Idiopathic Benign Paroxysmal Positional Vertigo. *Front Neurol* 9: 1049.
7. Iwasaki S, Chihara Y, Ushio M, Ochi A, Murofushi T, et al. (2011) Effect of the canalith repositioning procedure on subjective visual horizontal in patients with posterior canal benign paroxysmal positional vertigo. *Acta Otolaryngol* 131: 41-45.
8. von Brevern M, Bertholon P, Brandt T, Fife T, Imai T, et al. (2015) Benign paroxysmal positional vertigo: Diagnostic criteria. *J Vestib Res* 25: 105-117.
9. Jacobson GP, Newman CW (1990) The development of the dizziness handicap inventory. *Arch Otolaryngol Head Neck Surg* 116: 424-427.
10. Masuda K, Goto F, Fujii M, Kunihiro T (2004) Investigation of the reliability and validity of dizziness handicap inventory (dhi) translated into japanese. *Equilibrium Res* 63: 555-563.
11. Yardley L, Masson E, Verschuur C, Haacke N, Luxon L (1992) Symptoms, anxiety and handicap in dizzy patients: development of the vertigo symptom scale. *J Psychosom Res* 36: 731-741.
12. Hsu CL, Tsai SJ, Shen CC, Lu T, Hung YM, et al. (2019) Risk of benign paroxysmal positional vertigo in patients with depressive disorders: a nationwide population-based cohort study. *BMJ Open* 9: e026936.
13. Wei W, Sayyid ZN, Ma X, Wang T, Dong Y (2018) Presence of Anxiety and Depression Symptoms Affects the First Time Treatment Efficacy and Recurrence of Benign Paroxysmal Positional Vertigo. *Front Neurol* 9: 178.
14. Kitahara T, Ota I, Horinaka A, Ohyama H, Sakagami M, et al. (2019) Idiopathic benign paroxysmal positional vertigo with persistent vertigo/dizziness sensation is associated with latent canal paresis, endolymphatic hydrops, and osteoporosis. *Auris Nasus Larynx* 46: 27-33.