

Diagnosis of Violations of Microcirculation in the Late Recovery Period after Thrombosis of Cerebral Vessels

Vatnikov YA*, Kulikov EV, Kubatbekov TS, Kuznetsov VI, Sturov NV, Shirmanov VI, Parshina VI and Krotova EA

Department of Veterinary Medicine, Peoples Friendship University of Russia (RUDN University), Moscow, Russia

*Corresponding author: Vatnikov YA, Department of Veterinary Medicine, Peoples Friendship University of Russia (RUDN University), Moscow, Russia, Email: ilmedv1@yandex.ru

Received Date: June 27, 2018 Accepted Date: Jan 07, 2019 Published Date: Jan 14, 2019

Abstract

There is no objection that many pathological manifestations after thrombosis of cerebral vessels are associated in many respects with microcirculation disorders. These dysfunctions of microcirculation on the side of the lesion in patients are increasingly considered a point of influence for the recovery of this contingent of patients. In this connection, it is required to clarify the state of microcirculation in post-stroke patients at the late stage of their rehabilitation. In the article, the state of microcirculation in patients was clarified 7 months after the stroke. It was found that the deterioration of microcirculation processes is very typical for this contingent of patients, especially on the paretic side. Apparently, improvement in the state of motor function in patients after thrombosis of cerebral vessels is associated with increased microcirculation in affected limbs and in healthy limbs. The disturbances of microcirculation persisting in patients give grounds to recommend such a combination of drug treatment and non-medicamentous effects in the late stages of recovery, using the latter particularly actively on the affected side.

Keywords: Thrombosis of cerebral vessels; Late recovery period; Microcirculation; Vascular tone; Blood flow

Introduction

The viability of mammals has long been in the focus of attention of modern researchers [1-3]. It becomes clear that it depends on the mass of environmental factors [4-7] and shifts in the body's homeostasis [8-10]. Particular attention, among other things, attracts to itself blood, which is the environment of the organism and capable, by changing liquid properties, to influence the state of many vital processes [11].

Hemorheology is an important basis for all life processes of mammals [12-14] and human [15,16]. It provides the necessary metabolic processes in the body [17,18] and maintains the necessary level of adaptation [19-21] to the constantly changing environmental conditions [22-24]. In this connection, hemorheological properties of blood in vessels of different caliber and their connection with hemostasis are increasingly being studied in animals [25] and in humans [26,27] in norm and pathology [28-30]. The modern medicine

pays special attention to hemorheological changes in vascular thrombosis [31-33]. These include cerebral stroke, which does not currently reduce its prevalence among the population of developed countries [34]. The prevalence of thrombosis of cerebral vessels is now very high and continues to grow. In this regard, the continuation of the study of the pathological manifestations of thrombosis of cerebral vessels, which greatly contributes to the disability of the population, is of particular importance. In patients who have suffered it, the bulk does not return to its habitual way of life and work activity [35].

A major problem for post-stroke patients is motor impairment. Currently, it is recognized that they are based largely on violations of peripheral microcirculation [36]. In this connection, these microcirculation disorders in the paretic limb are attracting increasing attention of researchers seeking to increase the effectiveness of complex multidisciplinary rehabilitation in this contingent [37]. In this connection, it is necessary to assess the level of microcirculatory disorders that occur after thrombosis of cerebral vessels at the later stages of their rehabilitation. This information can help improve the efficiency of rehabilitation of this category of patients and prolong their life.

Objective: to establish the severity of microcirculatory disorders in patients in the late recovery period after a stroke.

Materials and Methods

The research was approved by the Ethics Committee of Peoples Friendship University of Russia (record №5 from 12.05.2014). 62 patients with an age from 45 to 75 y, including 36 men and 22 women who had had an ischemic stroke earlier, were examined. All patients gave written informed consent. Their examination was performed in the late (after 7.0 ± 0.47 months) recovery period after thrombosis of cerebral vessels.

The presence of ischemic stroke in each patient was confirmed by anamnesis, clinical picture, laboratory indicators, and the results of computed tomography of the brain. All patients had mild or moderate hemiparesis.

The study included patients with arterial hypertension. Their studies excluded patients who had diabetes mellitus, severe cardiac, hepatic or renal insufficiency and oncological diseases.

The control group was represented by 34 clinically healthy volunteers aged 45-75 (15 men and 19 women).

The study was carried out using laser Doppler flowmetry (LDF) using the LAKK-04 device manufactured by Lazma (Russia). In the course of this study, several indicators of microcirculation were taken into account: MP (microcirculatory parameter), expressed in terms of perfusion. He spoke about the state of perfusion associated with erythrocyte exponents. The mean square deviation (RMS) of the counted signals from the mean value or the value of the flag was taken into account. Analysis of amplitude-frequency characteristics of hemodynamic changes in tissue vessels was carried out at frequencies from 0.01 to 1.2 Hz. In the blood flow fluctuations, the following intervals were taken into account in studies: very low frequency (0.01-0.03 Hz) oscillations (VLF), they reflected the role of humoral-metabolic influences on microhemodynamics; low frequency (0.05-0.15 Hz) oscillations (LF) caused by periodic contraction of smooth myocytes of arterioles; high frequency (0.2-0.4 Hz) oscillations

(HF) associated with the dynamics of pressure in venous vessels; high-frequency (1.0-1.2 Hz) fluctuations (CF) of the pulse. The study took into account the contribution (P %) of different rhythmic oscillations on the LDF-gram, which were estimated from their percentage of the value of the flaxmation [38]. The value of the index of flaxemias was calculated as a partial active change in blood flow in the skin: $IFM=ALF / (ANF+ACF)$. In the work, respiratory and occlusive tests were performed.

Statistical analysis of the results obtained in the work was carried out with the help of the program "Statistics 6.0" by applying the methods of parametric and nonparametric analysis (Student and Mann-Whitney criteria). The level of significance in the study was $p<0.05$ (Table 1).

Results and Discussion

In patients in the study during the late recovery stage after thrombosis of cerebral vessels, mild hemiparesis was detected and only 2 patients had moderate hemiparesis. All patients on the side of the paresis recorded pronounced disorders of microcirculation associated with inhibition of blood flow in the lumen of the capillaries and due to excessive aggregation of erythrocytes. Microcirculation parameters were variable, and the level of the cell would be reduced. In the

frequency-amplitude spectrum in all cases, the LF-rhythm waves were reduced. In patients, a slight decrease in the amplitude and level of the contribution of the HF rhythm was noted. This was accompanied by a slight decrease in the amplitude and the value of the contribution of the CF rhythm with a decrease in the value of the index of fluctuations. On a healthy limb, all patients with microcirculatory disorders were not identified.

The negative changes in microcirculation in the affected limb found in patients are undoubtedly caused by a decrease in blood flow and a weakening of its outflow in the capillary bed. Apparently, this is due to a change in the lumen of the arteries and venules, their expansion and an increase in the degree of tortuosity of the venules.

All patients who underwent thrombosis of cerebral vessels, in addition to disorders of central hemodynamics, had microcirculation disorders mainly in the paretic limbs of patients.

Deterioration of peripheral hemodynamics was noted in all patients at a late stage of recovery after thrombosis of cerebral vessels on the affected side. Apparently, the positive dynamics achieved in the motor sphere in the patients observed at the late stage of recovery is associated with a positive dynamics of the microcirculation parameters on the lesion side and, to a lesser extent, on the healthy limbs.

Microcirculation parameters	Patients 7 months after the stroke, M ± m, n=62		Control group, M ± m, n=34
	Healthy limb	Paretic limb	
Microcirculatory parameter of perfusion unit	8.5 ± 0.41 p1<0.05	8.3 ± 0.34 p<0.05	8.6 ± 0.47
The value of the perfusion unit flag	1.02 ± 0.49 p1<0.05	0.94 ± 0.37 p<0.05	1.04 ± 0.055
The value of the index of fluctuations, conventional units	0.33 ± 0.021 p1<0.05	0.31 ± 0.016 p<0.05	0.34 ± 0.016
Very low frequency oscillations (VLF), Hz	0.42 ± 0.010 p1<0.05	0.37 ± 0.023 p<0.05	0.43 ± 0,014
Low frequency oscillations (LF), Hz	0.28 ± 0.007 p1<0.05	0.25 ± 0.009 p<0.05	0.28 ± 0.009
High frequency respiratory vibrations (HF), Hz	0.18 ± 0.009 p1<0.05	0.15 ± 0.012 p<0.05	0.18 ± 0.006
High-frequency pulse pulses (CF), Hz	0.65 ± 0.010 p1<0.05	0.58 ± 0.018 p<0.05	0.66 ± 0.012
Level of VLF contribution,%	27.0 ± 0.51 p1<0.05	26.0 ± 0.15 p<0.05	27.5 ± 0,59
Deposit level LF,%	17.6 ± 0.12 p1<0.05	15.0 ± 0.13 p<0.05	17.4 ± 0.25
Level of contribution HF,%	12.6 ± 0.17 p1<0.05	11.3 ± 0.14	12.7 ± 0.12
Deposit level CF,%	42.2 ± 0.19 p1<0.05	40.4 ± 0.15 p<0.05	42.3 ± 0.21
The value of vascular tone, unit	3.9 ± 0.20 p1<0.05	4.1 ± 0.16 p<0.05	3.8 ± 0.16
Severity of lowering of blood flow during a respiratory test,%	38.6 ± 0.23 p1<0.01	26.4 ± 0.20 p<0.01	40.0 ± 0,11
The size of the reserve of blood flow in capillaries,%	156.2 ± 0.62 p1<0.01	142.6 ± 0.033 p<0.01	159.1 ± 0.25

Note: p - reliability of differences between parameters of affected limb in patients and parameters of control group, p1 - reliability of differences between healthy and affected extremity in observed patients.

Table 1: The state of microrheological indicators in patients after ischemic stroke.

The microcirculation disorders that persist at the time of the examination give grounds to recommend the appointment of dezaggregants, venotonics, vasodilators, massage sessions, reflexology,

physiotherapy, and exercise therapy in patients with late stages of rehabilitation after thrombosis of cerebral vessels.

Conclusion

Impairment of peripheral hemodynamics is very typical for all patients in the late recovery period of ischemic stroke on the paresis side. There is reason to believe that the positive dynamics on the part of the motor sphere in patients in the late recovery period is associated with an improvement in microcirculation parameters on the paresis side and on healthy limbs. The remaining disorders of the microcirculatory link allow recommending patients who underwent ischemic stroke in the late recovery period the use of disaggregates and the use of non-drug effects, primarily on the diseased limb, to ensure the fullest possible rehabilitation.

References

1. Mitrokhina NV, Vatnikov YA, Sotnikova ED, Kulikov EV (2014) Assessment of the risk of osteosarcoma recurrence in canine long bone replants. *Biol Med* 6.
2. Kulikov EV, Vatnikov YA, Sotnikova ED, Seleznev SB, Troshina NI et al. (2015) Morphometric characteristics of the bone tissue structure in white Volga guinea-fowls. *Biol Med* 7.
3. Maksimov VI, Parakhnevich AV, Parakhnevich AA, Glagoleva TI, Kutafina NV (2017) Physiological Reaction of Erythrocytes' Micro Rheological Features in Newborn Piglets on Unfavourable Environmental Factors. *Ann Res Rev Biol* 17.
4. Vatnikov YuA, Sakhno NV, Sotnikova ED, Kulikov EV, Parshina VI et al. (2015) Clinical Control of Packed RBC Transfusion in Acute Surgical Pathology such as Gastric Dilation and Volvulus in Dogs. *Biomed Pharmacol J* 8.
5. Krasnikov AV, Annikov VV, Vatnikov YA, Sotnikova ED, Kulikov EV et al. (2016) Analysis of Dental Implants Biointegration in Animals. *Biol Med* 8.
6. Zavalishina SYu, Kutafina NV, Vatnikov YuA, Makurina ON, Kulikov EV et al. (2016) Platelet-Activity Dependence on the Age of Rats with Experimental Dyslipidemia. *Biol Med* 8.
7. Kulikov EV, Seleznev SB, Sotnikova ED, Yu AV, Kharlitskaya EV et al. (2016) The Morphological Aspects of Bone Marrow of Guinea Fowl of the Volga White Breed in Postembryonic Ontogenesis. *Res J Pharm Biol Chem Sci*. 7: 1148-1153.
8. Kulikov EV, Vatnikov YA, Parshina VI, Sotnikova ED, Vilkovskiy IF et al. (2017) Special aspects of the pathohistological diagnostics of familial shar-peï amyloidosis. *Asian J Pharm* 11: S152-S157.
9. Lenchenko EM, Vatnikov YA, Sotnikova ED, Kulikov EV, Gnezdilova LA et al. (2017) Experimental toxemia of chickens contaminated with yersinia enterocolitica bacteria. *Asian J Pharm* 11: S91-S96.
10. Seleznev SB, Kulikov EV, Vetoshkina GA, Vatnikov YA, Sotnikova ED et al. (2017) The evolution and structural organization of the organs of vertebrate immune system. *Asian J Pharm* 11: S84-S90.
11. Zavalishina SYu, Vatnikov YuA, Makurina ON, Kulikov EV, Sotnikova ED, Parshina VI, Rystsova EO, Kochneva MV, Sturov NV. (2017) Diagnostical Appreciation of Physiological Reaction of Intravascular Thrombocytes' Activity of Two-Years-Old Mice to Regular Physical Loads. *Biomedical & Pharmacology Journal* 10: 129-136.
12. Maksimov VI, Parakhnevich AV, Parakhnevich AA, Glagoleva TI, Kutafina NV (2017) Physiological Reaction of Erythrocytes' Micro Rheological Features in Newborn Piglets on Unfavourable Environmental Factors. *Ann Res Rev Biol* 17.
13. Maksimov VI, Parakhnevich AV, Parakhnevich AA, Glagoleva TI, Kutafina NV. (2017) Rheological Properties of Erythrocytes of Healthy Piglets during the Transition from Dairy to Vegetable Nutrition. *Ann Res Rev Biol* 16.
14. Maksimov VI, Parakhnevich AV, Parakhnevich AA, Glagoleva TI, Kutafina NV. (2017) Erythrocytes' Microrheological Features of Piglets during the Phase of Dairy-vegetable Nutrition after Damage or Common Supercooling. *Ann Res Rev Biol* 16.
15. Bikbulatova AA, Andreeva EG, Medvedev IN. (2017) Platelets' Functional Peculiarities in Persons of the Second Mature Age with Spinal Column Osteochondrosis of the Second Degree. *Ann Res Rev Biol* 21.
16. Medvedev IN, Skorjatina IA, Yu SZ (2016) Vascular control over blood cells aggregation in patients with arterial hypertension with dyslipidemia. *Cardior Ther Prevent* 15.
17. Zavalishina SYu, Medvedev IN. (2016) Features aggregation erythrocytes and platelets in old rats experiencing regular exercise on a treadmill. *Adv Gerontol* 29.
18. Makhov AS, Medvedev IN, Rysakova OG (2017) Functional features of hemostasis and physical fitness of skilled snowboarders with hearing impairment. *Teoriya i Praktika Fizicheskoy Kultury* 12: 27.
19. Skoryatina IA, Medvedev IN, Zavalishina SYu (2017) Antiplatelet control of vessels over the main blood cells in hypertensives with dyslipidemia in complex therapy. *Cardiovasc Therapy and Prevention* 16.
20. Simonenko VB, Medvedev IN, Tolmachev VV (2010) Effect of irbesartan of the function of hemocoagulative component of hemostasis in patients with arterial hypertension during metabolic syndrome. *Klinicheskaiya meditsina* 88: 27-30.
21. Yu ZS (2017) Physiological Features of Hemostasis in Newborn Calves Receiving Ferroglyukin, Fosprenil and Hamavit, for Iron Deficiency. *Ann Res Rev Biol* 14.
22. Medvedev IN (2017) The Impact of Durable and Regular Training in Hand-to-hand Fighting Section on Aggregative Platelet Activity of Persons at the First Mature Age. *Ann Res Rev Biol* 15.
23. Medvedev IN. (2017) Physiological Dynamics of Platelets' Activity in Aged Rats. *Ann Res Rev Biol* 18.
24. Medvedev IN (2016) Platelet functional activity in clinically healthy elderly. *Advances in gerontology* 29: 633-638.
25. Zavalishina SY (2017) Restoration of Physiological Activity of Platelets in New-Born Calves With Iron Deficiency. *Biomed Pharmacol J* 10.
26. Medvedev IN, Gromnatskii NI (2005) The influence of nebivolol on thrombocyte aggregation in patients with arterial hypertension with metabolic syndrome. *Klinicheskaiya Meditsina* 83: 31-33.
27. Medvedev IN (2015) Neuropsychological changes in cognitive function during the transition of the elderly in old. *Adv Gerontol* 28.
28. Medvedev IN, Amelina IV (2010) Evaluation of the relationship between chromosome aberrations and transcription activity of nucleolus organizer regions in indigenous Population of the Kursk Region. *Bull Exp Biol Med* 149.
29. Bikbulatova AA, Andreeva EG (2017) Dynamics of Platelet Activity in 5-6-Year Old Children with Scoliosis Against the Background of Daily Medicinal-Prophylactic Clothes' Wearing for Half A Year. *Biomed Pharmacol J* 10.

30. Bikbulatova AA. (2017) Dynamics of Locomotor Apparatus' Indices of Preschoolers with Scoliosis of I-II Degree Against the Background of Medicinal Physical Training. *Biomed Pharmacol J* 10.
31. Yu ZS, Medvedev IN (2017) Comparison of opportunities from two therapeutical complexes for correction of vascular hemostasis in hypertensives with metabolic syndrome. *Cardiovasc Res* 16.
32. Vorobyeva NV (2017) Physiological Reaction of Erythrocytes' Microrheological Properties on Hypodynamia in Persons of the Second Mature Age. *Ann Res Rev Biol* 20.
33. Skoryatina IA, Yu ZS, Makurina ON, Mal GS, Gamolina OV (2017) Some aspects of Treatment of Patients having Dislipidemia on the Background of Hypertension. *Prensa Med Argent* 103.
34. Ju LO, Glagoleva TI (2017) Physiological Activity of Platelet Aggregation in Calves of Vegetable Feeding. *Biomed Pharmacol J* 10: 1395-1400.
35. Ju LO, Medvedev IN, Fomina LL (2017) Platelets' Aggregative Properties of Irish Calves in the Phase of Dairy-vegetable Nutrition. *Ann Res Rev Biol* 16.
36. Kutafina NV (2017) Platelet Parameters of Holstein Newborn Calves. *Ann Res Rev Biol* 15.
37. Michael KM, Allen JK, Macko RF (2005) Reduced ambulatory activity after stroke: the role of balance, gait, and cardiovascular fitness. *Arch Phys Med Rehabil* 86: 1552-1556.
38. Kozlov VI, Azizov GA, Gurova OA (2012) Laser Doppler flowmetry in the evaluation of the condition and disorders of blood microcirculation. *Arch Balkan Med Union* 8.