



Evaluation of the Relationship between Primary Spontaneous Pneumothorax and Exercise and Patient Return to Previous Activities in Patients after Treatment

Manouchehr Aghajanzadeh^{1*}, Mostafa ziabari¹, Mohammad Reza Asgari¹, Mohamad Taghi Ashobi¹, Ali Alavi Fomani², Alirza Jafarnejad², Azita Tangestaninejad² and Yousha poorahmadi³

¹Department of Thoracic and General Surgery, Guilan University of Medical Sciences

²Department of pulmonology, Guilan University of Medical Sciences, Iran

³Inflammatory Lung Diseases Research Center, Guilan University of Medical Science, Iran

*Corresponding author: Manouchehr Aghajanzadeh, Department of Thoracic and General Surgery, Guilan University of Medical Sciences, Iran, E-mail: maghajanzadeh2003@yahoo.com

Received date: 24 December, 2021; Manuscript No. JSCP-21-34060;

Editor assigned date: 27 December, 2021; PreQC No. JSCP-21-34060(PQ);

Reviewed date: 07 January, 2022; QC No JSCP-21-34060;

Revised date: 18 January, 2022; Manuscript No. JSCP-21-34060(R);

Published date: 25 January, 2022; DOI: 10.4172/jscp.1000351.

Abstract

Introduction: The most common cause of Primary Spontaneous Pneumothorax (PSP) is sub pleural bleb apical rupture. Little is known about the relationship between PSP and exercise and return to exercise time. In this study, we tend to investigate the relationship between exercise and PSP and time of return to exercise and previous activities.

Method: This study was designed as a case series and the sample size included all patients diagnosed with PSP in Razi and Poursina and Aria hospitals of Rasht during 2015-2019 based on inclusion criteria. Variables were analyzed using Fisher's exact test, Chi square, Mann Whitney U and t-test (p<0.05).

Results: The most common type of Treatment in patients was Transaxillary Thoracotomy with Pleurodesis with iodine (TTP) in 58.2% and tube thoracostomy and pleurodesis in 41.7%, which was not statistically significant between athletes and non-athletes (p=0.806). Athletes who underwent TTP after 4 weeks and those treated with tube thoracostomy and pleurodesis after 8 weeks-12 weeks were advised to return to their previous activity. Of athletes, 9.5% had recurrence; of non-athletes, 9.8% had recurrence. Of athletes, 4.8% did not tolerate a return to their previous activity; of non-athletes, 7.3% did not tolerate returning to their previous activity regardless of treatment, and this difference was not significant.

Conclusion: Our study showed no significant difference between clinical manifestations and image findings as well as frequency of treatment and complications in both athlete and

non-athlete patients. There is no increase in recurrence and intolerance at the time recommended for return to previous activity.

Keywords: Primary spontaneous pneumothorax; Exercise and pneumothorax; Return to work

Introduction

Pneumothorax is defined as presence of air in the pleural cavity [1]. Pneumothorax can occur spontaneously or following direct trauma to the chest [2,3]. Spontaneous pneumothorax is divided into primary and secondary categories [4]. Primary Spontaneous Pneumothorax (PSP) occurs in the absence of any pathological underlying agent [5-7]. The most common cause of spontaneous pneumothorax is sub pleural bleb apical rupture [8]. PSP is more common in men [9]. Its incidence in men is 7 cases to 18 cases per 100,000 populations per year and in women 1 case to 6 cases per 100,000 populations per year [10-12]. Other risk factors include smoking, tall, young men aged 10 years to 30 years [13,14]. Moreover, 10% of PSP cases have a positive family history. Climate change including pressure and temperature reductions have also been identified as a contributing factor to PSP [15]. Symptoms of pneumothorax include sudden onset of severe chest pain in the affected side, dry cough and cyanosis, and even the patient may die of suffocation or acute cardiovascular failure [16]. Treatment can vary from inserting a chest tube by connecting to water seal to thoracoscopy and bleb resection and pleurodesis. But little has been done about the relationship between PSP and exercise and return to sport time, and there are little reports and surveys in literature on the relationship and there is disagreement about main question of when to return to exercise and daily activity. In this study we tend to investigate the relationship between exercise and PSP and time of return to exercise and previous activities.

Materials and Methods

Calculation of sample size, data collection, implementation and instruments

From 2013 to 2019, approximately 314 cases of pneumothorax were referred to Razi, Poursina and Aria hospitals of Rasht, of which 103 were included in our study of 103 patients studied, 21 were athletes and 82 were non-athletes. All of these patients underwent chest intubation after diagnosis of pneumothorax and were visited daily.

Statistical Analysis

SPSS software version 8 was used for data analysis. Descriptive data were calculated as mean and standard deviation. In this study, variables were analyzed using Fisher's exact test and Chi square test, Mann-Whitney-U test and t-test (p<0.05).

Results

In this study, 73.8% of patients were male and 26.2% were female; 20.4% of the patients were athletes and 79.6% were non-athletes; 85.7% were male athletes and 14.2% were female athletes resembling

the gender distribution of the disease. The most common clinical symptom was chest pain (41.7%). The most common exercise was bodybuilding (28.6%), followed by martial arts (19%), track and field and volleyball (14.3%). For 71.4% of athletes, the time interval between onset of symptoms and the last time of exercise was more than 7 days and 28.6% had activity during the last week. Of total subjects, only 3.9% had physical activity at the same time as PSP. The most common side of PSP was generally the right side (65%). The right side was the most prevalent side in athletes (52.4%) and the right side was the most common side in non-athletes (68.3%), which was

not statistically significant ($P=0.111$). Moreover, 36.9% of patients were smokers, 19% of athlete patients were smokers and 41.5% of non-athlete patients were smokers. Although the athletes were less smoker, this difference was not significant ($P=0.057$). Perhaps this difference was significant as the number of samples increased.

The most common finding in athlete and non-athlete patients was bleb observed on CT scan, which was not significantly different between athletes and non-athletes ($p=0.937$) (Table 1).

		Lung collapse	Air leak persistence for more than three days	Lung bleb in CT	Response to early treatment
Athlete	N	2	3	5	11
	%	9.50%	14.30%	23.80%	52.40%
Non-athlete	N	7	12	26	35
	%	8.50%	14.60%	31.70%	42.70%
Sum	N	9	15	31	46
	%	8.70%	14.60%	30.10%	44.70%

Table 1: Percentage frequency of findings in two athlete and non-athlete groups.

The most common type of treatment in patients was trans-axillary thoracotomy with pleurodesis with iodine (58.2%) and tube thoracostomy and pleurodesis (41.7%). Among athletes, TTP was the most common treatment (47.6%); among non-athletes, TTP was the most common treatment (60.9%), which is not significantly different between athletes and non-athletes based on Fisher exact test

($p=0.806$). Among athletes, 47.6% undergoing TTP were recommended to return to their previous activity after 4 weeks and 52.4% undergoing tube thoracostomy and pleurodesis were recommended to their previous activity after 8 weeks to 12 weeks. Among non-athletes, 60.9% undergoing TTP were recommended to return to their previous activity after 4 weeks and 39.1% treated conservatively were recommended to return to their previous activity after 12 weeks to 18 weeks (Table 2).

		Treatment		Sum
		Trans-axillary thoracotomy with pleurodesis with iodine	Tube thoracostomy and pleurodesis	
Athlete	N	10	11	21
	%	47.60%	52.40%	100%
Non-athlete	N	50	32	82
	%	60.90%	39.10%	100%
Sum	N	60	43	103
	%	43.70%	56.30%	100%

Table 2: Percentage frequency of treatments in athletic and non-athletic groups.

In this study, 93 patients did not have recurrence, 37.6% underwent tube thoracostomy and the rest underwent TTP. Not all patients with recurrence (10 patients) underwent thoracotomy and all underwent tube thoracostomy. This difference was statistically significant ($P<0.0001$) in all those who recurred and underwent tube thoracostomy and pleurodesis compared to patients who underwent

thoracotomy and did not recurred 96 patients had tolerated treatment, of which 44.8% underwent tube thoracostomy and 55.2% underwent thoracotomy. Of total number of patients who did not tolerate treatment, 28.6% underwent tube thoracostomy and 71.4% underwent thoracotomy. However, thoracotomy was more common among those who did not tolerate the treatment. But this difference was not significant ($P=0.464$) (Table 3).

Citation: Manouchehr Aghajanzadeh, Mostafa ziabari, Mohammad Reza Asgary, Mohamad Taghi Ashobi, Ali Alavi Fomani, Alirza Jafarnegad, Azita Tangestaninejad and Yousha poorahmadi (2022) Evaluation of the Relationship between Primary Spontaneous Pneumothorax and Exercise and Patient Return to Previous Activities in Patients after Treatment. *J Pls Sur Cos* 6:1.

			Treatment	
			Tube thoracostomy and pleurodesis	Trans-axillary thoracotomy with pleurodesis with iodine
Recurrence	Yes	N	10	0
		%	100%	0.00%
	No	N	35	58
		%	37.60%	62.40%
Sum		N	45	58
		%	43.70%	56.30%
Return to previous activity	Tolerated	N	43	53
		%	44.80%	55.20%
	Not tolerated	N	2	5
		%	28.60%	71.40%
Sum		N	45	58
		%	43.70%	56.30%

Table 3: Percentage frequency of recurrence and types of treatments.

Of 21 athletes, 52.4% underwent tube thoracostomy and returned to their previous exercise after 8 weeks to 12 weeks and 18.2% recurred. Of 21 athletes, 10 underwent thoracotomy and returned to exercise after 4 weeks with no recurrence. Of 82 non-athlete patients, 32

underwent tube thoracostomy, of which 25% recurred after 8 weeks. Comparison of athlete and non-athlete patients who underwent tube thoracostomy and had recurrence using fisher test showed no significant difference between them ($P=0.710$). Since there were no recurrences in patients undergoing thoracotomy, no comparison can be made between different subgroups (Table 4).

In people recurred after 8 weeks of conservative		Recurrence		Sum
		Yes	No	
Athlete	N	2	9	11
	%	18.20%	81.80%	100%
Non-athlete	N	8	24	32
	%	25%	75%	100%
Sum	N	10	33	43
	%	23.30%	76.70%	100%
Fisher's exact test: $P=0.710$				

Table 4: Percentage frequency of recurrence in patients of athletic and non-athletic groups after tube thoracostomy.

Of 21 athlete patients, 47.6% underwent Trans-axillary Thoracotomy and Pleurodesis (TTP), of whom 100% tolerated a return to their previous activity and no intolerance. Of 82 non-athlete patients, 60.9% underwent TTP, of whom 10% did not tolerate a return to their previous activity and 90% tolerated, and this difference between athletes and non-athletes was not significant ($P=0.578$). Of 21

athlete patients, 52.4% underwent tube thoracostomy, who were recommended to return to their previous activity after 8 weeks to 12 weeks, one of whom did not tolerate a return to previous activity and had recurrence. Of 82 non-athlete patients, 39.2% underwent tube thoracostomy, of whom 3.1% (one case) did not tolerate return to their previous activity after 8 weeks to 12 weeks, and this difference was not significant ($P=0.451$) (Table 5).

Citation: Manouchehr Aghajanzadeh, Mostafa ziabari, Mohammad Reza Asgary, Mohamad Taghi Ashobi, Ali Alavi Fomani, Alirza Jafarnegad, Azita Tangestaninejad and Yousha poorahmadi (2022) Evaluation of the Relationship between Primary Spontaneous Pneumothorax and Exercise and Patient Return to Previous Activities in Patients after Treatment. *J Pls Sur Cos* 6:1.

Patient		Tolerating return to previous activity (4 weeks after operation)		Sum
		Yes	No	
Athlete	N	10	0	10
	%	100%	0.00%	100%
Non-athlete	N	45	5	50
	%	90%	10%	100%
Sum	N	55	5	43
	%	91.70%	8.30%	100%
Patient		Tolerating return to previous activity (8 weeks after operation)		Sum
		Yes	No	
Athlete	N	10	1	11
	%	90.90%	9.10%	100%
Non-athlete	N	31	1	32
	%	96.90%	3.10%	100%
Sum	N	41	2	43
	%	95.30%	4.70%	100%
Fisher's exact test: P=0.578				

Table 5: Percentage frequency of tolerating return to previous activity among patients undergoing TTP in people who recurred after 4 weeks and 8 weeks.

Among athletes, recurrence rate was 9.5% and recurrence rate of non-athletes was 9.8%. Overall, this difference was not significant

(P=0.998). Of athletes, 4.8% did not tolerate their return to their previous activity, and 7.3% of non-athletes did not tolerate their return to their previous activity, regardless of the type of treatment, and this difference was not significant (P=0.998) (Table 6).

Patient		Recurrence		Sum
		Yes	No	
Athlete	N	2	19	21
	%	9.50%	90.50%	100%
Non-athlete	N	8	74	82
	%	9.80%	90.20%	100%
Sum	N	10	93	103
	%	9.70%	90.30%	100%
Patient		Return to previous activity	Sum	
		Tolerated	Not tolerated	
Athlete	N	20	1	21
	%	95.20%	4.80%	100%
Non-athlete	N	76	6	82
	%	92.70%	7.30%	100%

Sum	N	96	7	103
	%	93.20%	6.80%	100%

Table 6: Percentage frequency of recurrence and tolerance of return to previous physical activities in athlete and non-athlete groups.

Discussion

Recording PSP cases in athletes is particularly encouraged by practitioners to provide guidelines that improve the diagnosis and treatment of these conditions and they recommend providing guidelines in determining the safe return time. In this study, we tended to investigate the relationship between exercise and primary spontaneous pneumothorax and the time to return to exercise and previous activities of patients with PSP. The results of our study showed that 73.8% of the participants were male and 26.2% were female. The most common clinical symptom in our patients was chest pain (41.7%). Aghajanzadeh et al. studied PSP epidemiology in rasht during 2009-2014; in their study, 79.9% of patients were male. The most common clinical symptom (53.2%) was chest pain. Yoon et al. examined risk factors for spontaneous pneumothorax in 2013; they showed that 65.7% of patients were male, with the most common complaint being chest pain in 35.2% and shortness of breath in 21.7%. In addition to clinical findings, our study examined CT scans, with the most common finding in both athlete and non-athlete patients are multiple blebs ranging from 42.7% to 52.4%. In our study, 57.3% of patients did not give description of heavy activity. Yoon et al. noted that 53.4% of patients did not give a description of heavy activity, while Uramoto et al. showed that heavy activity is not a strong risk factor for predicting spontaneous pneumothorax.

The most common part of pulmonary involvement in spontaneous pneumothorax was in the right side of the chest (65%) and only 5.8% of patients had involvement in both sides of the lung. Yoon et al. reported that 3.2% of patients with spontaneous pneumothorax showed both sides of the lung involved. Few studies have investigated spontaneous pneumothorax in athletes, most of which were case reports. In our study, 103 patients with primary spontaneous pneumothorax were evaluated, of whom 21 were athletes (20.4%). In our study, we refer to an athlete who meets the following four criteria: 1) Performing sports to improve performance; 2) Participating actively in sports competitions; 3) Formally recognized in local or regional or national competitions; 4) Exercising and participating in most competitions included daily activity and mental focus. Of the sports studied, bodybuilding (n=6; 5.8%) was the most frequent sports in our study. It is worth noting that one of the most important issues in this disease is the high rate of recurrence despite different treatments available, ranging between 20%-50% for primary and 10%-20% for secondary cases. While recurrence rate in our study between athletes and non-athletes ranged from 9.5% to 9.8%, which may be due to the method used to examine patients undergoing CT scans and more patients receiving intervention. In our study, the most common type of treatment was transaxillary thoracotomy with pleurodesis with iodine, which had the same frequency in both groups. In athletic patients who underwent transaxillary thoracotomy, 100% had tolerated return to previous activity. Among non-athlete patients, only 10% could not tolerate a return to their previous activity. Of athlete and non-athlete patients who underwent tube thoracostomy and pleurodesis, 18.2%-25% had recurrence. None of our patients who underwent TTP

recurred. Overall, 9.5% of the athletes had recurrence and only 4.8% did not tolerate a return to their previous activity. We recommended 47.6% of athletes after four weeks based on TTP treatment received to return to their previous activity and recommended 52.4% to return to their activity after 8 weeks to 12 weeks. As noted earlier, only one case did not tolerate return to previous activity regardless of time interval of return to daily activity. Through a case report and empirical observation, Curtin et al. noted that, depending on general conditions and cause of spontaneous pneumothorax, it is possible to decide on sports activities of these athletes and the time to start exercises and sports activity should be limited at least for three weeks and activity must gradually increase at the time of return to exercises under supervision of the practitioner. In a case report by Davis et al., a track and field athlete was treated conservatively. The patient showed full recovery six days after onset of symptoms and started light activities after one week and started over the competition after three weeks. This patient had no problem after three months.

Conclusion

Overall, few studies have investigated spontaneous pneumothorax in athletes. Most studies have been in the form of case reports; however, in order to provide a guideline for diagnosis, treatment and timing of safe return of athletes with spontaneous pneumothorax to exercise, appropriate size of studies is required. Our study showed that both athlete and non-athlete patients were evaluated based on the same diagnostic and therapeutic indications. There were no significant differences in the results derived from clinical presentations and imaging findings and in terms of frequency of treatment and side effects (recurrence and tolerance to return to exercise and previous activity). At the recommended time to return to previous activity in our study, we did not see any increase in recurrence and intolerance compared to other studies. Therefore, it can be claimed that return to previous activity in athletes with spontaneous pneumothorax depends on general conditions and type of treatment, whether conservative or thoracotomy, and regardless of the exercise.

References

1. Sayar A, Kok A, Citak N, Metin M, Buyukkale S, et al. (2014) Size of pneumothorax can be a new indication for surgical treatment in primary spontaneous pneumothorax: A prospective study. *Ann Thorac Cardiovasc Surg* 20: 192-197. [Crossref], [Google Scholar], [Indexed]
2. Yoon JS, Suh JH, Choi SY, Kwon JB, Lee BY, et al. (2013) Risk factors for the development of reexpansion pulmonary edema in patients with spontaneous pneumothorax. *J Cardiothorac Surg* 8: 164. [Crossref], [Google Scholar], [Indexed]
3. Chou SH, Li HP, Lee YL, Lee JY, Chiang HH, et al. (2014) Video-assisted thoracoscopic surgery for postoperative recurrent primary spontaneous pneumothorax. *J Thorac Dis* 6: 52-5. [Crossref], [Google Scholar], [Indexed]
4. Inoue T, Ishida A, Nakamura M, Nishine H, Mineshita M, et al. (2013) Talc pleurodesis for the management of malignant pleural

- effusions in Japan. *Intern Med* 52: 1173-1176. [Crossref],[Google Scholar],[Indexed]
5. Xia H, Wang XJ, Zhou Q, Shi HZ, Tong ZH (2014) Efficacy and safety of talc pleurodesis for malignant pleural effusion: A meta-analysis. *PloS one* 9: e87060. [Crossref],[Google Scholar],[Indexed]
6. Isaka M, Asai K, Urabe N (2013) Surgery for secondary spontaneous pneumothorax: risk factors for recurrence and morbidity. *Interact Cardiovasc Thorac Surg* 17: 247-252. [Crossref],[Google Scholar],[Indexed]
7. Uramoto H, Shimokawa H, Tanaka F (2012) What factors predict recurrence of a spontaneous pneumothorax? *J Cardiothorac Surg* 7: 112. [Crossref],[Google Scholar],[Indexed]
8. Uramoto H, Tanaka F (2014) What is an appropriate material to use with a covering technique to prevent the recurrence of spontaneous pneumothorax?. *J Cardiothorac Surg* 9: 74. [Crossref],[Google Scholar],[Indexed]
9. Shi-ping L (2010) Diagnosis and treatment of primary spontaneous pneumothorax. *J Zhejiang Univ Sci B* 11: 735-744. [Crossref],[Google Scholar],[Indexed]
10. Tsukioka T, Inoue K, Oka H, Mizuguchi S, Morita R, et al. (2013) Pleurodesis with a 50% Glucose Solution in Patients with Spontaneous Pneumothorax in Whom an Operation is Contraindicated. *Ann Thorac Cardiovasc Surg* 19: 358-363. [Crossref],[Google Scholar],[Indexed]
11. Koji Y, Naoki H, Yasunori S, Ryo M, Hidenori K, et al. (2015) Initial Experience of single-incision thoracoscopic surgery for 100 patients with primary spontaneous pneumothorax. *Ann Thorac Cardiovasc Surg* 21: 513-516. [Crossref],[Google Scholar],[Indexed]
12. Haga T, Kurihara M, Kataoka H (2013) Spontaneous pneumothorax with persistent air leakage and invasive procedures. *Intern Med* 52: 2189-92. [Crossref],[Google Scholar],[Indexed]
13. Dearden AS, Sammon PM, Matthew EF (2013) In patients undergoing video-assisted thoracic surgery for pleurodesis in primary spontaneous pneumothorax, how long should chest drains remain in place prior to safe removal and subsequent discharge from hospital?. *Interact Cardiovasc Thorac Surg* 16: 686-91. [Crossref],[Google Scholar],[Indexed]
14. Noppen M (2010) Spontaneous pneumothorax: Epidemiology, pathophysiology and cause. *Eur Respir Rev* 19: 217-219. [Crossref],[Google Scholar],[Indexed]
15. Luh S (2010) Diagnosis and treatment of primary spontaneous pneumothorax. *J Zhejiang Univ Sci B* 11: 735-744. [Crossref],[Google Scholar],[Indexed]
16. Haga T, Kurihara M, Kataoka H, Ebana H (2013) Influence of weather conditions on the onset of primary spontaneous pneumothorax: Positive association with decreased atmospheric pressure. *Ann Thorac Cardiovasc Surg* 19: 212-215. [Crossref],[Google Scholar],[Indexed]