Research Article

Research and reviews in Psychology

A SCITECHNOL JOURNAL

Improving Psychotic Symptoms to Addicted Patients: the Antipsychotics Medication Helped Finding the Primary Diagnose


Vjollca N Ramigi*, Pierre Monpremier


Introduction

Does mental illness cause drug problems; it can be hard to tell which problem came first -the drugs or the mental illness. Having a mental illness can make a person more likely to abuse the drugs, make their symptoms feel better in the short term. Although there is a fewer study on comorbidity among youth, researcher suggests that adolescent with substance use disorders also have high rates of cooccurring mental illness; over 60 percent of adolescents in community -based substance use disorder treatment programs also meet diagnostic criteria for another mental illness. Data show high rates of comorbid substance using disorder and anxiety disorders, panic disorder, and post-traumatic disorder. The data show up the high rates also with affective disorder especially bipolar and psychotic disorder.

METHODOLOGY

Methodology of this investigation was discovering during the treatment for drugs abusing to find the primary illness. Well, finding the basis issue was helping for continuing treatment advice in future. Co-occurring Substance Use Disorder and Disorder and Serious Mental illness recently two years 2018 -2019 the patients of age 16 -30-year-old. In 2019, from 55 of patients, which were admitted in hospital; dominate age, over 30-year-old Female; 8; Male; 47?





CONCLUSION:

So, data showing, the response treatment of antipsychotic medication helped us discovering the primary diagnosis. Another side, it was important for parents which were interesting investing for rehabilitation treatment to understand the basic issue for their child. Knowing than they had another primary issue which inducted abusing the symptoms it's the low cost for them. Understanding than the primary treatment is with medication it's another way welcomed and satisfaction for

Research shows that people who have some kind of supportive relationship generally find it easier to tackle their problem. Having someone around to encourage. you is important because there is someone to talk to if times get tough, and to help you learn new ways of dealing with old problems. People with mental illness experience drug problems at far higher rates than general community. Studies suggest that both conditions are correctly diagnosed and receive the appropriate treatment.

Anyone who was vulnerable to, mental illness is therefore strongly discouraged from using drugs. Any time psychotic symptoms appear when taking prescription medication. Psychosis can also become more likely when these medication are abused .When it comes to nonprescription intoxicants the likelihood of psychotic symptoms appearing and what that looks like ,varies from substance to substance .For example ,taking a large amount of cocaine all at once can cause psychosis in minute .Psychosis from cocaine or amphetamine use typically produces persecutory delusions.

Received: September 24, 2020 Accepted: October 07, 2020 Published: October 14, 2020



All articles published in Research and reviews in Psychology are the property of SciTechnol, and is protected by copyright laws. Copyright © 2020, SciTechnol, All Rights Reserved.

^{*}Corresponding author: Vjollca N Ramiqi, Psychiatrist, Center Clinique University Republic of Kosovo, E-mail: cocir@hotmail.com

Citation: Ramiqi V N, Monpremier P, (2020) Improving Psychotic Symptoms to Addicted Patients: the Antipsychotics Medication Helped Finding the Primary Diagnose. rrpy 1:4.

References

- 1. The connection Between Substance use Disorder and Mental Illness: NIH USA government
- SAMHSA Substance Abuse and Mental Health Services Administration -2. USA government
- How to Identify Substance –Induced Psychosis _ American Addiction Centers; Reviewed by Scot Thomas. MD 3.
- 4. Substance -Induced psychosis an indicator of development of primary psychosis.; Am J. Psychiatry 2018,175(4);303-304.doi
- 5. Wofford CD.Kasckow JW,Scheller Gilkey G, Inderbitzin LB, .Substance use a powerful predictor of relapse in schizophrenia .Schizophrenic Resp, Dialogues in Clincal Neuroscience, 2006. 8(1): 37-43.
- AAC Remains Open and Committed Open Providing Critical Addiction 6. Treatment Amidst Coronavirus Sarah Harder March 16 2020
- 7. SANE Australia; Substance use &mental illness; do drug problems cause mental illness or does mental illness cause drug problems;

Author Affiliations

Тор

Psychiatrist, Center Clinique University Republic of Kosovo

Submit your next manuscript and get advantages of SciTechnol submissions

80 Journals

- 21 Day rapid review process
- 3000 Editorial team ٠
- 5 Million readers More than 5000 facebook ÷
- ۰.
- Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission