

Journal of Pharmaceutical Sciences and Emerging Drugs

A SCITECHNOL JOURNAL

Editorial

Maryam Hamzeloo-Moghadam

Editorial:

Statement of the Problem: Processed Capparis spinosa fruits, have been used in Iranian Traditional Medicine (ITM) for treatment of various diseases; the fruits are believed to have hot and dry temperament. The present study was designed in order to find the impact of processing on some phytochemical constituents and biological activities of Capparis spinosa. Methodology & Theoretical Orientation: The fruits were collected from Parsabad Moghan, Ardebil, Iran (2016) and processed with grape vinegar for 30 days. Total phenolics and alkaloids contents of the raw and processed fruits were determined. For quantitation of rutin and quercetin, R-HPLC with C18 column, methanol: formic acid 1% as the mobile phase (gradient), λmax 257

nm was used. Cytotoxic activity of the fruits was determined by MTT method. Antioxidant properties were evaluated by DPPH and FRAP techniques. Fingerpriting of raw and processed fruits were prepared using TLC with silica gel as the stationary phase and n-butanol: acetic acid: H2O 40:10:10 as the mobile phase. Findings: Total phenolics decreased in the processed fruits (7.5 times). Rutin concentration did not change during processing but quercetin amount decreased. Total alkaloids content was reduced as well. While, total phenolics decreased, antioxidant activities of the processed fruits increased using both DPPH and FRAP methods. No cytotoxic effect was observed for the raw fruits on normal human fibroblast cells (AGO1522) up to the concentration of 1000 µg/mL; however, decrease in the viability was observed for the processed fruits at this concentration. The fingerprints of the fruits were different which admitted the change in the fruit constituents due to processing. Conclusion & Significance: In general, it seems that processing with vinegar reduced the unpleasant taste of the plant due to alkaloids and increased the antioxidant effects; therefore it would be more suitable for use in some diseases such as diabetes and hepatitis.

