



## Outdoor Sports under UV Sunlight Exposure is not all Demerits - Considering its Advantages

Keiji Sugiura\* and Mariko Sugiura

### Abstract

Many physicians have warned of the risks of UV exposure during outdoor exercise. Outdoor exercise under sunlight, however, has not only adverse effects, but also many physical advantages. These include improved mental health, reduced mental stress, bone remodelling and increased muscle strength by increasing Vitamin D (25(OH) D3) levels, improved metabolism of glucose and lipids, growing up muscle mass and better skin by increased cortisol. Cortisol is also effective for antiaging by increasing anti-inflammatory factors, up-regulation of antioxidant enzymes, and improved skin blood flow by heating. Outdoor exercise with UV exposure has some good effects on the skin and physical body; females with regularly exercise outdoors are often perceived to be younger than their age and to possess good skin brightness and tone.

**Keywords:** Sports; Skin; Outdoor; UV; Athlete

### Introduction

Many physicians have warned of the adverse effects of exposure to outdoor Ultraviolet (UV) sunlight on the skin and eyes. We have met many females regularly engage in outdoor sports such as triathlon or marathons, however, have better skin brightness and tone than others their age. There are physical and psychological changes involved in enjoying outdoor exercise. These include mental hygiene, decreased stress, improved blood circulation and lip metabolism, increased muscular strength, bone remodelling (synthesizing vitamin D) and prevention of age-related neurodegenerative processes [1, 2]. These changes are affected by the quality and quantity of activity, environment and physical condition. One main factor of the environment is sunlight, UV. The skin of many female triathlon or marathon athletes showed good condition and tone not by skin care alone. They often appear cheerful and younger than their ages, and doing sports regularly with exposure to sunlight is not all about disadvantages. In this time, we focus on the merits of outdoor exercise in the sun.

### Environment and Exercise

Outdoor exercise involves a variety of environment factors such as weather, air, sunlight, landscape, people, insects, animals,

birds, bicycles and cars. These stimulate the brain or body via vision, respiration and skin, and such stimulation can cause physical changes. In particular, UV sunlight, only available outdoors, affects the human body differently than indoor sports.

Exercises are divided as follows: aerobic, resistance training, strength training and calisthenics. There are various reports about exercise types: aerobic exercise has therapeutic effects against mental illness [3, 4]; resistance training is effective for improving health [5, 6]; and physical exercise is the best way to prevent or delay aging [7]. The outdoor environment affects endurance sports such as triathlons and marathons because these sports are time consuming. Endurance sports should produce more multiplier physical and physis effects.

### Skin and Mental Hygiene

There have been many reports about the relation between physical exercise and mental health [8-11]. From the clinical evidence, exercise improves mental conditions (depression, Alzheimer's and Parkinson's disease) and the Quality of Life (QOL) [12-14]. It is well known that physic stresses often negatively affect skin conditions such as urticarial, acne, atopic dermatitis, prurigo and itching. On the other hand, improvement of mental hygiene can in some cases improve these skin conditions. Outdoor exercise reduces mental stress or confusion and promotes psychological benefits (happiness, satisfaction and improved QOL) [15-31]. Outdoor exercise involves one type of physical stress, the secretion of cortisol. UV sun exposure increases the active cortisol level in the skin by 11beta-hydroxysteroid dehydrogenase type 1 (11 $\beta$ -HSD1) [32, 33]. Cortisol is often used as a tablet or external medicine for treatment of dermatoses. Exercise under UV exposure should decrease mental stress and increase cortisol level, resulting in good skin tone and brightness. Moreover, dermatoses might be improved. Actually, UV is often used in treatment of skin diseases, and outdoor sports in sunlight could be a treatment for dermatoses.

### Skin, Bone and Muscle

Outdoor exercise with UV exposure is practical and effective for remodelling bone and increasing muscle. Remodelling bone is closely related to vitamin D (25(OH) D3) levels. There have been reports on 25(OH) D3 levels in athletes [34, 35], and 25(OH) D3 levels may depend on diet or exposure to sunlight. Vitamin D3 (25(OH) D3) from the diet has low-level absorption; therefore, most vitamin D3 (25(OH) D3) is synthesized by exposure to sunlight. Ultraviolet B (UVB) in the skin is crucial. Using SPF 15 sunblock decreases synthesis of vitamin D in the skin, and this reduction causes a phenomenon in the skin similar to that of melanin in African Americans and Africans [36, 37]. Protection from sunlight is effective for preventing skin cancer, but it is not favourable for the synthesis of vitamin D and bone remodelling. Sunlight is particularly essential to vitamin D synthesis for elder females, because they are likely to eat less and shelter from the sun using sunblock. Their bones are more fragile due to menopause. Recently, some females have taken extreme measures to shelter from sunlight using sunblock, long sleeves, or facemasks whenever they go out, and often exercise indoors to protect against sunburn. Because vitamin D absorption from the diet is low, these females often develop osteoporosis. Outdoor exercise is effective for menopausal-related

\*Corresponding author: Keiji Sugiura MD, PhD. Department of Environmental Dermatology & Allergology, Daiichi Clinic, Nittochi Nagoya Bld, 2F, 1-1 Sakae 2, Nakaku, Nagoya, 468-0008, Japan. Fax: +81-52-204-0835; E-mail: ksugiura@daichiclinic.jp

Received: November 11, 2020 Accepted: December 15, 2020 Published: December 23, 2020

and aging-related bone changes. It increases bone mass and strength, mitigates bone loss and prevents the death of bone cells. Exercise with UV exposure will be important for aging females.

Vitamin D and calcium are not absorbed sufficiently from the diet for bone remodelling; this requires exposure to sunlight (UV). Willis [38] reported that low levels of vitamin D in athletes could lead to reduced muscle strength and significantly increase the risk of bone injuries [38, 39]. Low levels of vitamin D can affect the exercise capacity of professional athletes [40, 41]. Outdoor sports with exposure to sunlight is necessary for remodelling bone and building muscle, and these physical changes result in better skin condition and health.

The synthesis of vitamin D in the skin during outdoor activities in sunlight (UVB) affects the circulating levels of 25(OH) D, which is related to glucose metabolism [42-44]. A 25(OH) D deficiency is a nascent symptom of metabolic disease [45]. In obesity, there can be hypovitaminosis D3, high BMI, and low physical performance [46]. Outdoor exercise is effective not only for slimming, but also to improve metabolism of glucose and lipids.

### Cortisol and skin

Human skin can produce and release cortisol in various environments; exposure to solar radiation [47-49], dryness [50], skin cancer [51], and trauma [52]. Exposure to UV increases active cortisol in the skin by 11beta-hydroxysteroid dehydrogenase type 1 (11 $\beta$ -HSD1) [32, 33]. This increasing cortisol might be caused by Interleukin-1 $\beta$  (IL-1 $\beta$ ) [50]. Outdoor sport is one kind of physical stress, and causes secretion of cortisol. Cortisol plays roles in increasing muscle mass and strength [53]. Most outdoor sports with a high degree of exposure to sunlight could be in a dry environment. Stress such as solar radiation, dry skin and physical exercise could produce cortisol secretion in the skin by IL-1 $\beta$ , causing increased muscle mass and strength, and better brightness and skin tone. The degree of improvement could depend on the quality and quantity of sports in sunlight.

### Exercise, Oxidative Stress and Aging

Dillard [54] first reported muscular exercise increases oxidant damage (stress), and Sies & Cadenas [55] defined oxidative stress as a disturbance between the pro-oxidant and antioxidant balance. Physical activity can cause oxidative stress, while it also involves preventive action of increasing antioxidants by multiple pathways [56-59]. Previous reports clarified that oxidative stress is an imbalance between the formation of oxidants (free radicals, reactive oxygen and nitrogen) and the body's antioxidant defence capacity [60, 61]. The low concentration of reactive oxygen species induces the antioxidant enzymes and other anti-inflammatory agents [62]. Exercise has the potential to increase anti-inflammatory factors, up-regulation of antioxidant enzyme gene promoter, and reduce adipose tissue [63].

Exercise-induced oxidative stress is subject to the age, gender, duration and intensity of exercise, training status, nutrition and oxidative stress levels at rest [64-69]. Long-duration high-intensity exercise can cause oxidative damage to skeletal muscle [70-72]. Chronic oxidative stress by overtraining is associated with chronic fatigue [73], illness [74], muscle atrophy [75] and long-term performance decrement [5]. Nutrition, especially vitamins C [76-78] and E [79-82], has a role in inhibiting exercise-induced oxidative stress and muscle injury. In athletes, reducing oxidative stress is crucial for recovery and maintenance of performance. Dietary vitamins C and

E after exercise are effective not only in athletes but also in people that exercise regularly (particularly endurance training or aerobic exercise) to reduce oxidative stress.

Previous reports showed regular swimming or running on a treadmill reduced oxidative damage in rats [83-85]. Mild oxidative stress induced by swimming or running may reduce oxidative damage to cells and tissues by operation of the antioxidant system. Mild oxidative stress plays a role in reducing risk of diseases associated with aging. The antiaging effects of physical exercise include increased muscle mass and bone density, enhanced metabolic function, improved respiratory function, increased neurogenesis, improved circulation and vascular systems, among others [86, 87]. People enjoying regular aerobic exercise such as swimming or running could reduce oxidative damage related to aging. Generally, skin blood flow is attenuated in aged skin, while in young skin, skin blood flow increases during whole body heating [88]. Aged skin in heating stress relies on NO-mediated vasodilation [89]. Endurance exercise causes functional vascular adaptations; elevation of skin blood flow [90] and endothelium-dependent vasodilatation [91-96]. This adaptation plays a role in maintenance of higher skin blood flow [97]. Moderate or endurance physical exercise and regular sports are effective for anti-aging and improve skin blood flow, resulting in younger appearance. In the result, female triathletes appear younger than their age.

### Conclusion

Endurance sports such as triathlons, marathons, long-distance bike rides, and open water swimming could reduce oxidative stress, produce antiaging effects, and increase skin blood flow. These effects could result in a younger appearance and improved skin tone and brightness. Outdoor exercise, especially endurance sports, under UV exposure is not all about demerits.

### Reference

1. Hillman CH, Erickson KI, Kramer AF (2008) Be smart, exercise your heart: exercise effects on brain and cognition. *Nature* 9: 58– 65.
2. Kramer AF, Colcombe SJ, McAuley E, Paige E Scalf, Kirk I Erickson (2005) Fitness, aging and neurocognitive function. *Neurobiol Aging* 26: 124–127.
3. Babyak M, Blumenthal JA, Herman SP, Khatri M, Doraiswamy, et al. (2000) Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosom Med* 62: 633-638.
4. Paluska SA, Schwenk TL (2000) Physical activity and mental health: Current Concepts. *Sports Med* 29: 167-180.
5. American Association of Cardiovascular and Pulmonary Rehabilitation. (1999) Guidelines for Cardiac Rehabilitation and Secondary Prevention Programs. 3rd Ed. Champaign, IL: Human Kinetics
6. American College of Sports Medicine (1998) Position stand: the recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sports Exerc* 30: 975-991.
7. Jonatan R, Francisco B, Ángel G (2006) Anti-aging therapy through fitness enhancement. *Clinical Interventions in Aging* 1: 213–220.
8. Stein D, Collins M, Daniels W, Timothy D Noakes, Michael Zigmond (2007) Mind and Muscle: the cognitive-affective neuroscience of exercise. *CNS Spectrum* 12: 19–22.
9. Vaynman S, Gomez-Pinilla F (2006) Revenge of the 'sit': how lifestyle impacts neuronal and cognitive health through molecular systems that interface energy metabolism with neuronal plasticity. *J Neurosci Res* 84: 699–715.
10. Duman R (2005) Neurotrophic factors and regulation of mood: role of exercise, diet and metabolism. *Neurobiol Aging* 26: S88–S93.
11. Dishman RK, Berthoud HR, Booth FW, Carl W Cotman (2006) Neurobiology of exercises. *Obesity* 14: 345–356.

12. Blumenthal JA, Babyak MA, Doraiswamy MP, Watkins L (2007) Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med* 69: 587–596.
13. Rolland Y, Pillard F, Klapouszczak A, Emma Reynish (2007) Exercise program for nursing home residents with Alzheimer's disease: a 1-year randomized, controlled trial. *J Am Geriatr Soc* 55: 158–165.
14. Hirsch M, Toole T, Maitland C, Robert A Rider (2003) The effects of balance training and high-intensity resistance training on persons with idiopathic Parkinson's disease. *Phys Med Rehabil* 84: 1109–1117.
15. Stuhl A, Porter H (2015) Riding the Waves. Therapeutic Surfing to Improve Social Skills for Children with Autism. *Ther Recreat J* 49: 253–256.
16. Zabriskie RB, Lundberg NR, Groff DG (2005) Quality of Life and Identity: The Benefits of a Community-Based Therapeutic Recreation and Adaptive Sports Program. *Ther Recreat J* 39: 176–191.
17. Mapes, N (2016) Green exercise and dementia. In *Green Exercise: Linking Nature, Health and Well-Being*:150–160
18. Crane M, Rissel C, Standen C, Greaves S (2014) Associations between the frequency of cycling and domains of quality of life. *Health Promot J Aust* 25: 182–185
19. Mutz M, Müller J (2016) Mental health benefits of outdoor adventures: Results from two pilot studies. *J Adolesc* 49:105–114.
20. Marselle MR, Irvine KN, Warber SL (2014) Examining Group Walks in Nature and Multiple Aspects of Well-Being. *Ecopyscholog* 6: 134–147.
21. Dorsch TE, Maxey M, Richards AR (2016) The effect of an outdoor recreation program on individuals with disabilities and their family members: A case study. *Ther Recreat J* 50:155–171.
22. Boddy K, Stein K, Whear R, Barton J, Depledge MH (2011) Does Participating in Physical Activity in Outdoor Natural Environments Have a Greater Effect on Physical and Mental Wellbeing than Physical Activity Indoors? A Systematic Review. *Environ Sci Technol* 45: 1761–1772.
23. Hartig T, Evans GW, Jamner LD, Davis DS, Garling T (2003) Tracking restoration in natural and urban field settings. *J Environ Psychol* 23:109–123.
24. Barton J, Griffin M, Pretty J (2011) Exercise, nature and socially interactive based initiatives improve mood and self-esteem in the clinical population. *Perspect Public Health* 132: 89–96.
25. Berman MG, Kross E, Krpan KM, Askren MK, Burson A, et al. (2012) Interacting with nature improves cognition and affect for individuals with depression. *J Affect Disord* 140: 300–305.
26. Pierskalla CD, Lee MA, Stein TV, Anderson DH, Nickerson R (2004) Understanding relationships among recreation opportunities: A meta-analysis of nine studies. *Leis Sci* 26: 163–180.
27. Hayhurst J, Hunter J, Kafka S, Boyes M. (2015) Enhancing Resilience in Youth through a 10-Day Developmental Voyage. *J. Adventure Educ. Outdoor Learn* 15: 40–52.
28. Barton J, Pretty J (2010) What is the Best Dose of Nature and Green Exercise for Improving Mental Health? A Multi-Study Analysis. *Environ Sci Technol* 44: 3947–3955.
29. Pretty J, Peacock, Hine R, Sellens M, South N, Griffin M (2007) Green exercise in the UK countryside: Effects on health and psychological well-being, and implications for policy and planning. *J Environ Plan Manag* 50: 211–231.
30. Clough P, Mackenzie SH, Mallabon L, Brymer E (2016) Adventurous physical activity environments: A mainstream intervention for mental health. *Sports Med* 46: 963–968.
31. Crust L, Henderson H, Middleton G (2013) The acute effects of urban green and countryside walking on psychological health: A field-based study of green exercise. *Int J Sport Psychol* 44: 160–177.
32. Boudon SM, Vuorinen A, Geotti-Bianchini P, Eliane Wandeler, Denise V Kretschmar, et al. (2017) 11 $\beta$ -hydroxysteroid dehydrogenase 1 inhibitors reduce cortisol levels in keratinocytes and improve dermal collagen content in human ex vivo skin after exposure to cortisone and UV. *PLoS ONE* 12: e0171079.
33. Nam J-J, Min J-E, Son M-H, Oh J-H, Kang S (2017) Ultraviolet- and infrared-induced 11 beta-hydroxysteroid dehydrogenase type 1 activating skin photoaging is inhibited by red ginseng extract containing high concentration of ginsenoside Rg3(S). *Photodermatol Photoimmunol Photomed* 33: 311-320.
34. Maimoun L, Mariano D, Couret I, Manetta J, Peruchon E (2004) Effects of physical activities that induce moderate external loading on bone metabolism in male athletes. *J Sports Sci* 22: 875-893.
35. K Solarz, A Kopeć, J Pietraszewska, F Majda, M Słowińska-Lisowska, et al. (2014) An Evaluation of the Levels of 25-Hydroxyvitamin D3 and Bone Turnover Markers in Professional Football Players and in Physically Inactive Men. *Physiol Res* 63: 237-243
36. Armas, L, Dowell, S, Akhter, M (2007) Ultraviolet-B radiation increases serum 25-hydroxyvitamin D levels: the effect of UVB dose and skin color. *J Am Acad Dermatol*. 57: 588-593.
37. Holick, M (2007) Vitamin D deficiency. *N Engl J Med* 357: 266-281.
38. Willis K, Peterson N, Larson-Meyer D (2008) willis ks, peterson nj, larson-meyer de: Should we be concerned about the vitamin D status of athletes? *Int J Sport Nutr Exerc Metab* 18: 204-224.
39. Tukaj C (2008) Adequate level of vitamin D is essential for maintaining good health. *Postepy Hig Med Dosw* 60: 502-510.
40. Cannell J, hollis w, sorensen M, taft T, anderson J (2009) Athletic performance and vitamin D. *Med Sci Sports Exerc* 41: 1102-1110.
41. Hamilton B, whiteley R, farooq A, chalabi H (2014) Vitamin D concentration in 342 professional football players and association with lower limb isokinetic function. *J Sci Med Sport* 17: 139-143.
42. Poolsup N, Suksomboon N, Plordplong N (2015) Effect of vitamin D supplementation on insulin resistance and glycaemic control in prediabetes: a systematic review and meta-analysis. *Diabet Med* 2016 33(3): 290-9.
43. Saraff V, Shaw N (2016) Sunshine and vitamin D. *Arch Dis Child* 101(2):190-2.
44. Peeling P, Fulton SK, Binnie M, Goodman C (2013) Training environment and Vitamin D status in athletes. *Int J Sports Med* 34: 248–252.
45. Sattar N (2012) Biomarkers for diabetes prediction, pathogenesis or pharmacotherapy guidance? Past, present and future possibilities. *Diabet Med* 29: 5–13.
46. Helmuth H, Sonja N, Delgerdalai B, Elisabeth P, Thomas P, et al. (2016) Low Vitamin D Levels Do Not Predict Hyperglycemia in Elderly Endurance Athletes (but in Controls). *PLoS ONE* 11.
47. Skobowiat C, Nejati R, Lu L, Williams RW, Slominski AT (2013) Genetic variation of the cutaneous HPA axis: An analysis of UVB-induced differential responses. *Gene*. 530 :1-7.
48. Skobowiat C, Sayre R, Dowdy J, Slominski AT (2013) Ultraviolet radiation regulates cortisol activity in a waveband-dependent manner in human skin ex vivo. *The British journal of dermatology* 168: 595–601.
49. Skobowiat C, Dowdy J, Sayre R, Tuckey R, Slominski A (2011) Cutaneous hypothalamic-pituitaryadrenal axis homolog: regulation by ultraviolet radiation. *Am J Physiol Endocrinol Metab* 301: 484-493.
50. Takei K, Denda S, Kumamoto J, Denda M (2013) Low environmental humidity induces synthesis and release of cortisol in an epidermal organotypic culture system. *Exp Dermatol* 22: 662–664.
51. Slominski T, Zmijewski A, Semak I, Blazej Z, Alexander P, et al. (2013) Cytochromes P450 and Skin Cancer: Role of Local Endocrine Pathways. *Anticancer Agents Med Chem* 14:77-96.
52. Vukelic S, Stojadinovic O, Pastar I, Morgan R, Agata K, et al. (2011) Cortisol synthesis in epidermis is induced by IL-1 and tissue injury. *J Biol Chem* 286: 10265-10275.
53. Gonzalo P , Raquel P , Nieves P , Beatriz M , Susana A, et al. (2015) EXERNET Study Group Biomarkers of physical activity and exercise *Nutr Hosp* 31: 237-244.
54. Dillard C, Litov E, Savin M, Dumelin E & Tappel L (1978) Effects of exercise, vitamin E, and ozone on pulmonary function and lipid peroxidation. *J Appl Physiol* 45: 927–932.
55. Sies H, Cadenas E (1985). Oxidative stress: damage to intact cells and organs. *Philos Trans R Soc Lond B Biol Sci* 311: 617–631.
56. McTiernan A (2008) Mechanisms linking physical activity with cancer. *Nature Rev* 8: 205-11.
57. Neilson H, Friedenreich C, Brockton N, Millikan R (2009) Physical activity and postmenopausal breast cancer: proposed biological mechanisms and areas for future research, *Cancer Epidemiol Biomarkers Prev* 18: 11-27.



58. Lynch B, Neilson H, Friedenreich C (2011) Physical activity and breast cancer prevention. In: Courneya KS, Friedenreich (eds) Physical Activity and Cancer, Recent Results in Cancer Research, Chap. 2. Springer Verlag, Berlin Heidenberg, 13-42.
59. Joanna K, Ewa D (2014) Oxidative Stress and Skin Diseases: possible role of physical activity. Asian Pac J Cancer Prev 15 (2): 561-568.
60. Sies H (1991) Oxidative stress. In: Sies H (ed.). Oxidative Stress: oxidants and antioxidants. San Diego: Academic Press 15-21.
61. Durackowa Z (2010) Some current sights into oxidative stress. Physiol Res 59: 459-469.
62. Gomes-Cabrera MC, Domenech E, Viña J (2008) Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. Free Radic Biol Med 44: 126-131.
63. Kaaks R, Lukanowa A, Kurzer MS (2002) Obesity, endogenous hormones, and endometrial cancer a synthetic review. Cancer Epidemiol Biomarkers Prev 11: 1531-1543.
64. Margaritelis NV, Kyparos A, Paschalis V, Theodorou AA, Panayiotou G, et al. (2014) Reductive Stress after Exercise: The Issue of Redox Individuality. Redox Biol 2: 520-528.
65. Bejma J, Ji LL (1999) Aging and Acute Exercise Enhance Free Radical Generation in Rat Skeletal Muscle. J Appl Physiol 87: 465-470.
66. Tiidus PM (2000) Estrogen and Gender Effects on Muscle Damage, Inflammation, and Oxidative Stress. Can J Appl Physiol 25: 274-287.
67. Paik IY, Jeong MH, Jin HE, Kim YI, Suh AR, et al. (2009) Fluid Replacement following Dehydration Reduces Oxidative Stress during Recovery. Biochem Biophys Res Commun 383: 103-107.
68. Bailey DM, McEneny J, Mathieu-Costello O, Henry RR, James PE, et al. (2010) Sedentary Aging Increases Resting and Exercise-Induced Intramuscular Free Radical Formation. J Appl Physiol 109: 449-456.
69. McGinnis G, Kliszczewicz B, Barberio M, Ballmann C, Peters B, et al. (2014) Acute Hypoxia and Exercise-Induced Blood Oxidative Stress. Int J Sport Nutr Exerc Metab 24: 684-693.
70. Powers S, Nelson W, Hudson M (2011) Exercise-induced oxidative stress in humans: cause and consequences. Free Radic Biol Med 51: 942-950.
71. Smith J (1995) Exercise, training and red blood cell turnover. Sports Med 19: 9-31.
72. Liu J, Yeo HC, Overvik-Douki E, Hagen T, Doniger SJ, et al. (2000) Chronically and acutely exercised rats: biomarkers of oxidative stress and endogenous antioxidants. J Appl Physiol 89: 21-28.
73. Kennedy G, Spence VA, McLaren M, Hill A, Underwood C, et al. (2005) Oxidative stress levels are raised in chronic fatigue syndrome and are associated with clinical symptoms. Free Radic Biol Med 39: 584-589.
74. Vider J, Lehtmaa J, Kullisaar T, Vihalemm T, Zilmer K, et al. (2001) Acute immune response in respect to exercise-induced oxidative stress. Pathophysiology 7: 263-70.
75. Xiao W, Chen P, Dong J (2012) Effects of overtraining on skeletal muscle growth and gene expression. Int J Sports Med 33: 846-853.
76. Alessio HM, Goldfarb AH, Cao G (1997) Exercise-Induced Oxidative Stress before and after Vitamin C Supplementation. Int J Sport Nutr 7: 1-9.
77. Ashton T, Young IS, Peters JR, Jones E, Jackson SK, et al. (1999) Electron Spin Resonance Spectroscopy, Exercise, and Oxidative Stress: An Ascorbic Acid Intervention Study. J Appl Physiol 87: 2032-2036.
78. Bryer SC, Goldfarb AH (2006) Effect of High Dose Vitamin C Supplementation on Muscle Soreness, Damage, Function, and Oxidative Stress to Eccentric Exercise. Int J Sport Nutr Exerc Metab 16: 270-280.
79. Sumida S, Tanaka K, Kitao H, Nakadomo F (1989) Exercise-Induced Lipid Peroxidation and Leakage of Enzymes before and after Vitamin E Supplementation. Int J. Biochem 21: 835-838.
80. Itoh H, Ohkuwa T, Yamazaki Y, Shimoda T, Wakayama A, et al. (2000) Vitamin E supplementation attenuates leakage of enzymes following 6 successive days of running training. Int J Sports Med 21: 369-374.
81. McBride JM, Kraemer WJ, Triplett-McBride T, Sebastianelli W (1998) Effect of Resistance Exercise on Free Radical Production. Med Sci Sports Exerc 30: 67-72.
82. Satchek JM, Milbury PE, Cannon JG, Roubenoff R, Blumberg JB (2003) Effect of Vitamin E and Eccentric Exercise on Selected Biomarkers of Oxidative Stress in Young and Elderly Men. Free Radic Biol Med 34: 1575-1588.
83. Radák Z, Kaneko T, Tahara S, Nakamoto H, Msasvai M, et al. (2001). Regular exercise improves cognitive function and decreases oxidative damage to proteins in rat brain. Neurochem Int 38: 17-23.
84. Radák Z, Kaneko T, Tahara S, Nakamoto H, Ohono H, et al. (1999) The effect of exercise training on oxidative damage of lipids, proteins and DNA in rat skeletal muscle: Evidence for beneficial outcome. Free Radic Biol Med 27: 69-74.
85. Nakamoto H, Kaneko T, Tahara S, Hayashi E, Naito H, et al. (2007) Regular exercise reduces 8-oxodG in the nuclear and mitochondrial DNA and modulates the DNA repair activity in the liver of old rats. Exp Gerontol 42: 287-295.
86. Garatachea N, Pareja-Galeano H, Sanchis-Gomar F, Santos-Lozano A, Fiuza-Luces C, et al. (2015) Exercise attenuates the major hallmarks of aging. Rejuvenation Res 18:57-89.
87. Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, et al. (2007) Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. Circulation 116: 1094-1105.
88. Minson CT, Wladkowski SL, Cardell AF, Pawelczyk JA, Kenney WL. (1998) Age alters the cardiovascular response to direct passive heating. J Appl Physiol 84: 1323-1332.
89. Holowatz LA, Houghton BL, Wong BJ, Wilkins BW, Harding AW, et al. (2003) Nitric oxide and attenuated reflex cutaneous vasodilation in aged skin. Am J Physiol Heart Circ Physiol. 284: H1662-1667.
90. Roberts MF, Wenger CB, Stolwijk JA, Nadel ER (1977) Skin blood flow and sweating changes following exercise training and heat acclimation. J Appl Physiol 43: 133-137.
91. Kvernmo HD, Stefanovska A, Kirkeboen KA, Osterud B, Kvernebo K (1998) Enhanced endothelium-dependent vasodilatation in human skin vasculature induced by physical conditioning. Eur J Appl Physiol Occup Physiol 79: 30-36.
92. Vassalle C, Lubrano V, Domenici C, Abbate AL (2003) Influence of Chronic Aerobic Exercise on Microcirculatory Flow and Nitric Oxide in Humans. Int J Sports Med 24: 30-35.
93. Franzoni F, Galetta F, Morizzo C, Lubrano V, Palombo C, et al. (2004) Effects of age and physical fitness on microcirculatory function. Clin Sci (Lond) 106: 329-335.
94. Lenasi H, Struel M (2004) Effect of Regular Physical Training on Cutaneous Microvascular Reactivity. Med Sci Sports Exerc 36: 606-612.
95. Wang JS (2005) Effects of exercise training and detraining on cutaneous microvascular function in man: the regulatory role of endothelium-dependent dilation in skin vasculature. Eur J Appl Physiol 93: 429-434.
96. Black MA, Green DJ, Cable NT (2008) Exercise prevents age-related decline in nitric-oxide-mediated vasodilator function in cutaneous microvessels. J Physiol 586: 3511-3524.
97. Grant H Simmons, Brett J Wong, Lacy A Holowatz, W Larry Kenney (2011) Changes in the control of skin blood flow with exercise training: where do cutaneous vascular adaptations fit in? Exp Physiol September 96: 822-828.

### Author Affiliations

Top

Department of Environmental Dermatology & Allergology, Daiichi Clinic, Nakaku, Nagoya, Japan