

Pathological Changes in Microcirculation in the Early Recovery Period of Ischemic Stroke

Binhong HU, Kulikov EV, Vatnikov YuA, Kuznetsov VI, Sturov NV, Shirmanov VI, Drukovsky SG, Petrov AK and Narresin Abdulaziz

Department of Veterinary Medicine, Peoples Friendship University of Russia (RUDN University), Moscow, Russia

*Corresponding author: Binhong Hu, Department of Veterinary Medicine, Peoples Friendship University of Russia (RUDN University), Moscow, Russia, E-mail: ilmedv1@yandex.ru

Received Date: May 21, 2018 Accepted Date: Jan 07, 2019 Published Date: Jan 14, 2019

Abstract

The rates of morbidity and mortality due to stroke among people of working age have been growing in recent years and have increased by more than 30% over the past 10 years. An important place among the complications of ischemic stroke is motor disorders, accompanied by persistent muscle hypertension and disorders of peripheral microcirculation. The purpose of the study: to reveal the features of microcirculatory disorders in patients in the early recovery period of ischemic stroke. Violations of peripheral hemodynamics are very typical for patients in the early recovery period of ischemic stroke. They are more pronounced on the side of the paresis. In the early recovery period, more severe forms of microcirculation disorder were observed more often on the paretic limb, accompanied by signs of a sharp decrease in blood flow, increased vascular tone with the phenomena of blood stagnation. The revealed disorders of the microcirculatory link allow the use of disaggregants, peripheral vasodilators, venotonics, massage, physiotherapy, reflexotherapy, and physical therapy with the involvement of the patient and healthy limbs to be recommended to patients in the early recovery period after ischemic stroke. All this should contribute to improving the effectiveness of rehabilitation of this category of patients during the early recovery phase.

Keywords: Ischemic stroke; Early recovery period; Microcirculation; Vascular tone; Blood flow

Introduction

Currently, researchers are paying increasing attention to the issues of hemostasis and hemorheology in mammals [1,2] and in humans [3,4]. This is due to the fact that dysfunctions leading to the violation of these indicators can strongly limit life expectancy [5-7], determine the state of health of the organism [8-10] and its overall functional status [11-13]. This has been proved at various experimental [14-17] and clinical models [18-20]. This is clearly confirmed by the example of patients with vascular dysfunction [21,22], arterial hypertension [23,24] who underwent thrombotic events, including ischemic stroke [25]. Particularly close attention of modern medicine to the latter

condition is due to the fact that cerebral stroke now occupies a leading position among the causes of disability and mortality of the population [26]. The rates of morbidity and mortality due to stroke among people of working age have been growing in recent years and have increased by more than 30% over the past 10 years. Stroke is very dangerous because about 30% of patients who have suffered from it constantly need outside help, another 20% cannot walk by them, and only 8% of surviving patients can later return to their previous work [27].

An important place among the complications of ischemic stroke is motor disorders, accompanied by persistent muscle hypertension and peripheral microcirculation disorders [28]. Currently, much attention is paid to secondary developmental disorders of microcirculation in the paretic limb [29]. It is believed that its optimization in the process of complex multi-profile rehabilitation is a reserve for the recovery of post-stroke patients [30,31].

The detection of microcirculatory disorders in patients with ischemic stroke and their timely correction in the early stages of neuro-rehabilitation will improve the quality of life and enhance the effectiveness of early rehabilitation in this category of patients.

The purpose of the study: to reveal the features of microcirculatory disorders in patients in the early recovery period of ischemic stroke.

Materials and Methods

The research was approved by the Ethics Committee of Peoples Friendship University of Russia (record №5 from 12.05.2014). A survey of 58 patients aged 43 to 75 years, of which 36 men and 22 women who underwent ischemic stroke were examined. All patients signed informed consent to participate in the study. The study was conducted in the early (after 4.0 ± 0.5 months) recovery period after the ischemic stroke.

The diagnosis was verified with the help of anamnesis, clinical picture, laboratory indicators, CT scan of the brain, ultrasonic dopplerography of the vessels of the head and neck. All patients had a motor deficit from mild to moderate hemiparesis, due to the presence of a lesion of the brain, confirmed by the method of neuroimaging. Data analysis with ultrasound Doppler scanning of the brachiocephalic arteries of patients showed the presence in half of patients of echographic signs of hypertensive angiopathy. In 34.5% of patients, stenosing arteriosclerosis of arteries of the carotid basin was visualized, 60% of them had moderate stenosis, and 40% had severe stenosis. 51.7% of patients had signs of non-stenosing atherosclerosis and only 13.8% of hemodynamically significant changes were not detected. Patients with hypertension could be included in the study. The study did not include patients with diabetes mellitus, with severe cardiac, hepatic or renal insufficiency. The control group is represented by 34 clinically healthy people aged 44 to 74 years, including 15 men and 19 women.

All the examined were carried out laser Doppler flowmetry (LDF) on the LAKK-04 instrument manufactured by Lazma (Russia). The study was carried out in the morning, at the same temperature in the room (about 21-24 ° C). Subjects during the study were in a sitting position. Before the study, the patients stayed calm for 15 minutes, did not smoke and did not eat or drink. The recording time was 6 minutes. The results were processed using a computer program (Table 1).

| Parameters of LDF-gram | Early recovery period, M ± m, n=58 | | Control, M ± m, n=34 |
|---|------------------------------------|-------------------------|----------------------|
| | Healthy limb | Paretic limb | |
| Parameter of microcirculation, perfusion units | 8.4 ± 0.35 | 8.2 ± 0.63 p<0.05 | 8.6 ± 0.47 |
| The level of the flask (mean square deviation), perfusion units | 0.98 ± 0.033 p1<0.05 | 0.91 ± 0.023 p<0.05 | 1.04 ± 0.055 |
| Index of flaxemias, conventional units | 0.33 ± 0.011 p1<0.05 | 0.29 ± 0.20 p<0.05 | 0.34 ± 0.16 |
| Very low frequency (VLF) oscillations, Hz | 0.42 ± 0.015 p1<0.05 | 0.35 ± 0.019 p<0.05 | 0,43 ± 0,014 |
| Low-frequency (LF) oscillations, Hz | 0.27 ± 0.008 p1<0.05 | 0.24 ± 0.010 p<0.05 | 0.28 ± 0.009 |
| High-frequency respiratory vibrations (HF), Hz | 0.16 ± 0.005 | 0.13 ± 0.011 p<0.05 | 0.18 ± 0.006 |
| High-frequency pulse (CF) oscillations, Hz | 0.63 ± 0.007 | 0.56 ± 0.020 p<0.05 | 0.66 ± 0.012 |
| %contribution VLF | 26.8 ± 0.40 p1<0.05 | 23.1 ± 0.33 p<0.05 | 27.5 ± 0.59 |
| %contribution LF | 18.5 ± 0.14 p1<0.05 | 16.2 ± 0.16 p<0.05 | 19.2 ± 0.25 |
| %contribution HF | 12.0 ± 0.20 p1<0.05 | 10.0 ± 0.15 p<0.05 | 12.7 ± 0.12 |
| %contribution CF | 42.0 ± 0.10 p1<0.05 | 38.1 ± 0.17 p<0.05 | 42.3 ± 0.21 |
| Vascular tone, units | 3.9 ± 0.15 p1<0.05 | 4.2 ± 0.20 p<0.05 | 3.8 ± 0.16 |
| Degree of decrease in blood flow in the respiratory sample,% | 33.2 ± 0.26 p1<0.01 | 24.1 ± 0.16 p<0.01 | 40.0 ± 0,11 |
| Reserve of capillary blood flow,% | 147.0 ± 0.71 p1<0.01 | 132.2 ± 0.045 p<0.01 | 159.1 ± 0.25 |

Table 1: Hemodynamic parameters in the early recovery period of ischemic stroke.

During the treatment of LDF-gram the following characteristics of microcirculation were determined: PM (microcirculation parameter), recorded in relative perfusion units (PE). Its value reflected the degree of perfusion, mainly the erythrocyte fraction, and the unit volume of tissue per unit time. The mean square deviation (RMS) of the detected Doppler signals from the mean value was taken into account. It characterized fluctuations in the flow of erythrocytes over time (or the level of the flask).

An amplitude-frequency analysis of hemodynamic rhythms of tissue blood flow oscillations was performed in the frequency range from 0.01 to 1.2 Hz. Among the considered fluctuations in blood flow reflected in the amplitude-frequency spectrum of the LDF-gram, very low-frequency (0.01-0.03 Hz) oscillations (VLF) are most

physiologically significant, characterizing the effect of humoral-metabolic factors on the state of microcirculation. Low frequency (0.05-0.15 Hz) oscillations (LF) were registered, which are caused by the spontaneous periodic activity of smooth myocytes in the arteriolar wall. A decrease in the amplitude of low-frequency flcomksmotion may indicate a spasm of the vessels of the microvasculature. High-frequency oscillations in the range of 0.2-0.4 Hz (HF-respiratory rhythms) were also taken into account. They are caused by changes in pressure in the venous section of the vascular bed. In addition, high frequency (1.0-1.2 Hz) pulse oscillations (CF) were recorded, which had a small fluctuation amplitude and caused changes in the rate of movement of erythrocytes in microvessels caused by changes in systolic and diastolic pressures.

In the course of the amplitude-frequency analysis of the LDF-gram, the contribution (P%) of various rhythmic components was considered, which was estimated from their power as a percentage of the total power of the spectrum of the flaxemias. An increase in the pulse wave amplitude means an increase in blood flow to the microcirculatory bed [32,33]. The ratio of active modulations of skin blood flow, caused by myogenic and neurogenic mechanisms, and passive modulations was calculated as the index of floxemias: $IFM=ALF / (ANF+ACF)$. It is known that the index of fluctuations decreases with decreasing mechanisms of active modulation (with spasm of bringing arterioles), as well as with strengthening of passive modulation mechanisms caused by stagnation of blood in the venous bed, changes in the rate of movement of erythrocytes in micro-vessels. In the study, respiratory and occlusive samples were performed [34].

Statistical analysis of the results of the study was carried out using the software "Statistics 6.0" with the use of parametric and nonparametric methods (Student's criteria, Mann-Whitney). Critical significance level was assumed to be $p < 0.05$.

Results and Discussion

In the early recovery period, all patients had hemiparesis of moderate severity. Analysis of their microcirculation showed that 100% of the patients had a stasis form of microcirculation due to a sharp decrease in blood flow in the capillary part of the microcirculatory bed and increased aggregation of erythrocytes. The microcirculation index was variable, the flax was reduced. In the frequency-amplitude spectrum, the vasomotor waves of the LF rhythm were reduced; there was a moderate decrease in the amplitude and percentage contribution of the HF rhythm, as well as a decrease in the amplitude and percentage contribution of the CF rhythm, as well as a decrease in the index of floxemia. At the same time, small breaches of microcirculation were observed on the healthy limb.

All patients with ischemic stroke along with a disorder of central hemodynamics had peripheral microcirculatory disorders on paretic and on healthy limbs. At the same time, violations of central hemodynamics were revealed in the form of echographically confirmed hypertonic angiopathy, stenotic and non-stenosing atherosclerosis of the vessels of the carotid basin.

Thus, violations of peripheral hemodynamics are very typical for patients in the early recovery period of ischemic stroke. They are more pronounced on the side of the paresis. In the early recovery period, more severe forms of microcirculation disorder were observed more often on the paretic limb, accompanied by signs of a sharp decrease in blood flow, increased vascular tone with the phenomena of blood stagnation. The revealed disorders of the microcirculatory link allow the active use of disaggregants, peripheral vasodilators, venotonics, massage, physiotherapy, reflexotherapy [35], therapeutic physical training [36] involving patients and healthy limbs, to be recommended to patients in the early recovery period after ischemic stroke. All this should help improve the rheological properties of red blood cells [37-41] and increase the efficiency of rehabilitation of this category of patients during the early stage of recovery.

Conclusion

In the early recovery period of ischemic stroke, violations of peripheral hemodynamics are noted. They are more pronounced on the side of the paresis. In the early recovery period, more severe forms of microcirculation disorder were observed more often on the paretic

limb, accompanied by signs of a sharp decrease in blood flow, increased vascular tone with the phenomena of blood stagnation. The revealed disorders of the microcirculatory link allow recommending to patients in the early recovery period after ischemic stroke active use of medicamentous and non-medicamentous agents that can improve the rheological properties of red blood cells and improve the efficiency of rehabilitation of this category of patients during the early stage of recovery.

References

1. Zavalishina SY, Kutafina NV, Vatnikov YA, Makurina ON, Kulikov EV, et al. (2016) Platelet-Activity Dependence on the Age of Rats with Experimental Dyslipidemia. *Biol Med* 8: 326.
2. Zavalishina SY, Vatnikov YA, Makurina ON, Kulikov EV, Sotnikova ED, et al. (2017) Diagnostics Appreciation of Physiological Reaction of Intravascular Thrombocytes' Activity of Two-Years-Old Mice to Regular Physical Loads. *Biomed Pharmacol J* 10: 129-136.
3. Medvedev IN, Skorjatina IA, Zavalishina SY (2016) Vascular control over blood cells aggregation in patients with arterial hypertension with dyslipidemia. *Cardiovas Ther Prevention* 15: 4-9.
4. Makhov AS, Medvedev IN, Rysakova OG (2017) Functional features of hemostasis and physical fitness of skilled snowboarders with hearing impairment. *Teoriya i Praktika Fizicheskoy Kultury* 12: 27.
5. Mitrokhina NV, Vatnikov YA, Sotnikova ED, Kulikov EV (2014) Assessment of the risk of osteosarcoma recurrence in canine long bone replants. *Biol Med* 6: 06-14.
6. Kulikov EV, Vatnikov YA, Sotnikova ED, Seleznev SB, Troshina NI, et al. (2015) Morphometric characteristics of the bone tissue structure in white Volga guineafowls. *Biol Med* 7: 111-115.
7. Vatnikov YA, Sakhno NV, Sotnikova ED, Kulikov EV, Parshina VI, et al. (2015) Clinical Control of Packed RBC Transfusion in Acute Surgical Pathology such as Gastric Dilation and Volvulus in Dogs. *Biomed Pharmacol J* 8.
8. Zavalishina SY, Medvedev IN (2016) Features aggregation erythrocytes and platelets in old rats experiencing regular exercise on a treadmill. *Adv Gerontol* 29: 437-441.
9. Krasnikov AV, Annikov VV, Vatnikov YA, Sotnikova ED, Kulikov EV, et al. (2016) Analysis of Dental Implants' Biointegration in Animals. *Biol Med* 8: 178-186.
10. Zavalishina SY, Kutafina NV, Vatnikov YA, Makurina ON, Kulikov EV, et al. (2016) Platelet-Activity Dependence on the Age of Rats with Experimental Dyslipidemia. *Biol Med* 8: 326.
11. Krasnikov AV, Annikov VV, Vatnikov YA, Sotnikova ED, Kulikov EV, et al. (2016) Analysis of Dental Implants' Biointegration in Animals. *Biol Med* 8: 178-186.
12. Bikbulatova AA, Andreeva EG, Medvedev IN (2017) Platelets' Functional Peculiarities in Persons of the Second Mature Age with Spinal Column Osteochondrosis of the Second Degree. *Ann Res Rev Biol* 21: 1-9.
13. Zavalishina SY (2017) Physiological Features of Hemostasis in Newborn Calves Receiving Ferroglyukin, Fosprenil and Hamavit, for Iron Deficiency. *Ann Res Rev Biol* 14: 1-8.
14. Kulikov EV, Seleznev SB, Sotnikova ED, Vatnikov YA, Kharlitskaya EV, et al. (2016) The Morphological Aspects of Bone

- Marrow of Guinea Fowl of the Volga White Breed in Postembryonic Ontogenesis. *RJPBCS* 7: 1148-1153.
15. Kulikov EV, Vatnikov YA, Parshina VI, Sotnikova ED, Vilkovskiy IF, Popova IA, Kochneva MV, Karamyan AS. (2017) Special aspects of the pathohistological diagnostics of familial shar-pei amyloidosis. *Asian J Pharma* 1: S152-S157.
 16. Lenchenko EM, Vatnikov YA, Sotnikova ED, Kulikov EV, Gnezdilova LA, et al. (2017) Experimental toxemia of chickens contaminated with yersinia enterocolitica bacteria. *Asian J Pharma* 11: S91-S96.
 17. Seleznev SB, Kulikov EV, Vetoshkina GA, Vatnikov YA, Sotnikova ED, et al. (2017) The evolution and structural organization of the organs of vertebrate immune system. *Asian J Pharma* 11: S84-S90.
 18. Bikbulatova AA. (2017) Dynamics of Locomotor Apparatus' Indices of Preschoolers with Scoliosis of I-II Degree against the Background of Medicinal Physical Training. *Biomed Pharmacol J* 10.
 19. Zavalishina SY, Medvedev IN (2017) Comparison of opportunities from two therapeutical complexes for correction of vascular hemostasis in hypertensives with metabolic syndrome. *Cardiovas Ther Prevent* 16: 15-21.
 20. Vorobyeva NV (2017) Physiological Reaction of Erythrocytes' Microrheological Properties on Hypodynamia in Persons of the Second Mature Age. *Ann Res Rev Biol* 20: 1-9.
 21. Skoryatina IA, Zavalishina SY, Makurina ON, Mal GS, Gamolina OV (2017) Some aspects of Treatment of Patients having Dislipidemia on the Background of Hypertension. *Prensa Med Argent* 103: 3.
 22. Medvedev IN (2017) The Impact of Durable and Regular Training in Hand-to-hand Fighting Section on Aggregative Platelet Activity of Persons at the First Mature Age. *Ann Res Rev Biol* 15: 1-6.
 23. Medvedev IN (2016) Platelet functional activity in clinically healthy elderly. *Adv Gerontol* 29: 633-638.
 24. Medvedev IN, Gromnatskii NI (2005) The influence of nebulolol on thrombocyte aggregation in patients with arterial hypertension with metabolic syndrome. *Klinicheskaja meditsina* 83: 31-33.
 25. Stakhovskaya LV, Kotov SV (2014) *Stroke. A guide for doctors.* Moscow, Russia.
 26. Suslina ZA, Tanashyan MM, Ionova VG (2005) *Ischemic stroke.* Moscow, Russia.
 27. Haeuber E, Shaughnessy M, Forrester LW (2004) Accelerometer monitoring of home- and community-based ambulatory activity after stroke. *Arch Phys Med Rehabil* 85: 1997-2001.
 28. Kutafina NV (2017) Platelet Parameters of Holstein Newborn Calves. *Ann Res Rev Biol* 15: 1-8.
 29. Medvedev IN (2017) Physiological Dynamics of Platelets' Activity in Aged Rats. *Ann Res Rev Biol* 18: 1-6.
 30. Medvedev IN (2015) Neuropsychological changes in cognitive function during the transition of the elderly in old. *Adv Gerontol* 28: 479-483.
 31. Zavalishina SY. (2017) Restoration of Physiological Activity of Platelets in New-Born Calves With Iron Deficiency. *Biomed Pharmacol J* 10: 711-716.
 32. Medvedev IN (2015) Neuropsychological changes in cognitive function during the transition of the elderly in old. *Adv Gerontol* 28: 479-483.
 33. Medvedev IN, Amelina IV (2010) Evaluation of the relationship between chromosome aberrations and transcription activity of nucleolus organizer regions in indigenous Population of the Kursk Region. *Bull Experi Biol Med* 149: 332-336.
 34. Bikbulatova AA, Andreeva EG (2017) Dynamics of Platelet Activity in 5-6-Year Old Children with Scoliosis against the Background of Daily Medicinal-Prophylactic Clothes' Wearing for Half A Year. *Biomed Pharmacol J* 10.
 35. Krupatkin AI (2012) *Laser Doppler flowmetry microcirculation: a guide for doctors.* Moscow, Russia.
 36. Oshurkova JL, Glagoleva TI (2017) Physiological Activity of Platelet Aggregation in Calves of Vegetable Feeding. *Biomed Pharmacol J* 10: 1395-1400.
 37. Oshurkova JL, Medvedev IN, Fomina LL (2017) Platelets' Aggregative Properties of Irishire Calves in the Phase of Dairy-vegetable Nutrition. *Ann Res Rev Biol* 16: 1-6.
 38. Maksimov VI, Parakhnevich AV, Parakhnevich AA, Glagoleva TI, Kutafina NV (2017) Physiological Reaction of Erythrocytes' Micro Rheological Peculiarities in Milk Fed Piglets after the Negative Impact of the Environment. *Ann Res Rev Biol* 17: 1-8.
 39. Maksimov VI, Parakhnevich AV, Parakhnevich AA, Glagoleva TI, Kutafina NV (2017) Rheological Properties of Erythrocytes of Healthy Piglets during the Transition from Dairy to Vegetable Nutrition. *Ann Res Rev Biol* 16: 1-7.
 40. Maksimov VI, Parakhnevich AV, Parakhnevich AA, Glagoleva TI, Kutafina NV (2017) Erythrocytes' Microrheological Features of Piglets during the Phase of Dairy-vegetable Nutrition after Damage or Common Supercooling. *Ann Res Rev Biol* 16: 1-8.
 41. Maksimov VI, Parakhnevich AV, Parakhnevich AA, Glagoleva TI, Kutafina NV (2017) Physiological Reaction of Erythrocytes' Micro Rheological Features in Newborn Piglets on Unfavourable Environmental Factors. *Ann Res Rev Biol* 16: 1-8.