



Plums as Potential Dietary Agents to Prevent Obesity and Obesity-Related Disorders

Rafat A Siddiqui*

Abstract

Obesity is a serious issue in both developed and developing world. The obese people are often hyperglycemia, dyslipidemia, and hypertensive and therefore, are pre-disposed to type 2 diabetes and cardiovascular diseases. It is generally known that eating fruits and vegetables can prevent body weight gain and can reduce the risk of chronic diseases. Recent research has identified several kinds of fruits and vegetables that are important in reducing inflammation and preventing metabolic syndrome. However, the sugar content of fruits can be problematic for obese patients with diabetes. Plums and prunes (dried plums) are rich in fiber and have low glycemic index. Consumption of plums (prunes) helps growth of beneficial gut bacteria, reduces fats deposit in tissues, improves lipid and glucose metabolism, and reduces inflammation. The consumption of plum or prunes causes modulation of cellular pathways that are deregulated in obesity. In conclusion regular consumption of plums or prunes can potentially be very helpful in preventing and/or managing obesity and obesity-related disorders.

Keywords

Obesity; Plums; Prunes; Gut-bacteria; Lipid metabolism; Diabetes

Obesity results from accumulation of fats in the body to an extent that it creates a serious risk to overall health. Excessive accumulation of fat in adipose, liver, heart, pancreases and other tissues results in a combination of metabolic disorders known as “metabolic syndrome” and characterized by hyperglycemia, dyslipidemia, and hypertension that predisposes obese people to type 2 diabetes and cardiovascular diseases [1-4]. Furthermore, obesity is also linked to several types of cancers [5-7]. According to Center for Disease Control (CDC) more than one-third (36.5%) of U.S. adults have obesity [8]. In 2014 the global economic impact of obesity was estimated to be US \$2.0 trillion or 2.8% of the global gross domestic product [9]. In developing countries, obesity is also becoming an emerging disease due to a shift toward a western life style for consuming processed food rich in fats and carbohydrates [3,4].

Nutrient and physical activity play an important role in the progression of obesity. The CDC reports that proper nutrition and exercise can prevent or reverse obesity [10]. It is generally known that eating fruits and vegetables can prevent body weight gain and can reduce the risk of chronic diseases [11]. Recent research has identified several kinds of fruits and vegetables that are important in reducing

inflammation and preventing metabolic syndrome [12] and this list is growing as new biological activities are discovered for the bioactive compounds present in the natural products. One such fruit is a plum, because the recent experimental evidences indicate that plums may be very beneficial for preventing or treating obesity.

Plum has about 40 species; however, only two species, the European plum (*Prunus domestica*) and the Japanese plum (*Prunus salicina*), are commercially recognized worldwide [13]. The European plum originated near the Caspian Sea about 2000 years ago and was introduced into the USA in the 17th century. The Japanese plum was originated in China but was mostly cultivated and developed in Japan and was introduced into the USA in the late 19th century [14]. Plums are dried for longer storage and often referred as prunes. Dried plums (prunes) are extremely low in fat, rich in macronutrients including both soluble (pectin) and insoluble fibers, oligosaccharides, simple sugars and sugar alcohol (sorbitol), as well as micronutrients like vitamins and minerals [14-16]. Several studies demonstrated that prunes are also rich sources of polyphenolic derivatives including chlorogenic acid, neochlorogenic acid, cryptochlorogenic acid, and oligomeric proanthocyanidin [16,17].

A growing number of studies have documented effects of plums or prunes that directly and indirectly relate to promote healthy metabolism and can be beneficial in controlling obesity. For example, plums or prunes promoted satiety when consumed as snacks; therefore, can limit food intake [18]. Interestingly the satiating effect of dried plums occurred concurrent with suppressed plasma glucose and insulin responses [18]. Furthermore, fibers and oligosaccharides in prunes act as prebiotics (food for microorganisms) and promote the growth of the healthy microorganisms in gut which help in nutrient absorption, removing toxins, improving mucosal immunity, decreasing fat metabolism and reducing pathogen colonization [19]. Plums associated- changes in gut bacteria (*Faecalibacterium*, *Lactobacillus*, and *Bacteroidetes* species) prevented weight gain in obese rats [20]. In human subjects, consumption of prunes essence concentrate also improved the population of beneficial bacteria, especially *Bifidobacterium*, *Lactobacillus* species [21]. It is also suggested that in addition to fibers, chlorogenic acid present in prunes may be metabolized by the intestinal microbiota to produce caffeic acid, which favors the proliferation of *Bifidobacterium* and *Lactobacillus* [20,22]. These bacteria colonization suppressed the intestinal pH, elevated production of short chain fatty acids, and also decreased intestinal cholesterol absorption [19,23,24]. Plums have low glycemic index because colonic fermentation results in a slower appearance of glucose into the blood stream [25].

Stacewicz-Sapuntzakis et al. [15,24] suggested that prunes lower the cholesterol level through enhancing bile acid excretion, and diverts excess cholesterol to bile acid synthesis. Several pre-clinical and clinical studies suggested that consumption of prunes extracts or concentrate suppressed the cholesterol level as well as lowered the LDL-oxidation [15,24,26]. Plum juice consumption decreased total cholesterol and triglycerides in plasma and significantly increased the ratio HDL-cholesterol/T-cholesterol in obese Zucker rats [27]. These study suggest that consumption of plums (prunes) may decrease the risk of obesity-induced CVD, because of its antioxidant activity [15,24,26,28]. An earlier study showed that plum treatment

*Corresponding author: Rafat A. Siddiqui, Nutrition Science and Food Chemistry Laboratory, Agriculture Research Station, Virginia State University, Petersburg, VA 23804, USA, Tel: (804) 524 5957; E-mail: rsiddiqui@vsu.edu

Received: May 02, 2017 Accepted: May 03, 2017 Published: May 06, 2017

for 2 weeks reduced areas under the curve (AUCs) for glucose and insulin during a glucose tolerance test in db/db mice [29]. Plum treatment significantly increased plasma adiponectin concentrations and PPAR γ mRNA expression in adipose tissue from Wistar fatty rats [29]. Plum thus may increase insulin sensitivity in these rats via adiponectin-related mechanisms. Plums juice also decreased the leptin levels in plasma of rats [27]. Furthermore, plum juice consumption decreased fat deposits in liver and heart tissues, inhibited adhesion molecules (ICAM-1 and VCAM-1), prevented NF- κ B activation, and decreased biomarkers of inflammation and hypertension [26]. These effects were suggested to be mediated through PPAR- γ -agonist activity. Activation of the PPAR- γ in adipose tissue is associated with regulation of adipose derived substances that are altered in obesity and insulin resistance [30]. Another study showed that Queen Garnet plum (GGP), a variety of the Japanese plum, improved cardiovascular and hepatic structure and function and reduced metabolic parameters such as bodyweight gain, visceral adiposity index and total body fat mass induced by the high fat diet [31]. The consumption of QCP also improved plasma lipid profiles by reducing plasma concentrations of triglycerides, total cholesterol and non-esterified fatty acids. Furthermore, the reduction in adipose tissue with QGP was correlated with decreased plasma leptin concentrations. Leptin is a hormone secreted from adipose tissue [32] and a leptin deficiency was found to promote weight regain by stimulating appetite and reducing energy output [33]. Population studies have also indicated that approximately 10% of obese individuals have low plasma leptin levels [34]. Leptin is a critical factor that link reduced energy stores to eating behavior. It is suggested that leptin therapy could sustain weight loss by overriding the tendency toward energy conservation [35].

The studies described above clearly show the beneficial effects of plums or prunes help to attenuate obesity and obesity-related disorders. It is evident that consumption of plums or prunes has resulted in modulation of cellular pathways that are deregulated in obesity. Consumption of prunes can also be very helpful for diabetic and CVD patients. Further studies are required in this area to understand the cellular and molecular mechanism for the beneficial effects of prunes on human health that are not yet commonly known. In conclusion regular consumption of plums or prunes, a low glycemic fruits, can potentially be very helpful in preventing and/or managing obesity and obesity-related disorders.

References

1. Ogden CL, Yanovski SZ, Carroll MD, Flegal KM (2007) The epidemiology of obesity. *Gastroenterology* 132: 2087-2102.
2. Gallagher EJ, Leroith D, Karnieli E (2010) Insulin resistance in obesity as the underlying cause for the metabolic syndrome. *Mt Sinai J Med* 77: 511-523.
3. Ford ES, Giles WH, Dietz WH (2002) Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 287: 356-359.
4. Ogunbode AM, Ladipo M, Ajayi IO, Fatiregun AA (2011) Obesity: an emerging disease. *Niger J Clin Pract* 14: 390-394.
5. Carroll K (1995) Obesity as a risk factor for certain types of cancer. *Lipids* 33: 1055-1059.
6. Bergstrom A, Pisani P, Tenet V, Wolk A, Adami HO (2001) Overweight as an avoidable cause of cancer in Europe. *Int J Cancer* 91: 421-430.
7. Peto J (2001) Cancer epidemiology in the last century and the next decade. *Nature* 411: 390-395.
8. Ogden CL, Carroll MD, Fryar CD, Flegal KM (2015) Prevalence of obesity among adults and youth. *CHS Data Brief* 219: 1-8.

9. Dobbs R, Sawers C, Thompson F, Manyika J, Woetzel JR, et al. (2014) *Overcoming Obesity: An Initial Economic Analysis*; McKinsey Global Institute: Jakarta, Indonesia.
10. Khan LK, Sobush K, Keener D, Godman K, Lowry A, et al. (2009) Centers for Disease Control and Prevention Recommended Community Strategies and Measurements to Prevent Obesity in the United States. *MMWR* 58: 1-26.
11. Boeing H, Bechthold A, Bub A, Ellinger S, Haller D, et al. (2012) Critical review: vegetables and fruit in the prevention of chronic diseases. *Eur J Nutr* 51: 637-663.
12. Esmailzadeh A, Kimiagar M, Mehrabi Y, Azadbakht L, Hu FB, et al. (2006) Fruit and vegetable intakes, C-reactive protein, and the metabolic syndrome. *Am J Clin Nutr* 84: 1489-1497.
13. Topp BL, Russell DM, Neumüller M, Dalbó MA, Liu W (2012) *Plum*. In *Fruit Breeding*. Springer: New York.
14. Igwe EO, Charlton KE (2016) A Systematic Review on the Health Effects of Plums (*Prunus domestica* and *Prunus salicina*). *Phytother Res* 30: 701-73.
15. Stacewicz-Sapuntzakis M, Bowen PE, Hussain EA, Damayanti-Wood BI, Farnsworth NR (2001) Chemical composition and potential health effects of prunes: a functional food? *Crit Rev Food Sci Nutr* 41: 251-286.
16. Reele SB, Chodos DJ (1985) Sorbitol induced diarrheal illness model. *Int J Clin Pharmacol Ther Toxicol* 23: 403-405.
17. Chok G, Lang K (1961) Action of chlorogenic acid in the gastrointestinal tract. *Arzneim-Forsch* 11: 545-549.
18. Furchner-Evanson A, Petrisko Y, Howarth L, Nemosek T, Kern M (2010) Type of snack influences satiety responses in adult women. *Appetite* 54: 564-569.
19. Alonso VR, Guarner F (2013) Linking the gut microbiota to human health. *Br J Nutr* 109: 21-26.
20. Noratto GD, Garcia-Mazcorro JF, Markel M, Martino HS, Minamoto Y, et al. (2014) Carbohydrate free peach (*Prunus persica*) and plum (*Prunus domestica*) juice affects fecal microbial ecology in an obese animal model. *PLoS One* 9: 101723.
21. Chiu HF, Huang YC, Lu YY, Han YC, Shen YC, et al. (2017) Regulatory/modulatory effect of prune essence concentrate on intestinal function and blood lipids *Pharma Biol* 55: 974-979.
22. Parkar SG, Trower TM, Stevenson DE (2013) Fecal microbial metabolism of polyphenols and its effects on human gut microbiota. *Anaerobe* 23: 12-19.
23. Fujimura KE, Slusher NA, Cabana MD, Lynch SV (2010) Role of the gut microbiota in defining human health. *Expert Rev Anti Infect Ther* 8: 435-454.
24. Stacewicz-Sapuntzakis M (2013) Dried plums and their products: composition and health effects-an updated review. *Crit Rev Food Sci Nutr* 53: 1277-1302.
25. Nilsson AC, Ostman EM, Holst JJ, Bjorck IM (2008) Including indigestible carbohydrates in the evening meal of healthy subjects improves glucose tolerance, lowers inflammatory markers, and increases satiety after a subsequent standardized breakfast. *J Nutr* 138: 732-739.
26. Tinker LF, Schneeman BO, Davis PA, Gallaher DD, Waggoner CR (1991) Consumption of prunes as a source of dietary fiber in men with mild hypercholesterolemia. *Am J Clin Nutr* 53: 1259-1265.
27. Noratto G, Martinob HSD, Simboc S, Byrned D, Mertens-Talcotte SU (2015) Consumption of polyphenol-rich peach and plum juice prevents risk factors for obesity-related metabolic disorders and cardiovascular disease in Zucker rats. *J Nutr Biochem* 26: 633-641.
28. Gallaher CM, Gallaher DD (2009) Dried plums (prunes) reduce atherosclerosis lesion area in apolipoprotein E-deficient mice. *Br J Nutr* 101: 233-239.
29. Utsunomiya H, Yamakawa T, Kamei J, Kadonosono K, Tanaka S (2005) Anti-hyperglycemic effects of plum in a rat model of obesity and type 2 diabetes Wistar fatty rat. *Biomed Res* 26: 193-200.
30. Sharma AM, Staels B (2007) Review: peroxisome proliferator-activated receptor gamma and adipose tissue—understanding obesity-related changes in regulation of lipid and glucose metabolism. *J Clin Endocrinol Metab* 92: 386-395.

31. Bhaswanta M, Fanning K, Netzel M, Mathai ML, Panchal SK, et al. (2015) Cyanidin 3-glucoside improves diet-induced metabolic syndrome in rats. *Pharmacol Res* 102: 208-217.

32. Havel PJ (2000) Role of adipose tissue in body-weight regulation: mechanisms regulating leptin production and energy balance. *Proc Nutr Soc* 59: 359-371.

33. Rosenbaum M, Murphy EM, Heymsfield SB, Matthews DE, Leibel RL (2002) Low dose leptin administration reverses effects of sustained weight-reduction on energy expenditure and circulating concentrations of thyroid hormones. *J Clin Endocrinol Metab* 87: 2391-2394.

34. Flier JS (1998) Clinical review 94: what's in a name? In search of leptin's physiologic role. *J Clin Endocrinol Metab* 83: 1407-1413.

35. Ahima RS (2008) Revisiting leptin's role in obesity and weight loss. *J Clin Invest* 118: 2380-2383.

Author Affiliations

[Top](#)

Nutrition Science and Food Chemistry Laboratory, Virginia State University, USA

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • www.scitechnol.com/submission