



Polychlorinated Biphenyl Exposure, Reproductive and Genital Abnormalities: Still a Problem for the World and Wildlife

Pinar E^{1*}, Kocer-Gumusel B¹, Elnour A² and Bhagavathula AS³

Introduction

United States Environmental Protection Agency (EPA) defined an endocrine-disrupting chemical (EDC) as “an exogenous agent that interferes with synthesis, secretion, transport, metabolism, binding action, or elimination of natural blood-borne hormones that are present in the body and are responsible for homeostasis, reproduction, and developmental process” [1]. These chemicals were originally thought to exert actions primarily through different hormone receptors, including estrogen receptors (ERs) and androgen receptors (ARs) [1]. However, basic scientific research shows that the mechanisms are much broader than originally defined.

The group of molecules identified as EDCs belongs to various chemical groups and have very different chemical structures. They can be both synthetic and natural. The synthetic chemicals include several industrial solvents/lubricants and their byproducts [polychlorinated biphenyls (PCBs), polybrominated biphenyls (PBBs), dioxins], plastics [bisphenol A (BPA), phthalates], pesticides [methoxychlor, chlorpyrifos, dichlorodiphenyltrichloroethane (DDT)], fungicides (vinclozolin), and pharmaceutical agents [diethylstilbestrol (DES)] [1,2].

Discussion

Polychlorinated biphenyls (PCBs) are a class of synthetic chlorinated aromatic compounds, used in cutting oils, lubricants, and as electrical insulators. They were widely used in industrial and consumer products for decades. Their production started in 1929 and was banned in the late 1970s [3]. The general population is exposed primarily through the consumption of contaminated foods like fish, meat, and dairy products and they can bio-accumulate up the food chain. As they are lipophilic, they are biologically concentrated and stored in human and animal adipose tissues. As a result of their persistence and ubiquity, measurable levels of serum PCBs are found in the majority of the general population [4].

The effects of PCBs on reproduction and hypothalamus-pituitary-gonadal (HPG) axis are shown in many rodent studies. In male rodents, phthalates and to a lesser extent PCBs are shown to induce

testicular dysgenesis syndrome (TDS) entirely or partly. Among wildlife, polar bears (*Ursusmaritimus*) showed waned testosterone production in males, elevated progesterone in females, altered behavior and thyroid hormone levels [5,6]. In humans, many PCBs have estrogenic or anti-androgenic activity. Although estrogenic activity of these compounds is a suspected mode of action, there is also evidence that PCBs inhibit estrogen sulfotransferase activity in the liver and effectively increase bioavailable estrogen in the body [7]. PCBs are also shown to be related to reproductive dysfunction (decreased sperm motility, decrease in fecundity, earlier menarche, altered sex ratio, and altered gonadal hormones in newborns) [8-11]. Several epidemiological studies suggest an inverse association between PCB concentrations in biological fluids and semen quality. High PCB levels in humans are specifically suggested to cause reduced sperm motility. These results were constantly reported from different parts of the world including the United States, The Netherlands, Taiwan, Sweden, and India [12-14]. Studies suggest that consumption of contaminated fish or exposure to contaminated rice oil could cause high PCB concentrations in humans [15,16]. However, there is limited epidemiological evidence that reveal the association between maternal serum concentrations of PCBs and cryptorchidism or hypospadias in the offspring [17-22]. Prenatally exposed infants to PCBs showed lower birth weight, smaller head circumference and alterations in the thyroid hormone homeostasis [23], while exposed children showed altered neural development, cognitive, motor and learning abilities [24,25].

PCBs were also shown to cause perturbations in the prostate gland. A recent study was conducted on Swedish men who did and did not have prostate cancer revealed that their PCB concentrations (specifically in the higher PCB level quadrants) of adipose tissue showed a significant correlation with prostate cancer odds ratio, with the most marked associations for PCB 153 and transchlordan [26]. The results of another important epidemiological study conducted on capacitor manufacturing plant workers exposed to high levels of PCBs also indicated a strong exposure-response correlation for prostate cancer mortality [27]. These results support previous findings of correlations between PCB 153 and PCB 180 and prostate cancer risk in electric utility workers [28,29].

Different PCB congeners have been banned more than 35 years ago; however they are still present in the environment and biological fluids. This phenomenon indicates their persistence in the environment as well as illegal usage. Although there is enough evidence that these compounds cause reproductive, genital and neurobehavioral anomalies in rodents, wildlife and humans, there are still gaps in knowledge for their mode/mechanism of action/s.

Conclusion

Further investigation using *in vitro* and *in vivo* models should be conducted to reveal the mechanism underlying their reproductive and developmental toxicity. Scientific societies should work together to increase the awareness to the hazard of EDCs, particularly PCBs. As toxicologists and pharmacists, we suggest that regulatory authorities take strict measures to decrease human exposure to different PCB congeners.

*Corresponding author: Pinar Erkekoglu, Faculty of Pharmacy, Department of Toxicology, Hacettepe University, 06100, Ankara, Turkey, Tel: +90 312 305 21 78; Fax: +90 312 311 47 77; E-mail: erkekp@yahoo.com

Received: November 04, 2015 Accepted: November 07, 2015 Published: January 14, 2016

References

1. Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, et al. (2009) Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocr Rev* 30: 293-342.
2. Maqbool F, Mostafalou S, Bahadar H, Abdollahi M (2015) Review of endocrine disorders associated with environmental toxicants and possible involved mechanisms. *Life Sci* S0024-3205: 30048-30045.
3. Thomas K, Xue J, Williams R, Jones P, Whitake D (2012) Polychlorinated Biphenyls (PCBs) in School Buildings: Sources, Environmental Levels, and Exposures EPA/600/R-12/051.
4. Longnecker MP, Rogan WJ, Lucier G (1997) The human health effects of DDT (dichlorodiphenyltrichloroethane) and PCBs (polychlorinated biphenyls) and an overview of organochlorines in public health. *Annu Rev Public Health* 18: 211-244.
5. Lie E, Larsen HJS, Larsen S, Johansen GM, Derocher AE, et al. (2005) Does high organochlorine (OC) exposure impair the resistance to infection in polar bears (*Ursus maritimus*)? Part I: Effect of OCs on the humoral immunity. *J Toxicol Environ. Health A* 67: 555-582.
6. Sonne C (2010) Health effects from long-range transported contaminants in Arctic top predators. An integrated review based on studies of polar bears and relevant model species. *Environ Int* 36: 461-491.
7. Kester MH, Bulduk S, van Toor H, Tibboel D, Meinel W, Glatt H, et al. (2002) Potent inhibition of estrogen sulfotransferase by hydroxylated metabolites of polyhalogenated aromatic hydrocarbons reveals alternative mechanism for estrogenic activity of endocrine disrupters. *J Clin Endocrinol Metab* 87: 1142-1150.
8. Meeker JD, Hauser R (2010) Exposure to polychlorinated biphenyls (PCBs) and male reproduction. *Syst Biol Reprod Med* 56: 122-131.
9. Faroon OM, Keith S, Jones D, de Rosa C (2001) Effects of polychlorinated biphenyls on development and reproduction. *Toxicol Ind Health* 17: 63-93.
10. Schell LM, Gallo MV (2010) Relationships of putative endocrine disruptors to human sexual maturation and thyroid activity in youth. *Physiol Behav* 99: 246-253.
11. Karmaus W, Huang S, Cameron L (2002) Parental Concentration of Dichlorodiphenyl Dichloroethane and Polychlorinated Biphenyls in Michigan Fish Eaters and Sex Ratio in Offspring. *J Occup Environ Med* 44: 08-13.
12. Cao Y, Winneke G, Wilhelm M, Wittsiepe J, Lemm F, et al. (2008) Environmental exposure to dioxins and polychlorinated biphenyls reduce levels of gonadal hormones in newborns: results from the Duisburg cohort study. *Int J Hyg Environ Health* 211: 30-39.
13. Dallinga JW, Moonen EJ, Dumoulin JC, Evers JL, Geraedts JP, Kleinjans JC (2002) Decreased human semen quality and organochlorine compounds in blood. *Hum Reprod* 17: 1973-1979.
14. Richthoff J, Rylander L, Jönsson BA, Akesson H, Hagmar L, et al. (2003) Serum levels of 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) in relation to markers of reproductive function in young males from the general Swedish population. *Environ Health Perspect* 111: 409-413.
15. Hauser R, Chen Z, Pothier L, Ryan L, Altshul L (2003) The relationship between human semen parameters and environmental exposure to polychlorinated biphenyls and p,p'-DDE. *Environ Health Perspect* 111: 1505-1511.
16. Rignell-Hydbom A, Rylander L, Giwercman A, Jönsson BA, Nilsson-Ehle P, et al. (2004) Exposure to CB-153 and p,p'-DDE and male reproductive function. *Hum Reprod* 19: 2066-2075.
17. Guo YL, Hsu PC, Hsu CC, Lambert GH (2000) Semen quality after prenatal exposure to polychlorinated biphenyls and dibenzofurans. *Lancet* 356: 1240-1241.
18. Hsu PC, Huang W, Yao WJ, Wu MH, Guo YL, et al. (2003) Sperm changes in men exposed to polychlorinated biphenyls and dibenzofurans. *JAMA* 289: 2943-2944.
19. Longnecker MP, Klebanoff MA, Brock JW, Zhou H, Gray KA, et al. (2002) Maternal serum level of 1,1-dichloro-2,2-bis(p-chlorophenyl)ethylene and risk of cryptorchidism, hypospadias, and polythelia among male offspring. *Am J Epidemiol* 155: 313-322.
20. Bhatia R, Shiau R, Petreas M, Weintraub JM, Farhang L, et al. (2005) Organochlorine pesticides and male genital anomalies in the child health and development studies. *Environ Health Perspect* 113: 220-224.
21. Mol NM, Sørensen N, Weihe P, Andersson AM, Jørgensen N, et al. (2002) Spermatid and serum hormone concentrations at the age of puberty in boys prenatally exposed to polychlorinated biphenyls. *Eur J Endocrinol* 146: 357-363.
22. Hosie S, Loff S, Witt K, Niessen K, Waag KL (2000) Is there a correlation between organochlorine compounds and undescended testes? *Eur J Pediatr Surg* 10: 304-309.
23. Sandau C, Ayotte P, Dewailly E, Duffe J, Norstrom RJ (2002) Pentachlorophenol and hydroxylated polychlorinated biphenyl metabolites in umbilical cord plasma of neonates from coastal population of Quebec. *Environ Health Perspect* 110: 411-417.
24. Roze E, Meijer L, Bakker A, Van Braeckel K, Sauer PJ, et al. (2009) Prenatal exposure to organohalogenated flame retardants, influences motor, cognitive and behavior performance at school age. *Environ Health Perspect* 117: 1953-1958.
25. Park HY, Park JS, Sovcikova EJ, Kocan A, Linderholm L, et al. (2009) Exposure to hydroxylated polychlorinated biphenyls (OH-PCBs) in the prenatal period and subsequent neurodevelopment in eastern Slovakia. *Environ Health Perspect* 117: 1600-1606.
26. Hardell L, Andersson SO, Carlberg M, Bohr L, van Bavel B, Lindström G, et al. (2006) Adipose tissue concentrations of persistent organic pollutants and the risk of prostate cancer. *J Occup Environ Med* 48: 700-707.
27. Prince MM, Ruder AM, Hein MJ, Waters MA, Whelan EA, et al. (2006) Mortality and exposure response among 14,458 electrical capacitor manufacturing workers exposed to polychlorinated biphenyls (PCBs). *Environ Health Perspect* 114: 1508-1514.
28. Ritchie JM, Vial SL, Fuortes LJ, Guo H, Reedy VE, et al. (2003) Organochlorines and risk of prostate cancer. *J Occup Environ Med* 45: 692-702.
29. Charles LE, Loomis D, Shy CM, Newman B, Millikan R, et al. (2003) Electromagnetic fields, polychlorinated biphenyls, and prostate cancer mortality in electric utility workers. *Am J Epidemiol* 157: 683-691.

Author Affiliation

Top

¹Hacettepe University, Faculty of Pharmacy, Turkey

²Department of Clinical Pharmacy, University of Gondar, College of Medicine and Health Sciences, School of Pharmacy, Gondar, Ethiopia

³Department of Pharmacology, College of Medicine and Health Sciences, UAE University, AlAin, UAE

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 50 Journals
- ❖ 21 Day rapid review process
- ❖ 1000 Editorial team
- ❖ 2 Million readers
- ❖ Publication immediately after acceptance
- ❖ Quality and quick editorial, review processing

Submit your next manuscript at ● www.scitechnol.com/submission

This article is published in the special issue **Testicular Dysgenesis Syndrome: Past, Present and Future** and has been edited by Dr. Lawrence S. Amesse, Wright State University Boonshoft School of Medicine, USA