

Reticular erythematous mucinosis, A rare cutaneous mucinosis

Amer Almohssen

American Society of Dermatologic Surgery, USA



Abstract

Abstract Reticular erythematous mucinosis (REM) is a rare form of primary cutaneous mucinosis, most often involving sun exposed skin of the midline of the upper chest or back in middle-aged women. REM bears clinical and histopathologic resemblance to lupus erythematosus tumidus (LET), dermatomyositis, scleredema, and lichen myxedematosus. Early recognition and diagnosis of REM is particularly relevant as REM is more benign and has fewer systemic consequences.

Biography

Amer Almohssen has completed his Dermatology residency training Rutgers University in New Jersey, USA, completed his Fellowship training in Dermatopathology at the State University of New York and The Ackerman Academy of Dermatopathology in New York City, USA. He is an Attending in Dermatology and has many publications that have been cited numerous times.

Publications

Almohssen AA, Schwartz RA., J Eur Acad Dermatol Venereol. 2019 Mar;33(3):e103-e104

Weiss A, Shah R, Almohssen AA, Lambert WC, Skinmed. 2020 Mar 1;18(2):109-111. eCollection 2020

Global Summit on Dermatology and Cosmetology, London, UK | 8th-9th June, 2020

Citation: Amer Almohssen, *R Reticular erythematous mucinosis, A rare cutaneous mucinosis*, Dermatology Summit 2020, Global Summit on Dermatology and Cosmetology, London, UK, June 8-9, 2020, 03:02-3