



Research Article

A SCITECHNOL JOURNAL

Erosive Tooth Wear and its Risk Factors among 9-14 Years Children Undergoing Orthodontics

Gupta A^{1*}, Anur G², Singh K³, Singh S⁴, Jossan AS⁵ and Singh A⁵

Abstract

Objective: The purpose of this study was to assess the prevalence of dental erosion and its associated risk factors among 9-14 year-old children with orthodontic braces.

Methods: This cross-sectional survey was performed by calibrated examiners in a random sample of 447 patients (250 - boys and 197 - girls). Clinical examination was done for dental erosion using the diagnostic criteria of Eccles and the Index of O'Sullivan to record the distribution of the dental erosion. A self-administered questionnaire was developed to record information about consumption of drinks. The analyses were performed using SPSS software 16.

Results: Overall significant difference in the prevalence of dental erosion was seen in all the factors except age group. The most frequently affected surface was the incisal or occlusal edge (43.2%). The results showed that males, children who consumed carbonated drinks, fruit juice and those who had habit of swimming had more dental erosion.

Conclusions: Dental erosion is a major public health problem especially among children with orthodontic cases. The prevalence of erosion is commonly associated with carbonated drinks and swimming. So, orthodontists should offer preventive care and promote a healthier lifestyle by education to children, who are at increased risk of dental erosion.

Keywords

Beverages; Children; Dental erosion; Orthodontists

Introduction

Non-carious lesions are becoming a progressively significant problem in oral health. Dental erosion is the one of the commonest non carious lesion affecting mainly child population. It is defined as the chronic irreversible chemical dissolution of the dental hard tissues caused by acids or chelants without bacterial involvement [1]. It has important risk factors including both extrinsic and intrinsic aspects. Intrinsic factors include ailment that cause vomiting or regurgitation such as gastro-oesophageal reflux and, anorexia nervosa [2] that cause a reduction in saliva flow. While the extrinsic ones include dietary habits such as acidic drinks and

foods, sports like swimming in chlorinated pools and occupational factors [3]. Many surveys have shown upto 30% of erosion among 12 year old children and upto 53% among 5 year old children [3,4]. According to Xhonga and Valdmanis, Lussi et al. Smith and Robb, the prevalence of dental erosion ranges between 25% and 60% in child population. The most frequent origin of erosion is due to acidic foods as orange, lemon juices and drinks as colas, pepsi [2], as these drinks turn down the pH of oral cavity below 5.0–5.7. This reduction in pH cause demineralization of enamel and dentin which show the way to severe loss of tooth surface and ultimately results in sensitivity [5]. It is observed that frequent intakes of acidic juices are greater factor for dental erosion. Patients suffering from Sjogren's syndrome are predominantly at higher risk of developing erosion since reduction in salivary flow results in decreased ability to balance the acidic environment of mouth therefore prevents remineralization [6].

Studies done by Williams et al. Milosevic et al. and Auad et al. have described the distribution, prevalence and severity of erosion in adolescents but literature is lacking in children especially undergoing orthodontic treatment [7]. Hence, this study aimed to determine the prevalence and severity of erosion in children aged 9-14 years.

Methodology

A descriptive cross sectional survey was conducted among 447 children with braces aged between 9 – 14 years in Jammu, India. Ethical clearance was obtained from ethical committee of the Institute and informed consent was taken before starting the examination. All the available children undergoing orthodontic appliances of this age group during study schedule were selected and those who were on medication, having symptoms of bruxism, reflux and Sjogren's syndrome were excluded.

Calibration of examiners

Training and calibration of the examiner was done in the department before onset of the study. The diagnostic criteria of dental erosion were appropriately discussed. The results of the duplicate examinations were utilized for estimation of intra-examiner reliability.

Clinical examination

The clinical examination was done in a well illuminated classroom using mouth mirrors and sterilized cotton rolls. The teeth examined were central incisors, lateral incisors, and first molars in both the arches. The diagnostic criterion for measuring dental erosion proposed by Eccles was used [8]. The index of O'Sullivan was adopted to record the distribution of affected teeth [9].

O'sullivan index for measurement of dental erosion:

Site on erosion on each tooth

Code A: Labial or buccal only

Code B: Lingual or palatal only

Code C: Occlusal or incisal only

Code E: Multi-surface

*Corresponding author: Akshay Gupta, Department of Orthodontics and Dentofacial Orthopaedics, Indira Gandhi Government Dental College and Hospital, Jammu, India, Tel: +918437112344; E-mail: drkarankahlon@gmail.com

Received: May 04, 2017 Accepted: July 05, 2017 Published: July 12, 2017

Questionnaire

A self prepared questionnaire was used for recording demographic details, hobby of swimming, fruits and beverages intake.

Data analysis

Data was analyzed using SPSS version 15.0. Student's t test was used to find the significance of difference in the means of dental erosion at p value <0.05. To test the association between independent sample proportions, chi-square test was done.

Results

The total number of the sample who was examined was 447 including 250 - males and 197 - females. Further they were categorized according to age, habit of swimming, consumption of fruit juices and carbonated drinks.

In the present study, prevalence rate of dental erosion among all participants was 31.5% in orthodontic cases. According to gender a significant difference in the mean scores was obtained, where male children having more erosion compared to their counterparts **Table 1**. Similarly children in the older age group and with regular habit of swimming were having higher level of dental erosion **Tables 2** and **3**. The study subjects found with more intake of fruit juices and carbonated drinks, had comparatively more erosion at p=0.000 **Table 4**. Overall significant difference in the frequency of dental erosion was seen in all the factors except age group as shown in **Table 5**. According to the surfaces affected, most commonly tooth surfaces involved were incisal/occlusion (52.3%), followed by the labial/ buccal surfaces (26.5%), whereas in 9.6% of the cases, multi-surfaces were involved as shown in **Figure 1**.

Discussion

The overall prevalence of dental erosion in the present study was 31.5%, which was higher than the results of Caglar et al. in 2005 among school children in an Istanbul and Wang et al. in 2010 among 12-13-year-old children in Southern China [10,11]. But this prevalence is lower according to studies done by Deery et al. and Al-Majed et al. this difference could be due the present of orthodontic appliances [12,13].

The most commonly affected surfaces were incisal or occlusal (52.3%), followed by labial or buccal surfaces (26.5%). The application of etchants on the labial or buccal surface causes erosion. Similar results were obtained by Al-Malik et al. study that showed 91% of dental erosion on the incisal/occlusal surfaces of 12-14-year-old Saudi boys [14]. However, Auad et al. and Mangueira et al. reported that the palatal surfaces were the most affected [7,15].

The consumption of carbonated drinks showed positively correlation with dental erosion on the facial and occlusal surfaces, whereas severe palatal erosion occurred infrequently due to chronic vomiting [16]. The major affected surfaces of erosion are incisal/occlusal as these surfaces being more prone to physical forces from mastication, that leads to more chances of dental erosion.

A significant relationship was seen between the consumption of fruit juices and carbonated drinks in association with dental erosion. The results of the present study reported are similar with the previous literature [10,11,17-19]. However study conducted by Arnadóttir et al. in among Icelandic teenagers did not show any correlation [20]. Owens also showed carbonated drinks as cola have strong association

with tooth erosion, as it has a lot of carbonic acids which can rapidly dissolve the enamel surface of teeth [21].

It was also found that erosion was significantly higher in those who used to swim regularly and the findings were similar to study conducted by Talebi et al. among 12-year-old School Children in Mashhad [22]. The proposed pH of swimming pools is 2.7 which lead to acidic environment of oral cavity and results in erosion of teeth.

The present findings showed that male children had significantly more erosion than their counterparts. Similar results were obtained by Al Dlaigan et al. in Birmingham, UK, where male's participants had more buccal/ labial and lingual/palatal tooth surface erosion, and similarly Dugmore et al. in a cohort of adolescents of different ethnics in UK also showed same results [17,23]. However there was no difference in the prevalence of dental erosion between males and females in study by Deery et al. in the UK. The lack of significant differences in the prevalence of erosion according to gender in this study could be due to similar patterns of exposure to the risk factors in this community [12].

Conclusion

The present paper stated that dental erosion is one of the most important problems among children with orthodontic appliances

Table 1: Dental erosion according to gender using student's t test.

Sex	No	Mean	SD	p-value
Male	250	.39	.490	
Female	197	.22	.421	.000*

Table 2: Dental erosion according to age groups using student's t test.

Age	No	Mean	SD	p-value
9-11 years	240	.29	.458	
12-14 years	207	.35	.478	.007*

Table 3: Dental erosion according to swimming using student's t test.

Swimming	No	Mean	SD	p-value
Yes	161	.55	.500	
No	286	.18	.392	.000*

Table 4: Dental erosion according to consumption of fruit juice/ carbonated drinks using student's t test.

Fruit juice / Carbonated drinks	No	Mean	SD	p-value
≥5 times in a week	124	.56	.498	
Occasionally	323	.22	.417	.000*

Table 5: Percentage of children having dental erosion according to different factors.

Characteristics	Dental erosion (%)	p-value
Sex		
Male	39.3%	
Female	22.7%	0.006*
Age		
9-11 years	29.6%	
12-14 years	34.8%	0.231**
Fruit juices/ Carbonated drinks		
≥5 times in a week	57.1%	
Occasionally	22.2%	0.000*
Swimming		
Yes	55.6%	
No	18.8%	0.000*




Figure 1: Showing areas affected with dental erosion.

as they have more prone to habits of consuming carbonated drinks. Also children with habit of swimming tended to have more erosion. Dentists should pay more attention in identification of this lesion at its primary stage. And make an approach of preventive services and promote healthier lifestyles who are at increased risk of dental erosion.

References

1. Gandara BK, Truelove EL (1999) Diagnosis and management of dental erosion. *J Contemp Dent Pract* 1: 16-23.
2. Lussi A (2006) Erosive tooth wear a multifactorial condition of growing concern and increasing knowledge. *Monogr Oral Sci* 20: 1-8.
3. Litonjua LA, Andreana S, Bush PJ, Cohen RE (2003) Tooth wear: attrition, erosion and abrasion. *Quintessence Int* 34: 435-446.
4. Office for national statistics. Children dental health in United Kingdom 2003, London 2005.
5. Lussi A, Schaffner M, Jaeggi T (2007) Dental erosion - diagnosis and prevention in children and adults. *Int Dent J* 57: 385-398.
6. Kitchens M, Owens BM (2007) Effect of carbonated beverages, coffee, sports and high energy drinks and bottled water on the in vitro erosion characteristics of dental enamel. *J Clin Paediatr Dent* 31: 153-159.
7. Auad SM, Waterhouse PJ, Nunn J, Steen N, Moynihan PJ (2007) Dental erosion amongst 13- and 14-year old Brazilian school children. *Int Dent J* 57: 161-167.
8. Eccles JD (1979) Dental erosion of nonindustrial origin. A clinical survey and classification. *J Prosthet Dent* 42: 649-653.
9. O'Sullivan EA (2000) A new index for measurement of erosion in children. *Eur J Paediatr Dent* 2: 69-74.
10. Caglar E, Kargul B, Tamboga I, Lussi A (2005) Dental erosion among children in an Istanbul public school. *J Dent Child* 72: 5-9.
11. Wang P, Lin HC, Chen JH, Liang HY (2010) The prevalence of dental erosion and associated risk factors in 12-13-year-old school children in Southern China. *BMC Public Health* 10: 2-9.
12. Deery C, Wagner ML, Longbottom C, Simon R, Nugent ZJ (2000) The prevalence of dental erosion in a United States and a United Kingdom sample of adolescents. *Pediatr Dent* 22: 505-510.
13. Al-Majed I, Maguire A, Murray JJ (2002) Risk factors for dental erosion in 5-6 year old and 12-14 year old boys in Saudi Arabia. *Community Dent Oral Epidemiol* 30: 38-46.
14. Al-Malik MI, Holt RD, Bedi R (2001) The relationship between erosion, caries and rampant caries and dietary habits in preschool children in Saudi Arabia. *Int J Paediatr Dent* 11: 430-439.
15. Mangueira DF, Sampaio FC, Oliveira AF (2009) Association between socioeconomic factors and dental erosion in Brazilian schoolchildren. *J Public Health Dent* 69: 254-259.
16. Lussi A, Schaffner M, Hotz P, Suter P (1991) Dental erosion in a population of Swiss adults. *Community Dent Oral Epidemiol* 19: 286-290.
17. Al-Dlaigan YH, Shaw L, Smith A (2001) Dental erosion in a group of British 14-year-old, school children. Part I: Prevalence and influence of differing socioeconomic backgrounds. *Br Dent J* 190: 145-149.
18. El Aidi H, Bronkhorst EM, Huysmans MC, Truin GJ (2011) Multifactorial analysis of factors associated with the incidence and progression of erosive tooth wear. *Caries Res* 45: 303-312.
19. Mungia R, Zarzabal LA, Dang SC, Baez M, Stookey GK, Brown JP (2009) Epidemiologic survey of erosive tooth wear in San Antonio, Texas. *Tex Dent J* 126: 1097-109.
20. Arnadóttir IB, Saemundsson SR, Holbrook WP (2003) Dental erosion in Icelandic teenagers in relation to dietary and lifestyle factors. *Acta Odontol Scand* 61: 25-28.
21. Owens BM (2007) The potential effects of pH and buffering capacity on dental erosion. *Gen Dent* 55: 527-531.
22. Talebi M, Saraf A, Ebrahimi M, Mahmodi E (2009) Dental Erosion and Its Risk Factors in 12-year-old School Children in Mashhad. *Shiraz Univ Dent J* 9: 13-18.
23. Dugmore CR, Rock WP (2003) The progression of tooth erosion in a cohort of adolescents of mixed ethnicity. *Int J Clin Pediatr Dent* 13: 295-303.

Author Affiliation

Top

- ¹Department of Orthodontics and Dentofacial Orthopaedics, Indira Gandhi Government Dental College and Hospital, Jammu, India
²Department of Orthodontics and Dentofacial Orthopaedics, India
³Department of Public Health Dentistry, Luxmi Bai Institute of Dental Sciences & Hospital, Patiala, India
⁴Luxmi Bai Institute of Dental Sciences & Hospital, Patiala, India
⁵Genesis Institute of Dental Sciences, Ferozepur, India